



**DISEASES OF THE  
CHEST**

**VOLUME 2**

*AFRICA* BUTTERWORTH & CO (AFRICA) LTD  
DURBAN GOODRICKE'S BUILDINGS MASONIC  
GROVE

*AUSTRALIA* BUTTERWORTH & CO (AUSTRALIA) LTD  
SYDNEY 8 O CONNELL STREET  
MELBOURNE 430 BOURKE STREET  
BRISBANE 240 QUEEN STREET

*CANADA* BUTTERWORTH & CO (CANADA) LTD  
TORONTO 1367 DANTORTH AVENUE

*NEW ZEALAND* BUTTERWORTH & CO (AUSTRALIA) LTD  
WELLINGTON 49/51 BALLANCE STREET  
AUCKLAND 35 HIGH STREET

# DISEASES OF THE CHEST

*Edited by*

SIR GEOFFREY MARSHALL

K C V O , C B E , M D ( L O N D ) , F R C P ( L O N D )

CONSULTING PHYSICIAN, GUY'S HOSPITAL,  
BROMPTON HOSPITAL, AND KING EDWARD VII SANATORIUM MIDHURST, SUSSEX

*and*

KENNETH M. A PERRY

M A , M D ( C A N T A B ) , F R C P ( L O N D )

ASSISTANT PHYSICIAN THE LONDON HOSPITAL,  
VISITING PHYSICIAN, PAPWORTH VILLAGE SETTLEMENT, CAMBRIDGE

*(In two volumes)*

BUTTERWORTH & CO. (PUBLISHERS) LTD.

BELL YARD, TEMPLE BAR

1952



# BUTTERWORTHS MEDICAL PUBLICATIONS

## "MODERN TRENDS" SERIES

- DERMATOLOGY—Edited by R M B MacKenna, MD, FRCP  
 DIAGNOSTIC RADIOLOGY—Edited by J W McLaren, MA, MRCS, LRCP,  
 DMRE  
 GASTRO ENTEROLOGY—Edited by F Avery Jones, MD, FRCP  
 NEUROLOGY—Edited by Anthony Feiling, MD, FRCP  
 OBSTETRICS AND GYNAECOLOGY—Edited by K Bowes, MD, MS, FRCS  
 OPHTHALMOLOGY, Second Series—Edited by Arnold Sorsby, MD, FRCS  
 ORTHOPAEDICS—Edited by Sir Harry Platt, MD, MS, FRCS  
 PAEDIATRICS—Edited by the late Sir Leonard G Parsons, FRCS, MD, FRCP  
 PSYCHOLOGICAL MEDICINE—Edited by Noel G Harris, MD, FRCP  
 PUBLIC HEALTH—Edited by Arthur Massey, CBE, MA, MD, DPH

## "MODERN PRACTICE" SERIES

- ANAESTHESIA—Edited by Frankis T Evans, MB, BS, DA  
 DERMATOLOGY—Edited by G B Mitchell-Heggs, OBE, MD, FRCP  
 INFECTIOUS FEVERS (2 Vols)—Edited by H Stanley Banks, MD, FRCP, DPH  
 OPHTHALMOLOGY—Edited by H B Stallard, MBE, MD, FRCS  
 PSYCHOLOGICAL MEDICINE—Edited by J R Rees, CBE, MD, BCh, FRCP  
 TUBERCULOSIS (2 Vols)—Edited by T Holmes Sellors, DM, MCh, FRCS,  
 and J L Livingstone, MD, FRCP

## TEXT BOOKS

- ANKYLOSING SPONDYLITIS—The late F Hernaman Johnson, MD, FFR,  
 DMRE, and W Alexander Law, OBE, MD, FRCS  
 RADIOLOGY—William Evans, DSc, MD, FRCP  
 CARDIOGRAPHY—William Evans, DSc, MD, FRCP  
 CHRONIC BRONCHITIS—Trevor H Powell, MRCP (Ed)  
 CARDIOSCOPY—William Evans, DSc, MD, FRCP (*In preparation*)  
 DISEASES OF THE EAR, NOSE AND THROAT—Edited by W G Scott-Brown, CVO,  
 MD, BCh, FRCS  
 FOETAL AND NEONATAL PATHOLOGY—J Edgar Morison, MD, BSc  
 GENETICS IN OPHTHALMOLOGY—Arnold Sorsby, MD, FRCS  
 MANAGEMENT OF BRONCHIAL ASTHMA—Herbert G J Herxheimer, MD  
 (Frankfort), LRCP (Ed), LRCS (Ed), LRFPS (Glas)  
 MEDICAL TREATMENT PRINCIPLES AND THEIR APPLICATION—Edited by the late  
 Geoffrey Evans, MD, FRCP  
 PATHOLOGY OF TUMOURS—R A Willis, DSc, MD, FRCP  
 POSTGRADUATE OBSTETRICS AND GYNAECOLOGY—F J Browne, MD (Aberd),  
 DSc, FRCS (Ed), FRCOG  
 PRINCIPLES OF PATHOLOGY—R A Willis, DSc, MD, FRCP  
 SYSTEMIC OPHTHALMOLOGY—Edited by Arnold Sorsby, MD, FRCS  
 UROLOGY OF CHILDHOOD—T Twistington Higgins, OBE, MB, ChB, FRCS,  
 D Innes Williams, MD, MChir, FRCS, and D F Ellison Nash,  
 FRCS  
 VARICOSE VEINS—R Rowden Foote, MRCS, LRCP, DObst RCOG  
 BRITISH DENTAL ANNUAL (1952)—Edited by Evelyn Sprawson, MC, DSc,  
 FDS, MRCS, LRCP

# TABLE OF CONTENTS

## VOLUME 1

CHAPTER	PAGE
LIST OF CONTRIBUTORS - - - - -	viii
PREFACE - - - - -	xi
1 BRONCHO PULMONARY ANATOMY - - - - - A F Foster-Carter	1
2 APPLIED PHYSIOLOGY OF RESPIRATION - - - - - F J Prime and J McMichael	23
3 ACUTE TRACHEO-BRONCHITIS - - - - - J G Scadding	43
4 CHRONIC BRONCHITIS - - - - - J G Scadding	55
5 THE PNEUMONIAS - - - - - J G Scadding	60
6 CHEMICAL, LIPOID AND IRRADIATION PNEUMONIAS - - - - - A Thelwall Jones	141
7 WHOOPING-COUGH - - - - - Kenneth M A Perry	154
8 FUNGUS INFECTIONS OF THE LUNGS - - - - - R W Riddell	162
9 LUNG ABSCESS - - - - - W P Cleland	197
10 EPIDEMIOLOGY AND IMMUNOLOGY OF PULMONARY TUBERCULOSIS - - - - - The Late S Roodhouse Gloyne	245
11 PULMONARY TUBERCULOSIS IN CHILDHOOD - - - - - A Margaret Macpherson	276
12 HAEMATOGENOUS TUBERCULOSIS OF THE LUNGS - - - - - N Lloyd Rusby	294
13 BRONCHIOGENIC TUBERCULOSIS - - - - - F H Young	314
14 PLEURAL TUBERCULOSIS - - - - - F H Young	355
15 CLINICAL PATHOLOGY OF PULMONARY TUBERCULOSIS - - - - - J W Clegg	365
16 TREATMENT OF PULMONARY TUBERCULOSIS - - - - - Andrew Morland and Sir Clement Price Thomas	376

# TABLE OF CONTENTS

CHAPTER		PAGE
17	THE CHEMOTHERAPY OF PULMONARY TUBERCULOSIS - - - John Crofton	431

## INDEX TO VOLUME 1

## VOLUME 2

1	PLEURISY - - - - - Kenneth Robson	1
2	EMPYEMA - - - - - Oswald Tubbs	18
3	HAEMOTHORAX - - - - - Kenneth M A Perry	49
4	SPONTANEOUS PNEUMOTHORAX - - - - - Kenneth M A Perry	54
5	CALCIFICATION OF THE PLEURA - - - - - I C Lodge Patch and Kenneth M A Perry	60
6	TRANSIENT PULMONARY INFILTRATION WITH EOSINOPHILIA - - - James L Livingstone	65 ✓
7	POLYARTERITIS NODOSA (PERIARTERITIS NODOSA) - - - James L Livingstone	74
8	ASTHMA - - - - - E R Bofand	82
9	EMPHYSEMA OF THE LUNGS - - - - - Ronald V Christie	101
10	PULMONARY OEDEMA - - - - - G W Hayward and Ronald V Christie	123
11	ATELECTASIS - - - - - F P Lee Lander	137
12	BRONCHIECTASIS - - - - - F P Lee Lander	156
13	INTRA-THORACIC MANIFESTATIONS OF SARCOIDOSIS - - - J G Scadding	185
14	THE RETICULOSES AND ALLIED DISORDERS - - - - - Neville Oswald	192
15	HONEYCOMB LUNGS - - - - - Neville Oswald	200
16	CARDIO-PULMONARY SCHISTOSOMIASIS - - - - - D Evan Bedford	216
17	DISEASES OF THE LUNG CAUSED BY FLUKES, ROUND-WORMS AND AMOEBAE - - - - - Kenneth M A Perry	223

# TABLE OF CONTENTS

CHAPTER							PAGE
18	HYDATID DISEASE OF THE LUNG	-	-	-	-	-	229
	Geoffrey Flavell						
19	SYPHILIS OF THE LUNG	-	-	-	-	-	235
	Kenneth M A Perry						
20	PULMONARY EMBOLISM	-	-	-	-	-	238
	W D W Brooks						
21	BLAST INJURIES OF THE LUNG	-	-	-	-	-	246
	Kenneth M A Perry						
22	PULMONARY HAEMOSIDEROSIS	-	-	-	-	-	248
	Kenneth M A Perry						
23	OCCUPATIONAL LUNG DISEASES	-	-			-	252
	Kenneth M A Perry						
✓ 24	CYSTS OF THE LUNG	-	-	-	-	-	294
	Geoffrey Flavell						
<u>25</u>	TUMOURS OF THE LUNG	-		-	-	-	306
	Vernon C Thompson						
26	BORNHOLM DISEASE	-	-	-	-	-	342
	Kenneth M A Perry						
27	TUMOURS OF THE CHEST WALL	-	-	-	-	-	344
	Geoffrey Flavell						
28	DISEASES OF THE MEDIASTINUM	-	-	-	-	-	350
	Kenneth M A Perry						
29	CYSTS AND TUMOURS OF THE MEDIASTINUM			-	-	-	358
	Geoffrey Flavell						
30	DISEASES OF THE DIAPHRAGM	-	-	-	-	-	375
	Kenneth M A Perry						
31	ANAESTHESIA IN CHEST DISEASE	-	-	-	-	-	388
	A I Parry Brown						
32	PHYSIOTHERAPY FOR CHEST DISEASES	-	-	-	-	-	395
	Jocelyn M W Reed						

## INDEX TO BOTH VOLUMES



## CHAPTER 1

### PLEURISY

KENNETH ROBSON

#### AETIOLOGY

PLEURISY implies inflammation of the pleural surfaces, whether (1) fibrinous, (2) serous with effusion or (3) purulent. The most important single cause of pleurisy, whether acute or chronic, dry or with effusion, is tuberculosis. Pneumonic consolidation—whatever the causative organism—single or multiple pulmonary abscess and primary or secondary neoplasia may also be accompanied by pleurisy, as may complicated bronchiectasis. Rarely, pleurisy complicates acute rheumatism. It may occur in nephritis, often terminally, or during the course of blood diseases such as leukaemia. Diaphragmatic pleurisy, with shoulder-tip pain, and mediastinal pleurisy—both localizations with or without effusion—have been described, as isolated phenomena they are rare.

It is probable that many of the cases of so-called simple or primary pleurisy are, in reality, due to underlying foci of inflammation in the lung, whilst others are tuberculous. The latter aetiology should always be suspected if the pain or rub lasts more than a few days. It is very doubtful whether pleurisy ever follows mere exposure to cold—"Pleuritis a frigore" of the older writers. Epidemics of primary pleurisy have been described (Scadding, 1946) and are probably to be distinguished from Bornholm disease. Pleurisy, evidenced by friction, occurs with varying frequency in the latter, and there is still doubt whether the primary site of inflammation is in the pleura or in the muscles and fasciae of the parietes. It is the epidemic, sometimes explosive, nature of an outbreak which stamps it as something distinct. In most outbreaks of Bornholm disease, relatively few of the individuals who have the severe, sudden chest pain actually show friction.

#### CLINICAL PICTURE

##### Pleurisy without effusion

Dry pleurisy whatever its cause, is a painful condition characterized by knife-like pain related to the respiratory excursion, which it tends to inhibit locally. The pain is not of critically sudden onset, and careful interrogation will generally distinguish it from intercostal neuralgia, fibrositis and pre-eruptive herpes. Friction is the cardinal sign and must always be sought. The friction has a rubbing, grating or scratching quality and, if coarse, can be palpated with the flat of the hand. Pressure with the hand will often cause the pain to disappear, and pressure with the chest-piece of the stethoscope may diminish, or even abolish, the friction sound while pressure is maintained. Friction in a peculiar way sounds very "near to the ear". It is often useful to posture the patient so as to stretch the side of the thorax on which the rub is being sought, and to listen exactly over the site of the pain as indicated by the patient. With pleural friction, as with other physical

signs in the chest, its absence in a given case by no means excludes the possibility of pleurisy

Radiographic evidence of dry pleurisy is not constant and fluoroscopy only sometimes affords evidence in the form of abnormalities of diaphragmatic function and behaviour. X-ray examination, however, will eliminate such conditions as spontaneous pneumothorax, and will help to establish the presence of any intrapulmonary condition of which pleurisy is symptomatic

### **Pleurisy with effusion**

Effusion may develop insidiously or it may follow or be accompanied by pleuritic pain. Fluid, in considerable quantities, can accumulate without symptoms if it collects slowly and the patient is thus given time to adjust to its presence gradually. On the other hand, it may sometimes accumulate with great rapidity, in which case distress and dyspnoea may become urgent.

The physical signs of fluid, whether it be transudate or exudate, pus, chyle or blood, are characteristic and, provided that the quantity is sufficient, unmistakable. The apex beat is displaced away from the fluid, movement on the affected side is reduced, and there is decrease in tactile fremitus and impairment of percussion note on the side of the fluid. The impairment of percussion note is so striking that the word "stony" has come to be associated with this type of dullness, and eliciting the physical sign may often cause pain in the fingers of the percussor. The breath sounds are absent or altered in strength and may, in certain circumstances, be bronchial. Added sounds are no part of the picture unless there be a rub at the upper limit of the fluid. The spoken voice often takes on that peculiar nasal quality known as "aegophony". Physical signs are found in the presence of moderate or large collections of fluid, but if the fluid is confined to an interlobar fissure or is less than 150 millilitres approximately in total amount in an adult, it may be unrecognizable by physical examination.

Fluoroscopy will show gross collections of fluid immediately. The affected part of the thorax is poorly lit and the illumination does not improve with inspiration. Turning the patient slowly into the lateral and oblique positions may help to define the extent of the fluid, and the lordotic view will generally show involvement of the greater fissure. The x-ray plate provides further evidence and a permanent record. Fluid gives a shadow as opaque as the heart shadow and diaphragm, with which, according to the amount which is present, it merges. A well penetrated view may show its distribution but, despite the older teaching on the subject, the upper limit of the fluid does not invariably run in a gentle curve towards the axilla, and change of posture only slowly brings about re-distribution in the closed pleural cavity, too slowly to permit regular and convincing demonstration of shifting dullness (Figs 1-3, 7a and b).

### **TYPES OF PLEURISY WITH EFFUSION**

The following are the important groups of causes of a pleural effusion.

- (1) tuberculous effusion,
- (2) effusion complicating disease in the subjacent lung, other than tuberculosis, of all gradations up to frank empyema,

# TYPES OF PLEURISY WITH EFFUSION

FIG 1 — Total left-sided effusion (By courtesy of the Institute of Diseases of the Chest, Brompton)



FIG 2 — Left sided lateral effusion (By courtesy of the Institute of Diseases of the Chest Brompton)



# PLEURISY




FIG 3 — (a) Postero-anterior and (b) lateral views of interlobar effusion on the right side showing opening up of the greater and lesser fissures (By courtesy of the Institute of Diseases of the Chest, Brompton)

a)



(b)

## TUBERCULOUS EFFUSION—TUBERCULOUS PLEURISY

- (3) effusion complicating sub diaphragmatic or hepatic disease
- (4) effusion in cardiac and renal disease ,
- (5) effusion accompanying new growths ,
- (6) chylous and pseudo chylous effusions

## TUBERCULOUS EFFUSION—TUBERCULOUS PLEURISY

To Laennec belongs the credit for the demonstration of the relationship between pleurisy and tubercles in the lung Either dry pleurisy or pleurisy with effusion




FIG 4 —Graph showing age incidence of primary serous pleurisy (idiopathic)  
(From Seel and Foren 1928 )

can occur at any time during the course of pulmonary tuberculosis, and the underlying cause presents no difficulty when established disease with tubercle bacilli in the sputum comes to light in investigation Tuberculous dry pleurisy and more particularly, pleural effusion also occur in the absence of any readily obtained confirmation and in this group the danger of misdiagnosis is very great

### Types of tuberculous pleurisy

There are two main varieties of tuberculous effusion

- (1) occurring as a late primary manifestation ,
- (2) due to the presence sub-pleurally of post primary foci

Both groups sometimes continue to a tuberculous empyema The second occurs perhaps more usually in middle aged or elderly subjects and is generally very chronic

*A late primary manifestation*  
 Occurring as a late manifestation of the primary infection, tuberculous effusion has its greatest incidence between the ages of 15 and 25 years (Fig 4)

*Pleurisy as an allergic phenomenon*—The view that the outpouring of fluid represents an allergic phenomenon is maintained by Wallgren (1930) and Rich (1946). An individual who experienced his primary infection with tubercle bacilli, its development, the irritation of the pleural onchial adenitis with the discharge of bacilli may also be the source of hypersensitivity developed varies with the constitution of the individual patient. The effusion may be of sudden onset or it may develop insidiously. It may be associated with acute symptoms or it may be a casual discovery. The latter fact suggests that certain individuals pass through an effusion of limited extent without its ever coming to clinical recognition. Proof of this is difficult and an exact expression of its frequency equally so. Some idea may be obtained from figures supplied by Brooks. Of some 420,000 individuals between the ages of 15 and 75 years examined in the course of a fluorographic survey, 56 cases showed radiographic evidence of recent primary tuberculosis with a symptomless pleural effusion. The opportunity arose to observe 63 out of 259 individuals who showed radiographic evidence of recent primary tuberculosis, and in 6 cases a pleural effusion developed within 6 months.

Thompson (1946) found that the effusion tended to form on the same side as the pulmonary or glandular lesion. There is general agreement that the tuberculous primary complex is right-sided more often than left-sided (Hayne, Pagel and O'Shaughnessy, 1947) and in most cases the initial effusion is on the right side.

#### *Post-primary effusion*

Either dry pleurisy or pleurisy with effusion may occur at any time during the life-time of a consumptive patient. Dry pleurisy is often evanescent and marks advance or extension of disease. Effusion may develop and, when it does so, the accumulation of fluid is often gradual and insidious. The older the patient, the more likely is the effusion to be loculated, the more gradual the accumulation, the slower is the absorption so that it is typical of tuberculous effusion in middle-aged subjects to pursue a chronic course, which may, however, eventually have a satisfactory outcome. Nevertheless it carries a higher mortality correspondingly than does the juvenile variety. The constitutional disturbance may be less severe and the fluid more often blood stained than in the juvenile type. The cytology and cultural characteristics are otherwise essentially similar in the two varieties. The age incidence of the post-primary effusion, as might be expected, shows a broader base than does the primary type, being similar, in fact, to the age incidence in pulmonary tuberculosis as a whole.

#### *Clinical course*

Perhaps most typically the primary effusion proclaims its presence with malaise and discomfort in breathing. The course of the illness is generally favourable

The temperature subsides in the space of 2 weeks or so the fluid is absorbed function returns and the patient recovers. Death in the acute phase of the effusion is rare so that opportunities for post mortem examination are not frequent. In some of the subjects which have been examined tubercles have been found on the pleural surface and sometimes there has been evidence of dissemination to other organs.

### Complications

These can be considered in two groups (1) those which are complications of an active tuberculous process and (2) those which are complications of the presence of fluid in the chest.

#### *Complications of the tuberculous process*

The complications of the tuberculous process are chronicity and dissemination. These are of course interconnected and the chronicity of an effusion and the serial involvement of the opposite pleural space often go hand in hand. Dissemination may be on a limited scale or it may be widespread resulting in miliary tuberculosis with the picture dominated by meningitis. Other sites to which dissemination commonly takes place are the pericardium the peritoneum and the genito urinary system. In the unstable and febrile phase of an effusion dissemination and acute miliary spread are ever present dangers.

Concato's disease—polyorrrhomenitis or polyserositis—in which there is effusion in both pleurae the peritoneum and the pericardium is nearly always of tuberculous character. Rare cases are met with in which the condition remains quite unexplained.

#### *Complications of the effusion*

The fact of the presence of fluid carries with it the risk of secondary infection either endogenous in intercurrent disease or exogenous. Pleural thickening and adhesions are not uncommon sequels of uncomplicated effusion and sometimes marked pleural calcification develops in the course of time. Mechanical hydrostatic disturbances may result from the fact of the presence of fluid. As is mentioned elsewhere redistribution of the fluid by adjustment due to movement within the thorax may take place in the natural course of events or during aspiration. If the redistribution is rapid changes may ensue in the aeration of large districts of the sound lung or in the pulmonary circulation with resulting oedema of the lungs. The gravity of either of these events or of a combination of them cannot be over emphasized.

### Prognosis

Any individual with a hemithorax half filled or more with fluid is in danger on this account alone. It is true that the danger is not great but the potentialities constitute a definite threat to life. The more immediate prognosis in general terms has been well summarized (Wood 1946) as depending not so much on the extent of the infection as on the unpredictable liability at any stage to acute miliary dissemination.

The more remote prognosis has always been a matter of interest and speculation. It is said above that a patient recovers. But he is a marked man and whether

# PLEURISY

the illness has been short and sharp or long and smouldering, whether there has been total effusion or a small one, whether serial involvement of one pleural space after another has occurred with recurrences of fever or not, patients who have been through an effusion of this sort have a 25 per cent chance of developing clinical pulmonary tuberculosis within 5 years. It was the inescapable fact, of the toll in tuberculous morbidity following pleural effusion, which gradually compelled the adoption of the view that all primary pleurisy is of tuberculous origin unless clearly proved otherwise.

Much laborious work has been put in at follow-up inquiries, but there are few series of cases which are free from some sort of objection. It is important that a

TABLE I  
FOLLOW-UP STUDIES OF TUBERCULOUS PLEURISY

Authors	Percentage of total which are untraced	Size of actual series (cases)	Method of diagnosis	Use of x ray		Observation times		Follow up	Morbidity from tuberculosis
				During attack	Follow up	Minimum	Maximum		
Hedges (1900)	59	130	Needling (67 cases) Phys. signs (63 cases)	No	No	Years 2	Years 8	Personal examination	43%
Allard and Koster (1912)	43.2	514	Not stated	No	No	2	28	Mainly personal examination	47.7%
Scheel and Foien (1928)	15	812	Not stated	No	No	1	10	Notification	22.4%
Gaarde (1930)	Not stated	126	Not stated	Partial use		1	12	Partial personal	40%
Borelius (1933)	15	194	No clear information	Not available	Yes	7	17	Personal examination mainly	39.8% 66.3%
Kallner (1937)	15	580	X ray (76%) Clinical (24%)	Yes (76%)	Yes	6	20	Personal examination	39%
Brompton Hospital (1939)	30	111	Needling (76 cases) Clinical and x ray (35 cases)	Yes	Yes	5	12	Personal examination	23% (42%)
Thompson (1946)	Single case (?)	190	Needling (125 cases) Probably all x rayed	Probable	All except 19 cases	1	8	Majority personal	25%
Sinding-Larsen and others (1947)	20.2	474	Not stated	Not stated	Not stated	5	19	Notification	Pulmonary 6.4% All forms 12.4%
Finkler (1947)	33	548	X-rays. Some by needling	Not stated	Not stated	5	25	Notification	18% 25%
Prest (1947)	Not known	1192	X ray	Yes	Not stated	Not stated		Observation	30%
Maclean (1948)	16.5	183	Not stated	Not stated	Not stated	5	5	Notification	20.5%

series should comprise as large a number of patients as possible and that the number untraced should be minimal. Strict criteria are necessary in diagnosis at the time of the effusion (in order to eliminate cases of pulmonary tuberculosis of outspoken type), and the follow-up examinations should be as comprehensive as possible. The period of observation should be at least 5 years.

The accompanying table summarizes some of the more important facts to be obtained from certain of the published follow up inquiries. Each individual series comprises more than 100 cases.

### **Sampling the fluid**

When fluid is present in amounts sufficient to give rise to physical signs, it is always wise to take a sample, this is done to confirm the presence of fluid in a doubtful case and to proceed from the recognition of its presence to an attempt to discover why it is there. Sampling should always be carried out with full aseptic precautions. The best site for exploratory puncture (or thoracentesis) is decided after careful scrutiny of postero-anterior and lateral films, and the puncture is made with a needle mounted on a syringe after infiltration with a local anaesthetic right down to the pleura. In an average case the seventh or eighth interspace in the mid axillary line is a suitable site. The patient may either be sitting up and leaning slightly forwards with the arms resting on a bed-table, or lying on his side with the site of the effusion uppermost.

The fluid is generally straw-coloured (very rarely, if trauma be excluded, haemorrhagic) and clear. Sometimes a little clot may form on standing. Lymphocytes are abundant and generally predominate, although occasionally even tuberculous fluid is distinctly polymorphonuclear in its cellular content in the earlier phases (Wihman, 1948, Kraft, 1949). Usually no organism can be seen and culture is sterile on all ordinary media. Inoculation into a guinea pig may produce tuberculous lesions in up to 50 per cent of cases, and culture on Löwenstein Jensen's (or Dubos') medium may be positive for tubercle. Increasing use will probably be made of cultural methods as techniques are improved and standardized.

Close (1946) gives some interesting figures concerning the relative merits of guinea pig inoculation and culture in the detection of tubercle bacilli in pleural fluid, and also of the increasing number of positive results which come with experience. Out of 34 tubercle positive fluids, 10 were positive to both guinea pig inoculation and culture, and 24 were positive to culture but negative to guinea-pig inoculation. Of 11 attempts to demonstrate tubercle bacilli in fluid, made in 1941, none was successful, in 1945, however, 16 were positive out of 23. There is no doubt that the technique employed is of the greatest importance. More recently (Gelenger and Wiggers, 1949, Calnan and his colleagues, 1951) the glucose content of the fluid has been investigated. There is some doubt about the level of glucose which is critical, but low values strongly suggest a tuberculous origin whilst high levels are found in non-tuberculous conditions.

### **Treatment**

#### *General management*

With tuberculous pleurisy this is a long-term matter. In the absence of any reliable specific remedy, the indications are to support the patient by every means

at our disposal while his own recuperative powers and specific resistance develop. Rest is the "sheet anchor"—general bodily rest during the febrile phase and stiling of the lung and pleural movement until absorption has taken place. Strict rest can be a most valuable measure if the temperature is obstinate in settling, and it is essential in total effusion. Patients with large effusions are best nursed propped up and must be very well supported with pillows. Just as too rapid decompression of the thorax in these cases can lead to fatal consequences, so can sudden redistribution within the thorax of a large effusion. Fidgeting and restlessness can initiate such a redistribution. This state of affairs sometimes arises very critically, the aspect of the patient changing in a moment from one of relative comfort and calm to one of alarm, apprehension and respiratory embarrassment. He becomes collapsed, cold and clammy, with a rapid pulse and shallow respiration, and it is very difficult to rescue him. Oxygen and restoratives should be used at once and aspiration should be carried out. (*See also under "Complications" and "Management of the fluid"*)

The temperature takes a variable time to subside, sometimes up to 6 weeks. From 10 to 14 days after it has become normal provided that absorption is proceeding satisfactorily and there are no complications, a patient may start getting up. By the end of a further 6 weeks—that is approximately 3 months from the beginning of the illness—he will have arrived at the stage of being up for 4 hours a day and doing incidental walking of, perhaps, half a mile a day. *Graduated walking exercise is now instituted with increased time up, and by the end of a further 3 months he will be up all day and walking 5 miles per day.* In most cases it will be permissible at this stage to let him resume work, but supervision and serial x-ray observation must continue for 5 years.

A good working rule is to examine clinically and radiographically patients who have had an effusion at the following intervals: every 3 months for a year, every 4 months for the second year, every 6 months up to the end of 5 years. In the clinical examination, special importance attaches to the weight, and to the possibility that metastatic tuberculous deposits have made their appearance in the interim.

Attempts should not be made during convalescence to promote absorption of the fluid or adhesions by expansion breathing exercises or to "unfreeze" a lung (unless after some months of constitutional stability). The aim is a compromise—to obtain settlement of an inflammatory process, which carries with it always the hazard of dissemination. Within limits, the anatomical result *per se* is unimportant, and the existence of a poorly expanding hemithorax with movements reduced to a minimum may have to be accepted, at least for a time. What should be emphasized is the functional result, as seen by a patient's ability to get about and earn his own living or continue his training. Any decision about

### *Management of the fluid*

A sample should always be taken, but it is less easy to lay down rules about

removal of fluid therapeutically. The general principle is that so far as possible, the fluid shall be allowed to become absorbed of its own accord. Over enthusiasm in aspiration is more likely to do harm than would conservatism even allowing for those patients whose breathing becomes urgently and critically embarrassed. There are three indications to interfere

- (a) in fever lasting a disproportionately long time
- (b) in fluid persisting a disproportionately long time
- (c) to relieve respiratory distress

It will often be found that removal of up to 1 pint of fluid will suffice to tip the scale towards stabilization or absorption in the case of (a) or (b). The method of removal is immaterial provided that the process is slow, is conducted under sterile precautions and does not involve the introduction of air into the free pleural space. This point of view is maintained although the policy of complete evacuation of the fluid in successive stages and pneumothorax induction is not without its supporters (Feldman and Lewis 1946). For the aspiration a large capacity syringe fitted with a two way tap is convenient. Local anaesthesia must be used down to the pleura and a stout needle preferably incorporating some means of clearing with a stylette is the most serviceable. Tubing other than stout walled pressure tubing often collapses under suction and aspiration becomes impossible. Alternatively Potain's aspirator can be used.

When aspiration is undertaken to relieve dyspnoea it is generally because fluid has accumulated rapidly and the circumstances which have initiated the rapid accumulation will in all probability operate again. It is unlikely that the removal of a pint or two from a space containing 10 or 12 pints total effusion will contribute much to the comfort of the patient and theoretically the removal of a large amount will be necessary. To remove amounts in excess of 3 pints presents technical difficulties.

Rapid removal is dangerous involving as it does the risk of disturbing the pulmonary circulation and producing oedema of the lung with albuminous expectoration and death. Warning of this development is given by the patients experiencing discomfort and oppression in the chest and by the appearance of

in position connected to an underwater drain or simply to a bottle on the floor by the bedside is very apt to be followed by sepsis and this is a very serious complication indeed. Some authors advocate a really big volume aspiration with air replacement compensating for the removal of the fluid by the introduction of air. Theoretically this is a sound method but practically it cannot always be carried out and it should certainly not be attempted by the inexperienced. Another plan (perhaps the method of choice) is to remove 20-30 ounces daily for 4-6 days although with this method one is very apt to encounter a dry tap at the third or fourth attempt and to fail of one's purpose.

Among prophylactic measures are limitation of fluid intake (allowing 30-50 fluid ounces in 24 hours in all) and the administration of diuretics by the mouth



(Diuretin, 10 grains thrice daily) or of one of the mercurial preparations by intramuscular injection. There are instances in which these measures appear to have been of use.

#### *Other methods of treatment*

The institution of artificial pneumothorax by air replacement of the fluid is mentioned above. The field of application of this technique is, however, probably very limited. Pneumoperitoneum with phrenic interruption has its advocates, but insufficient time has as yet elapsed for assessment of its worth as compared with the ordinary conservative regimen.

Although there is at present insufficient experience on which to base a final judgment—and here again insufficient time has as yet elapsed for an assessment to be made—there is good reason to think that streptomycin has a definite place in the management of large effusions or those associated with prolonged fever. Certainly the immediate benefit is striking. The dose for an adult is 1 gramme daily intramuscularly in a single dose. The antibiotic is given alone for 7–10 days. As soon as the general condition of the patient permits, para-aminosalicylic acid (15 grammes *per diem* in divided doses) is added. After a loading dose of 30 grammes of streptomycin in daily injections, the frequency of the injections can be reduced to 3 times a week, perhaps some 45 grammes of streptomycin in all being used.

### POST PNEUMONIC EFFUSION

This title embraces effusions complicating disease in the subjacent lung, other than tuberculosis, of all gradations up to frank empyema (post-pneumonic effusion).

#### *Aetiology*

Empyema thoracis most commonly occurs as a complication of pneumonia, generally pneumococcal, although any organism may be involved, including *Bacterium coli* and the streptothrix. It may also be associated with other infective conditions of the lung, particularly (i) pulmonary abscess, (ii) bronchiectasis or (iii) infected lung and bronchial tissue distal to an obstructing neoplasm. What is true of empyema can be regarded as true also of post-pneumonic effusion, which is, in a sense, an attenuated form of empyema. The increasing and early use of all forms of chemotherapy in pneumonic illness has led to considerable modifications in the form of pneumonia and to a probable reduction in the over all occurrence of empyema.

#### *Differential diagnosis*

It is of the greatest importance to distinguish between a post-pneumonic collection of fluid and one due to tuberculosis. The whole management and handling of the case is different, but the distinction is sometimes a matter of some difficulty. All too frequently a case only comes under close scrutiny after the fluid has formed, perhaps about the tenth day of the illness. It will be known that there have been some days of illness, but in all likelihood no information will be available about the leucocytosis and no systematic observation of temperature may have been carried out. Some of the differentiating points are now formulated.

## EFFUSION COMPLICATING SUB DIAPHRAGMATIC OR HEPATIC DISEASE

### *Tuberculous effusion*

Patient commonly 10-25 years of age  
 Pain often early  
 High fever, not responding to sulphonamides or penicillin  
 No leucocytosis  
 Fluid obliteration of space on x-ray film from the start  
 Fluid lymphocytic  
 No organisms seen in the smear, except rarely tubercle bacilli  
 Culture on ordinary media sterile  
 Special culture and guinea-pig inoculation may be positive for tubercle  
 Prolonged course with slow absorption

### *Post pneumonic effusion (attenuated empyema)*

Any age or sex  
 Pain not an essential feature, especially early  
 May be high fever responding to chemotherapy  
 Leucocytosis early  
 Clear pleural space early on skiagram  
 Fluid *may* be polymorphonuclear  
 Coccal organisms *may* be seen  
 Culture on ordinary media *may* be positive for streptococci or pneumococci  
 Negative for tubercle  
 Often absorbed within 3-4 weeks

### Treatment

The treatment of post-pneumonic effusion, if the disease is of moderate extent, consists in the evacuation of the fluid by aspiration. Small collections of fluid may be allowed to be absorbed spontaneously. Observation with radiography or under the fluoroscope is necessary to verify that absorption is taking place, and the final check should be made one month after absorption appears to be complete, before the patient is released from observation. If the fluid is considerable in amount, it is a good plan, after aspiration, to introduce penicillin intramuscularly. If there has been time to do sensitivity tests on the organisms grown from a culture of pleural fluid, then a different chemotherapeutic agent may have been found appropriate and this should be used. Penicillin, up to 1,000,000 units, should be injected after aspiration, and as soon as possible expansion exercises should be initiated. This active rehabilitation is in sharp contrast with what is practised in tuberculous effusion, but it is of the greatest importance, since re-expansion of the underlying lung is the only satisfactory way to obliterate what is potentially an empyema cavity.

## EFFUSION COMPLICATING SUB DIAPHRAGMATIC OR HEPATIC DISEASE

Pleural effusion occurs as part of the general dropsical process when cirrhosis of the liver has reached an advanced stage. Occasionally the effusion may be the

## PLEURISY

primary tuberculous pleurisy, but sometimes the reaction is strongly polymorphonuclear. A careful history may be of the greatest use, supplying evidence of a severe upper abdominal catastrophe within recent weeks. Many years of good health may have intervened between the original symptoms of amoebiasis and the development of hepatitis. The induction of an artificial pneumoperitoneum may assist in differentiation, since those conditions which are associated with a sympathetic effusion generally bring about adhesion between the liver and the right side of the diaphragm.

## EFFUSION IN CARDIAC AND RENAL DISEASE

The collection of fluid in the pleural space in cardiac and renal disease is referred to as hydrothorax—a term which serves to differentiate the condition from that in which the fluid results from inflammatory lesions of the pleura. In cardiac and renal conditions, the fluid is part of the general dropsical process and the treatment is that of the underlying state. In renal disease, subacute nephritis with oedema, both pleural spaces are usually involved, but in cardiac disease, congestive heart failure, the right side may be affected alone. This may be related to pressure on the azygos vein by the enlarged right auricle. Aspiration may be required, particularly in heart failure. The fluid in hydrothorax has a low specific gravity (generally less than 1.015), and only small amounts of protein and a few cells are present, in contrast with the high protein content and cellularity of inflammatory effusions.

## EFFUSION ACCOMPANYING NEW GROWTH

New growths in the thorax, whether primary or secondary, innocent or malignant, may be accompanied by pleural effusion. Effusion is usually not an important part of the picture in innocent growths of the lung or chest wall, but it may be the presenting feature in primary malignant pulmonary disease or the first evidence of intrathoracic metastasis in breast cancer. The poorly differentiated form of bronchial carcinoma, the so called oat cell type, is more likely to present in this way than is the squamous variety. The effusion is an index of the proximity of the growth to the pleura or of massive involvement of root and mediastinal glands by the growth, perhaps, also, it is indicative of the poorly formed vascular channels which characterize a rapidly growing undifferentiated neoplasm.

The effusion is often frankly blood stained, even at the very first puncture. This is also true of secondary neoplasia, for instance from carcinoma of the breast, with which the pleural surface is studded. Cells, showing mitotic figures and with other features suggesting that they may have become detached from malignant growths, may be found in the fluid. In order to obtain such evidence it is particularly worth while to examine sections prepared from strands of blood clot and other debris remaining in the lumen of the needle or adhering to its end after aspirations of a few ounces of the fluid.

Effusion develops in connexion with benign or malignant growths of the pleura, whether primary or secondary. Primary malignancies are excessively rare, but cases are on record, some of which are claimed to be carcinoma and others sarcoma.

## CHYLOUS AND OTHER MILKY EFFUSIONS

At the present time it seems fairly certain that some of the cases should be regarded as endothelioma (Buxton and Wilcox 1950). Other tumours of the pleura which may be accompanied by effusion are lipoma and fibroma which are benign whilst in addition to the breast the stomach and oesophagus are sometimes primary sites of growth from which metastasis takes place to the pleura. Medial dermoid neoplasia in which malignant change has taken place, is sometimes accompanied by a pleural effusion. Tinney and Olsen (1945) dealing with 141 examples of pleural effusion due to neoplasms give the distribution of primary sites of origin as shown in Table II

Site of primary growth	No. of cases	Percentage
Intrathoracic	51	36
Bronchial carcinoma	47	—
Medastinal carcinoma	4	—
Breast	42	30
Ovary	12	8
Kidney	6	4
Stomach	5	4
Colon and rectum	4	3
Lower urinary tract	4	3
Oesophagus	3	1
Thyroid	1	1
Jejunum	1	1
Not established	14	9

In Meigs syndrome serous pleural effusion sometimes with ascites is found in association with fibroma of the ovary (Meigs and Cass 1937). Since removal of the fibroma is curative an exploratory laparotomy is indicated even when the case appears to be one of inoperable ovarian malignant disease with pleural metastasis.

## CHYLOUS AND OTHER MILKY EFFUSIONS

Chylothorax is the name applied to an effusion containing chyle the exudate appears as milky fat-containing fluid. About 33 per cent of cases follow trauma (MacNab and Scarlett 1932) and of the remainder malignant neoplasms involving the thoracic duct and its tributaries, are the commonest cause but especially in the tropics filariasis is also a cause. The diagnosis is made from the milky appearance of the pleural fluid its only nature being established by staining with Sudan III. Olsen and Wilson (1944)

## PLEURISY

suggested that the condition was more common than is usually thought. In certain cases of reticulosis, tuberculosis and carcinoma in which pleural effusions develop, the fluid may have a milky appearance and contain fat, although it does not contain chyle, such effusions have been described as chyliform. Pseudo chylous effusions also occur in which the milky appearance is not due to fat but to other particles causing opalescence, such as a lecithin globulin complex, calcium phosphate, cholesterol or filarial embryos. These effusions occur rarely in the reticuloses, in tuberculosis and in neoplastic conditions, and sometimes also in heart or kidney disease.

The prognosis in all these cases is grave, the seriousness depending on the underlying condition. The best results may be expected in the less severe traumatic cases. The effusions tend to recur and, in the case of true chylous effusions, aspiration is not advisable except for diagnostic purposes, because the loss of fat, which results, may be serious.

## BIBLIOGRAPHY

- Allard H and Koster H (1912) "Idiopathic Pleurisy and Pulmonary Tuberculosis" *Brit med J* 1, eptome 17
- Borelius R (1933) On the Question of Pleurisy and Pulmonary Tuberculosis After Examination *Acta med scand* 79, 391
- Brooks W D W (1951) Personal communication *Brit med J* 1, 281
- Buxton P H and Wilcox A (1950) Endothelioma of the Pleura *Brit med J* 1, 281
- Calnan W L (1951) Diagnostic Value of the Glucose Content of Serous Pleural Effusions *Brit med J* 1, 1239
- Close H G (1946) Tubercle Bacilli in the Pleural Effusion of Young Adults *Lancet* 1, 193
- Erwin G S (1944) Pleural Effusions Arising from Tuberculous Tracheo-bronchial Adenitis (Primary Effusion) *Tubercle* 25, 44
- Feldman D J and Lewis H P (1946) Tuberculous Pleural Effusion *Med clin N Amer*, 30, 245
- Finkler E (1947) Pleuritis Exsudativa und spätere Tuberkulose Erkrankung (statistische Erhebung an 548 Militär patienten) *Schweiz Ztschr Tuberk* 4 372
- Gaard F W (1930) The Prognosis of Pleurisy With Effusion *J Amer med Ass* 95, 249
- Gelenger S M and Wiggers R F (1949) Relationship of the Pleural Fluid Sugar to Pulmonary Tuberculosis *Dis Chest* 15, 325
- Hedges C E (1900) The Aetiology Immediate and Remote Prognosis of Primary Pleurisy with Serous Effusion *St Bart's Hosp Rep* 36, 75
- Kallner S K (1937) On the Prognosis of Exudative Pleurisy *Acta med scand* 92, 549
- Kayne Pagel and O Shaughnessy (1948) *Pulmonary Tuberculosis* London Oxford University Press
- Kraft, K R (1949) The Diagnostic Problem of Primary Pleural Effusions *Amer Rev Tuberc* 59, 259
- Maclean K S (1948) The Relation of Idiopathic Pleurisy and Pleural Effusion to Tuberculosis *Guy's Hosp Rep* 97 133
- MacNab D S and Scarlett E P (1932) Traumatic Chylothorax Due to Intrathoracic Rupture of the Thoracic Duct *Canad med Ass J* 27, 29
- Meigs J V, and Cass J W (1937) Fibroma of the Ovary with Ascites and Hydrothorax *Amer J Obstet Gynec* 33 249
- Olsen A M and Wilson G T (1944) Chylothorax *J thorac Surg* 13, 53
- Press P (1947) Les Pleuresies Exudatives Tuberculeuses Antecedents et Pronostic *Rev med Suisse Romande* 67, 100
- Rich A (1946) *The Pathogenesis of Tuberculosis* 2nd ed Springfield Ill, Thomas
- Robson, K (1944) Modern Views on Primary Pleurisy with Effusion *Practitioner*, 153, 344

# BIBLIOGRAPHY

- Seadding J G (1946) Acute Benign Dry Pleurisy in the Middle East *Lancet* **1**, 763
- Scheel and Foien (1928) Pleuresie et Tuberculose Ulterieure *Acta med scand*  
Suppl No 26 187
- Sinding Larsen C M and Heldaas O (1947) Tuberculous Pleurisy *Nord Med* **34** 1255
- Thompson B C (1946) Pathogenesis of Pleurisy with Effusion *Amer Rev*  
*Tuberc* **54** 349
- Tinney W S and Olsen A M (1945) The Significance of Fluid in the Pleural Space  
A Study of 274 cases *J thorac Surg* **14** 248
- Wallgren A (1930) On the Pathogeny of Acute Sero fibrinous Pleurisy *Acta med scand*  
Suppl No 34 349
- Wihman G (1948) A Contribution to the Knowledge of the Cellular Content in Exudates  
and Transudates *Acta med scand* Suppl No 205
- Wood S R (1948) Serous Tuberculosis in East African Natives *Brit J Tuberc* **42** 38

## CHAPTER 2

### EMPHYEMA

OSWALD TUBBS

#### NON-TUBERCULOUS EMPHYEMA THORACIS

STRICTLY speaking the term, "empyema thoracis", should be confined to the description of a localized collection of pus within the pleural cavity. The early stages of bacterial invasion of the pleural cavity and the consequent formation of fluid which is not localized by adhesions must then be described by some other expression, such as generalized or diffuse suppurative pleurisy with effusion. However, in this chapter, in order to conform with common practice and for the sake of convenience, the term, empyema, will be used to cover all stages of suppurative pleurisy.

#### ACUTE EMPHYEMA

##### *Aetiology and pathology*

Because pneumonia is the most common precursor of pyogenic infection of the pleural space, there is a tendency to forget the many other conditions which may lead to suppurative pleurisy.

##### *Infections derived from the lungs*

When the lung is the source of the infection, it is always possible that some lesion other than pneumonia—such as bronchogenic carcinoma with secondary suppuration, lung abscess, bronchiectasis or actinomycosis—may be responsible for infecting the pleural cavity. Owing to the great increase in the incidence of carcinoma of the bronchus, it is particularly important to remember that this dread disease is commonly complicated by an empyema. Furthermore, when pneumonia is the cause of the pleural infection, the pulmonary lesion is often not of the classical pneumococcal or streptococcal type. For example, an empyema is not an uncommon complication of staphylococcal pneumonia in infants and small children, and is often preceded by a spontaneous pneumothorax, due to the rupture of a small subpleural abscess. The pleural space may also be infected from the lung to which the bacteria are carried by the blood stream as a result of a generalized pyaemia or septicaemia, but these cases are now rare owing to the widespread use of chemotherapy.


Before the introduction of the antibiotic drugs, pneumonia was frequently complicated by empyema, but there has been a very remarkable decrease in the incidence of this complication during recent years.

##### *Infections not derived from the lungs*


The infection causing an empyema, however, is not always derived from the lung. Bacteria may be carried into the chest by a penetrating wound, but suppuration is not common unless foreign matter is retained within the thorax, organic material such as clothing being more likely to lead to infection than

# ACUTE EMPYEMA

FIG 5 —(a) Right sided subphrenic abscess with a fluid level  
(b) a total pyopneumothorax after trans pleural aspiration of the subphrenic abscess



(a)



(b)



are metallic foreign bodies. However, accidental penetrating wounds of the thorax are not common in peace time, but the enormous increase in transpleural surgical operations, such as those on the lung, heart, oesophagus and sympathetic chain, have inevitably led to the surgeon's being responsible for a certain number of cases of empyema, although, with proper post-operative care, the incidence of pleural suppuration following such operations is very low. Other sources of infection of the pleural cavity include subphrenic abscess (Fig. 5), lesions of the oesophagus, and osteomyelitis of the ribs or spine. According to Harley (1949), who reviewed 182 cases of subphrenic abscess, empyema occurred in 29 per cent. Diseases of the oesophagus are not frequently complicated by an empyema, because the common lesions of this organ, such as carcinoma, are chronic, thus they allow time for the development of peri-oesophageal adhesions, which form a barrier to infection of the pleural space.

### Sex incidence

Because respiratory infections are more common in males, it naturally follows that pleural empyema has a similar sex incidence.

### Bacteriology

The bacteria most commonly responsible for pleural suppuration are (1) pneumococci, (2) haemolytic streptococci and (3) staphylococci, in that order of frequency. A mixed flora of anaerobic organisms is usually responsible for those cases of empyema which are secondary to foetid pulmonary suppuration, but these organisms may also be found in patients who have no clinical evidence of lung suppuration. In a few cases a pure growth of anaerobic streptococci is obtained from culture of the pleural pus. More commonly a non-haemolytic micro-aerophilic streptococcus is the cause of the infection, particularly in patients suffering from multiple areas of suppurative infection of the lung, most frequently seen as a sequel to a major abdominal operation. Occasionally an empyema develops as a result of invasion of the pleural space by coliform bacilli; these cases are almost invariably associated with gastro-intestinal disease, such as acute appendicitis. *Clostridium welchii* alone may be responsible for an empyema, but this is rare except in time of war, for this infection is usually introduced by a penetrating wound. Actinomycosis may give rise to suppuration in the pleural cavity and this possibility should not be forgotten, because culture of the pus will not yield any organisms after 48 hours' incubation and this may lead to a report of "sterile on culture", the nature of the infection can, however, usually be recognized by the presence of the classical 'sulphur granules'. In patients who have received a course of systemic chemotherapy, particularly when one of the antibiotics has been used, it is not uncommon to find that the pleural pus shows no organisms on microscopic examination and is sterile on culture. These findings may lead the clinician to suppose that tuberculosis is the most likely cause of the empyema, and to treat the patient according to this erroneous supposition—that is to say, external drainage is avoided and the empyema is allowed to become chronic.

### Morbid anatomy

The initial response to invasion of a normal pleural space by pyogenic bacteria

is acute inflammation of the two pleural surfaces, which is quickly followed by the effusion of a serous exudate, this is turbid on account of the presence of polymorphonuclear cells. Although the inflammatory process is unlikely to be of uniform severity throughout the pleural space, nevertheless, at this stage, the fluid is free in the pleural cavity. Subsequently the fluid becomes thicker, as the result of an increase in the number of dead and living polymorphonuclear cells and of a rise in the protein content of the fluid. This is usually accompanied by the deposition of fibrin, sometimes in very large amounts, particularly when pneumococci are responsible for the infection. In cases due to haemolytic streptococci, there is usually very little fibrin present, for these organisms produce a fibrinolysin which breaks down any fibrin deposit. On the other hand, an

of the fluid, the pus usually becomes localized to one part of the pleural cavity, most commonly the lower posterior part, as a result of fibrinous adhesion between the two pleural layers. In a few cases the pus may be subdivided into two or more separate pockets, because the lung becomes adherent to the parietes between these pockets (Fig 6). Much more commonly, the fluid pus is held in a number of loculi, which are separated only by a layer of fibrin that can easily be broken up with a finger. This loculation by fibrin is particularly pronounced in cases arising from infection of a haemothorax.

Occasionally the lung is everywhere in contact with the chest wall, and the pus is entirely confined to the space between two lobes. However, an interlobar empyema generally extends to the parietes, which means that a portion of the general pleural space is also involved. Very rarely the pus becomes isolated to a pocket lying between the lung and mediastinum ("mediastinal empyema"), so that the abscess has no contact with the chest wall. It is not strictly accurate to use the term "mediastinal empyema" for cases in which the pus, although largely confined to the space between the lung and mediastinum, also extends to reach part of the chest wall. Exceptionally an empyema is entirely confined to the space between the inferior surface of the lower lobe and the diaphragm, and may therefore be described as a diaphragmatic empyema.

With the lapse of time, further fibrin is deposited and organization of the deeper layers into fibrous tissue commences. Occasionally the pus perforates into a bronchus (broncho pleural fistula) and is expectorated.

### Clinical picture

#### *Comparative and clinical types*

in the course of a streptococcal broncho pneumonia, which was often a sequel to influenza. The patients were already seriously ill from the effects of the pneumonia, and infection of the pleural cavity led to a further deterioration of the general condition, marked by distressing dyspnoea and an increase in fever and pulse rate. Aspiration of the thin pleural fluid containing haemolytic

# EMPYEMA



FIG 6—Skilograms showing right sided empyema which was divided into two separate cavities (a) Taken after drainage of a basal para-vertebral cavity the upper cavity, which was situated posteriorly and extended into the upper part of the oblique fissure, was subsequently drained by resection of a portion of the eighth rib (b)



(b)

streptococci had to be repeated frequently, sometimes more than once a day, owing to its rapid re-accumulation, in order to relieve the patient's dyspnoea. Death often occurred and was inevitable if early drainage by rib resection was undertaken.

In the metapneumonic cases, the pleural fluid gradually formed after an attack of lobar pneumonia had passed its crisis. The pus, which was usually caused by pneumococci, thickened rapidly and soon became localized to one part of the pleural space. These patients were not seriously ill but, instead of making a progressive return to health following the febrile crisis, they either continued to run a low grade-fever or, if the temperature had returned to normal, again became febrile about a week after the pneumonic crisis.

Because pneumonia is now almost invariably controlled within 72 hours by appropriate chemotherapy, it is no longer practical to divide empyemas into synpneumonic and metapneumonic categories, but this old classification serves to remind the surgeon never to undertake drainage by rib resection until the empyema is localized by adhesions and the pneumonic process completely resolved.

#### *Symptoms and signs*

Nowadays the presence of pleural fluid following or accompanying pneumonia is usually recognized from routine physical or radiological examination, although severe pleural pain in the early stages of the pneumonia, or persistence of pyrexia in spite of adequate chemotherapy, may have already led the clinician to speculate on the likelihood of this complication. It is important to realize that the absence of fever by no means excludes the presence of pus, particularly if the pus is sterile

is usually denoted by a sudden deterioration in the general condition and by increased dyspnoea, but some cases develop remarkably insidiously. Occasionally, when an empyema does not receive prompt and adequate treatment, the pleural pus finds its escape within a few weeks by perforating into a bronchus. This complication leads to the sudden expectoration of large quantities of pus, but this development is often preceded by a period lasting several days, during which the patient develops an increasingly "wet" cough, due to irritation of the bronchus into which the pus later perforates. Just before the fistula is formed, the sputum sometimes becomes slightly blood-stained. After the initial episode occasioned by the formation of the broncho-pleural fistula, the patient will continue to expectorate pus and usually notices that this is exacerbated by a change in posture.

With regard to the physical signs, every student is taught the classical findings of impaired movement, absent tactile fremitus, dullness or impaired note on percussion, absent or reduced breath sounds, and diminished voice conduction

pression collapse of the lung tissue beneath the fluid. The position of the mediastinum is useful in distinguishing a pleural effusion from pulmonary atelectasis, but it cannot be relied on to differentiate between the two conditions,

for the mediastinum may not shift towards the opposite side unless an effusion is very large, and collapse of a whole lobe often occurs without causing the mediastinum to move towards the side of the lesion

### *Radiological appearances*

All types of pleural fluid—serous, purulent, bloody and chylous—produce a similar homogeneous opacity on x ray examination of the chest. In the earliest stages of an empyema the fluid is free in the pleural space, and its position is influenced to a certain degree by gravity, but to a much less extent than would be expected (Fig 7). With moderate-sized effusions, skiagrams taken with the patient sitting upright show a diffuse opacity at the base, and this often rises highest in the axilla. In films taken with the patient supine, the opacity becomes slightly less dense at the base and extends further upwards, but the change is usually not great. The skiagrams taken with the patient sitting upright may show a fluid level, but *only* if there is gas as well as fluid in the pleural cavity. A fluid level therefore indicates one of three possible alternatives: (1) air has been allowed to enter the pleural cavity from the exterior, either through a needle introduced at a previous paracentesis or as a result of a surgical or accidental wound, (2) a broncho-pleural fistula has formed, permitting air to pass from the lung into the empyema, or (3) the suppuration is caused by gas-producing organisms (Fig 8). With regard to the last-mentioned, the appearances after infection of a haemothorax by gas-producing organisms, such as *Clostridium welchii*, may be very striking for the gas is prone to become trapped in many fibrinous loculi, giving rise to multiple fluid levels.

As soon as the empyema becomes localized by adhesions, the radiological opacity remains constant whatever the position of the patient. Frontal and lateral skiagrams should be taken, using a penetration above that normally employed for routine chest radiography, in order to show the exact position and extent of the empyema. Empyemas of medium or small dimensions are usually located in the paravertebral gutter and extend upwards from the diaphragm, less commonly the pus is localized to the axillary region. The uncommon interlobar empyema gives rise in a lateral skiagram to an elliptical shadow in the line of one of the fissures. The rare mediastinal empyema appears in a frontal view as an opaque bulge, continuous with the opacity normally produced by the mediastinal structures.

### *Diagnostic aspiration*

The diagnosis requires confirmation by the aspiration of a sample of pleural fluid. In order to avoid unnecessary discomfort to the patient, it is preferable that the position of the fluid should be demonstrated radiologically before aspiration is attempted.


Paracentesis is performed under local anaesthesia, with the patient firmly but comfortably supported in bed. If the pus is located posteriorly, the patient may either lean forwards across a cardiac table or sit sideways on the edge of the bed (Fig 9a and b). If the axilla is chosen as the site for puncture, the patient need not move in bed, except to put the hand of the affected side behind his head so that the arm is lifted out of the operator's way (Fig 9c). Because the initial aspiration

# ACUTE EMPYEMA

is made primarily for diagnostic purposes it is preferable to select an intercostal space which is considered to lie well above the lowest limit of the empyema for attempts to enter the bottom of the cavity may lead either to penetration of the diaphragm which may be higher than expected or to obstruction of the needle by fibrin which tends to gravitate to the bottom of an empyema. The skin and soft

FIG. 7—Fluid free in the left pleural cavity (a) With the patient sitting up (b) with the patient recumbent

(a)



## EMPYEMA



FIG 8—Fetid empyema due to gas forming organisms. This is not a total empyema as the lung is adherent to the chest wall at the apex.

tissues over the selected site for aspiration are infiltrated with procaine solution through a fine hypodermic needle, and the skin is punctured with the point of a scalpel. A large bore needle (at least size 17 B W G) is then fixed to the end of a two way adaptor, which is connected to a 20-millilitre syringe filled with procaine solution. This needle is introduced into the skin puncture and steadily advanced through the chest wall, simultaneously infiltrating the tissues freely with procaine solution in order to ensure a painless puncture. Three test tubes, one of which contains a small quantity of 3.8 per cent sodium citrate solution, should be ready to receive samples of the aspirate, the tube containing citrate is used only if the fluid is found to be serous and therefore likely to clot. One of the non citrated specimens is placed in a rack in the ward and the others are sent for pathological examination.

If pus is obtained, it is worth while to continue the aspiration for therapeutic purposes, using the two way adaptor to discharge the fluid into a receiver, thus

pleural space at the end of the aspiration.

### Treatment

#### *Aspiration and chemotherapy*

If the presence of a pleural effusion is recognized early, if aspiration produces fluid from which the solid deposit on standing is less than one third of the total volume, and if the bacteria are sensitive to some antibiotic suitable for intra-

## ACUTE EMPYEMA




FIG. 9—Positions for pleural aspiration (see text)

may entail aspiration on alternate days or even daily. At the end of each aspiration the appropriate antibiotic is injected into the empyema and this should not be omitted until it is found impossible to demonstrate bacteria in a film of the pus on at least two consecutive occasions. It is more satisfactory to rely on microscopic examination of the pus than on cultures, for bacteria may be present and yet fail to grow when incubated on suitable media. In the case of penicillin sensitive bacteria, it is suggested above that the initial dose should be in the region of 500 000 units but, subsequently, it is doubtful whether anything is gained by exceeding 100 000 units at each intrapleural injection.

It cannot be over-emphasized that treatment solely by aspiration and local chemotherapy is justified only when the infected pleural effusion is recognized early, and it should not be continued unless the fluid becomes progressively thinner and ceases to reaccumulate within 2-3 weeks. There is a great tendency



almost entirely from the character of the pus. When the pus was thick, so that, on standing, the solid deposit exceeded half the total volume, the empyema was presumed to have become localized by adhesions. This was a fair assumption and still applies to-day. A sample of pus should therefore be taken at the time of each aspiration and placed in a rack in the ward, so that the amount of deposit on standing may be seen at a glance. If local chemotherapy is used, however, the pus may not become increasingly thick, although the empyema has in fact become localized. Consequently, evidence that the empyema has become

adhesions. In cases of doubt a small quantity of air, but not more than 10 millilitres, may be injected into the empyema, skiagrams being subsequently taken in the upright position, in order to determine whether this air is free to rise to the apex of the pleural cavity or whether, on the other hand, it is restricted to a limited empyema cavity (Fig. 11).

All localized empyemas should be drained by rib resection, even if the pus is sterile and the patient apparently free from toxæmia. Before doing so, 5 millilitres of iodized oil are injected into the cavity and frontal and lateral skiagrams taken, using increased penetration. The oil falls to the bottom of the empyema, and it is therefore possible to determine from the skiagrams the exact site at which

oil falls below this level, because it is claimed that drainage is likely to be hindered by a rise in the position of the diaphragm following operation. It is certainly uncommon for the oil to drop below the levels mentioned, but, when it does do so, there should be no hesitation in resecting a portion of rib at the level demonstrated as the bottom of the cavity, for the tube can always be lengthened after operation if the diaphragm rises.

## *Technique of rib resection and external drainage*

The operation is performed under local anaesthesia with the patient lying in that position which provides the surgeon with the most easy access to the site selected for drainage. For resection of a rib posteriorly or in the axilla, the patient lies on the "good" side, buttressed by a large support placed anteriorly. An operation should never be performed until the patient is fit enough to lie comfortably in this position for as long as is required. This does not apply to cases complicated by a broncho-pleural fistula, because operation must then be performed with the patient firmly supported in the sitting position, for recumbency is likely to cause the pus to enter the bronchial tree through the fistula and thus promote incessant coughing or even drown the patient.

An incision 3-4 inches long is made over the portion of rib selected for resection, . . . . . If the resection . . . . . along the lateral . . . . . in all other cases, the incision is preferably made in the line of the rib, although some surgeons

# ACUTE EMPYEMA

FIG 11—Frontal and lateral skiagrams of a localized empyema. Iodized oil has been injected to show the bottom of the cavity. Air in the upper part of the empyema shows that the cavity is limited by adhesions. (The air in the cavity was allowed to enter accidentally in this case and far exceeds the recommended maximum of 10 cc.)



(a)



(b)

recommend a vertical incision for all cases. At least 3 inches of the rib should be resected subperiosteally. A needle is then passed through the rib bed, in order to confirm that the site chosen for operation does in fact overlie the empyema. As soon as pus is obtained, a small incision is made through the rib bed so that the pus may escape slowly. When the flow of pus diminishes, the opening into the empyema is extended for the full length of the portion of rib resected. The rest of the pus is subsequently removed by suction and all fibrinous masses are extracted by means of sponge-holding forceps. The empyema is then inspected with the aid of a light suitable for introduction into the cavity, the light on the end of a cystoscope will serve this purpose if there is no special instrument available. Any fibrin adhering to the parietes or lung surface should be gently wiped away and extracted. The mobility of the lung may be ascertained by getting the patient to cough gently. If the empyema has not been entered at its lowest limit, a further portion of the same rib or a segment of another rib must be resected, so that the cavity may be drained from the bottom. A large rubber tube—at least  $\frac{1}{2}$  inch in internal diameter for an adult—is inserted so that the inner end projects about 1 inch inside the cavity. Finally the superficial muscles and skin are sutured around the drainage tube. This tube is held in position by transfixing it on a level with the skin with a safety-pin, which is anchored to the chest by long pieces of adhesive strapping. The dressings are applied around the tube and held in position with gate strapping. Bandages or strapping which encircle the chest should be avoided because they restrict respiratory movement. Unless the empyema is very small, the drainage tube should be connected to a water-seal bottle, in order that the patient may develop a moderate negative pressure within the empyema and so encourage re-expansion of the lung. Water-seal drainage also provides for the discharge to be carried away from the dressings (Fig. 12).

#### *Drainage by an intercostal tube—disadvantages*

In this discussion on treatment, the use of intercostal tube drainage has not been mentioned. Such drainage should never be employed for an adult in the treatment of a localized empyema, because the introduction of the tube does not provide an opportunity to remove fibrin and the calibre of the tube is necessarily too small to provide efficient drainage. Furthermore, an intercostal tube often becomes painful after a week or so, because of irritation of the intercostal nerve, and it sometimes causes pressure necrosis of part of one of the adjacent ribs. However, some clinicians maintain that, prior to localization of an empyema, the use of an intercostal tube connected to a water-seal bottle is preferable to repeated aspiration. Before the introduction of effective local chemotherapy, it could be argued that continuous drainage through an intercostal tube was better than intermittent aspiration, particularly in those cases of streptococcal effusion which re-collected very rapidly. But, with the use of antibiotics, it is rare to see an effusion re-collect quickly, and most empyemas soon become localized and ready for drainage by rib resection. It is therefore doubtful whether there is now any place for intercostal drainage in the treatment of empyema. If it is employed it should always be followed by rib resection, although an exception to this rule may sometimes be made in the case of small children.

*Post operative care*

period depending on the size of the empyema and the mobility of the walls of the cavity. In the case of large cavities re-expansion of the lung will be assisted



FIG. 12.—This photograph illustrates water seal drainage: the method of fixing the tube to the chest wall and the 'gate-strap' used for keeping the dressings in position.

by the negative pressure induced by water seal drainage provided that the lung is not bound down by rigid scar tissue. As a practical guide it is probable that water seal drainage is serving a useful purpose so long as the column of water in the glass tube shows a large respiratory "swing" and a mean negative pressure exceeding 10 centimetres. With an empyema of less than 500 millilitres capacity at the time of drainage it is usually unnecessary to persist with water seal drainage for more than a week but with large cavities it may be advisable to continue for 2-3 weeks. When the water seal is deemed to have completed its purpose the tube is cut off close to the safety pin with which it is anchored to the chest wall and any discharge subsequently drains into the dressings. This has the great advantage of allowing the patient to move more freely.

It is very important that the patient should perform breathing and postural exercises under the direction of a physiotherapist who is specially versed in this form of treatment. The exercises are started on the day following operation; they are continued until the empyema has completely healed and further until respiratory movements have been restored to normal. The patient should practice these exercises as much as possible throughout the day and not only in the presence of the physiotherapist.

Re-expansion of the lung will also occur more quickly if the patient is physically active. He should therefore leave his bed early and undertake as much exertion as the general condition permits. Most patients are able to get up within 24-48 hours of operation.

Irrigation of the empyema cavity which was popular at one time and is still

# EMPHYEMA

FIG 13—Serial "sino-grams" to show progress in obliteration of an acute empyema cavity. (a) The tube is correctly placed at the bottom of the cavity; (b) the cavity is slightly smaller. In (c) a tube of excessive length has been inserted and a bronchial fistula has developed (probably due to pressure from the end of the tube) (d) The fistula has healed although some residual oil can be seen in the alveoli. The tube is again too long and requires shortening. The tube is not removed until the sino-gram shows complete obliteration of the intra-thoracic cavity.


(a)

(b)

ACUTE EMPYEMA



(c)



(d)

frequently employed, should not be done, for it serves no useful purpose and may be harmful, because there is always a risk that secondary infection will be introduced. It is obviously strongly contra indicated if a bronchial fistula is present.

## *Control of the drainage tube*

The rapid obliteration of an empyema depends more on the maintenance of perfect drainage than on any other factor. Lateral and frontal skiagrams should be taken within 48 hours of surgical drainage. A fluid level in the empyema indicates imperfect drainage, this may be due to blockage of the tube by fibrin, in other cases the tube is too long and so projects above the bottom of the cavity, but sometimes it is too short, so that the inner end has been withdrawn from the cavity into the chest wall. In any of these circumstances, the tube must receive immediate and appropriate adjustment.

The length of the tube is subsequently controlled by frontal and lateral skiagrams, taken after introducing 5 millilitres of iodized oil into the empyema through the tube. The oil should be injected with the patient lying down, the tube is then occluded with a spigot and the skiagrams are taken with the patient in the upright position. From these "sinograms" or "pleurograms", which should be taken at intervals not exceeding 2 weeks, it is possible to determine the relation of the inner end of the tube to the bottom of the cavity (Fig 13). The tube can then be adjusted so as to provide dependent drainage. These films usually also show the size of the empyema. The tube must not be removed until "sinograms" have shown that the empyema cavity has been completely obliterated. This usually occurs 4-10 weeks after operation. Large cavities and those drained after the optimal time require the longest period of drainage. Patients who require prolonged drainage should be encouraged to return to their employment with the tube still in position.

## *Possible future advances in treatment*

One of the strongest arguments in favour of performing rib resection in all cases of localized empyema is that this provides an opportunity to remove all the fibrin from the cavity. It is now possible to obtain commercially enzymes which are capable of breaking down within the human body the solid constituents of an inflammatory exudate. These enzymes, which are derived from the

and Read, 1950). It is possible that the intrapleural injection of these enzymes may obviate the necessity for drainage of some cases of acute empyema but it would be unwise to recommend this as a routine form of treatment until these ferments have received more extensive trial. They are likely to be of particular value in empyemas which arise from infection of a haemothorax.

During recent years some surgeons have tried to reduce the period of convalescence by first sterilizing the empyema with intrapleural chemotherapy and then treating the residual cavity by decortication (see page 44). Complete decortication followed by an uncomplicated convalescence certainly gives quick results but decortication is a major operation compared with rib resection drainage

## ACUTE EMPYEMA

and is not always followed by complete and rapid re-expansion of the lung. It is therefore advisable to reserve this form of treatment for those patients who are seen in the first instance several weeks after the optimum time for external drainage.

**Empyema complicating diseases other than pneumonia**

(1) *Secondary to septicaemia or pyaemia*—In patients with a generalized blood infection, the occurrence of pleural pain, dyspnoea or an unexplained deterioration in general condition should lead to a careful investigation for evidence of infection of the pleural space. Fortunately, these cases, which were often bilateral, are now rare, as the result of the widespread use of chemotherapy.

(2) *Empyema secondary to subphrenic abscess*—Reference is made above to Harley's review of 182 cases of subphrenic abscess (Harley, 1949), in which he found that empyema occurred in 29 per cent, but that 50 per cent of these cases followed injury to the diaphragm, caused by transpleural drainage, a penetrating wound or transpleural aspiration (Fig 5). On the other hand, he found that a serous pleural effusion occurred in 25 per cent. This means that, if the diaphragm is uninjured, a pleural effusion arising in the course of a subphrenic abscess is more likely to be serous than suppurative, which emphasizes the importance of diagnostic aspiration in these cases. A serous effusion will clear up spontaneously provided that the subphrenic abscess is drained satisfactorily. An infected effusion should receive treatment similar to that described above for a post-pneumonic empyema. When the empyema is localized and ready for external drainage, the site for rib resection should be chosen by the method described above and should be separate from the wound made to drain the subphrenic abscess.

(3) *Empyema secondary to bronchiectasis*—If the infection associated with bronchiectasis extends into the lung parenchyma, the patient suffers an acute febrile episode, usually described as pneumonia. In some cases, if the pleural space is not obliterated by adhesions, the infection spreads further and gives rise to an empyema. This possibility should not be forgotten because the diagnosis is apt to be missed, so that the empyema becomes chronic, and then, if a broncho-pleural fistula develops, the increased expectoration is attributed to an exacerbation of the bronchial infection.

(4) *Empyema secondary to lung abscess*—If a foetid lung abscess ruptures into the pleural space, the patient often becomes dangerously ill within a few hours. In these cases the injection of penicillin into the stinking pyopneumothorax is often life-saving.

(5) *Empyema secondary to bronchiogenic carcinoma*—In patients of cancer age who develop an empyema, the possibility of an underlying growth should always be considered, particularly in males, and if the acute illness was preceded by a considerable period of lassitude or by symptoms of bronchial irritation. In cases of doubt the patient should be subjected to bronchoscopy before deciding on treatment of the empyema. If a growth is found, the presence of the empyema does not exclude the possibility of successful resection. If there are other factors indicating that the growth is inoperable, every effort should be made to control the empyema by aspiration and intrapleural chemotherapy, so that external



drainage may be avoided, for a discharging wound only adds to the miseries of a dying patient

(6) *Empyema due to actinomycosis*—Infection of the pleural cavity by actinomycosis characteristically produces a thin layer of pus with much loculation, but, exceptionally, a single cavity simulating a postpneumonic empyema occurs. Infection of the pleura is often followed by much pain and later by induration of the chest wall and sinus formation. Drainage operations are rarely required for almost all cases clear up completely and permanently as a result of prolonged and intensive systemic penicillin therapy. It is suggested that at least 2,000,000 units should be given daily for a minimum period of 6 weeks.

### Complications of acute empyema

(1) *Broncho pleural fistula*—Fistulae which complicate an acute empyema almost invariably heal after dependent drainage of the empyema.

(2) *Failure of the lung to re expand and obliterate the empyema cavity*—See "Chronic Empyema"

(3) *Infection of the chest wall*—Before antibiotics were used locally to control the infection in empyemas, it was common to see an abscess in the chest wall at the site of a previous paracentesis, but such cases are now rare. In a few instances, spreading infective necrosis of the skin has been observed, usually after drainage of the empyema. These serious cases should be treated by immediate excision of the affected tissue with the diathermy knife, and the raw area so caused should be subsequently grafted with skin.

(4) *Haemorrhage from the intercostal vessels, following drainage of the empyema*—This rare complication may be recognized by the occurrence of persistent and profuse haemorrhage around the tube. The tube should be removed at once and the wound packed with a gauze roll. The patient is then taken to the operating theatre so that the wound may be re opened and the vessels ligated. Some surgeons claim that this complication does not occur if the vessels are routinely divided between ligatures at the time of draining the empyema.

(5) *Brain abscess*—Because it is rare for a straightforward acute empyema to give rise to a cerebral abscess, this complication is considered at the end of the discussion on chronic empyema.

## CHRONIC EMPYEMA

It is obvious that all cases of chronic empyema have passed through an acute stage and that the transition from the acute to the chronic stage is a gradual one, so that there is no clear line dividing the two conditions. Furthermore the conditions under which an empyema may be reasonably defined as "chronic" are subject to the personal views of individual clinicians, but it is suggested here that the term, chronic, should be applied to those cases in which a cavity persists for more than 3 months after the original pleural effusion.

### Aetiology and pathology

If an acute pyogenic empyema is not drained externally or aborted by aspiration and local chemotherapy, more and more fibrous tissue is laid down, particu-

larly on the parietal layer of pleura thus immobilizing the chest wall and preventing expansion of the lung. It is necessary to emphasize that the fibrous tissue is laid down on the surface of the pleura and therefore should not be called pleural thickening, although it is often described as such. Sometimes the pus finds its escape through a broncho pleural fistula and is expectorated this is mentioned above under Acute Empyema. In other cases the chest wall is eroded and some of the pus forms an abscess beneath the overlying skin and superficial muscles—a condition designated empyema necessitatis (Fig 14) this is soon followed




Fig. 14—Empyema necessitatis due to actinomycosis. (a) the swelling above and medial to the left nipple is a subcutaneous abscess communicating with an empyema. (b) 4½ months later after treatment with systemic penicillin (440 mega units were given over a period of 3 months) therefore no operative treatment was employed.

by necrosis of the tissues covering the abscess and the pus escapes externally leaving a sinus which may heal intermittently or discharge persistently.

The fibrous tissue formed during this chronic stage gradually contracts so that there is progressive flattening of the chest wall which is accompanied by a scoliosis with the concavity towards the affected side. In addition the mediastinum which may have been displaced towards the healthy side during the acute phase is slowly drawn towards the side of the empyema (Fig 15). The chronic inflammation also causes an irritation of the periosteum of the ribs overlying the empyema and consequently new bone is laid down on the inner surface of the ribs so that they become triangular in cross section (Fig 16).

These changes are not confined to persistent undrained cavities for exactly the same series of events occurs when a drained empyema is not rapidly obliterated by re-expansion of the lung. The most common cause of persistence of an empyema in spite of external drainage is premature removal of the tube. A



FIG 15—Left sided chronic empyema in a girl aged 18 years which started as an acute empyema at the age of 6 years. Iodized oil has been injected into the cavity and is just visible at its base

FIG 16—Chronic empyema showing an extreme degree of periosteal new bone formation. The cavity has been drained and a tube which is far too long inserted. Iodized oil injected through the tube has entered a bronchial fistula and outlined the bronchial tree



residual cavity is also likely to occur if there is any interference with free drainage, as, for example, when the tube is too long or too short, or when the drainage opening is not at the bottom of the cavity. Sometimes an empyema fails to heal because the pleural infection is maintained by the presence of a foreign body, such as a piece of drainage material or a sequestered portion of rib. In other cases the infection may be kept up by an unsuspected aetiological factor, such as tuberculosis, actinomycosis or pleural carcinomatosis. A chronic cavity may also persist because the lung is not free to expand. This may be due to the presence of intrinsic disease, such as carcinoma, or fibrosis associated with a chronic lung abscess or bronchiectasis. More commonly, however, the lung is healthy, but is unable to re-expand because it is covered with a rigid layer of fibrous tissue. The latter may occur as the result of a delay in draining the empyema or because the surgeon omitted to remove all the fibrin at the time of operation. The lung is also likely to become covered with a thick layer of scar tissue if a total empyema has been caused by drainage of a septic pleural effusion before it has become localized by adhesions. A broncho-pleural fistula sometimes leads to a delay in re-expansion of the lung but, unless the fistula is very large, it does not often cause an empyema to become chronic, provided that the underlying lung is otherwise healthy.

### Clinical picture

#### *Clinical types*

Two clinical types of chronic pyogenic empyema may be recognized: (1) "latent", in which drainage has never been performed, either because the diagnosis was not made during the acute stage or because the clinician did not recognize the need for external drainage; (2) "persistent", in which a cavity persists in spite of external drainage.

*Latent empyema*—A patient with a latent empyema may be free from symptoms, and may have been referred for treatment because the signs of an encysted collection of fluid have been discovered on clinical or radiological examination. This applies particularly to those cases in which the pus is sterile as a result of either local or systemic chemotherapy. In other cases the patient complains of persistent fever, chest pain or dyspnoea following an acute respiratory infection. Sometimes the patient notices that, after recovering from an attack of pneumonia, he has not returned to the same state of vigorous health which he enjoyed before his illness. Then, after an interval which varies from a few weeks to several years, he may suffer an acute febrile episode accompanied by pleural pain. If the correct diagnosis is not recognized and appropriate treatment instituted, these symptoms may gradually subside but are liable to recur again and again. As long as a latent empyema persists, it may at any time develop into an empyema necessitatis: the patient will then complain of a tender swelling in the chest wall.

*Persistent empyema*—A patient with a "persistent" chronic empyema most commonly complains because there is a continuous or intermittent discharge from the drainage wound following removal of the tube. Sometimes the wound heals and remains healed for quite a long time, perhaps for a period of several years, and then becomes the site of an abscess which is, in fact, an empyema necessitatis. In other cases of "persistent" empyema the tube has never been

removed, and further treatment is sought because the empyema cavity does not show signs of undergoing progressive obliteration

*Broncho-pleural fistula*—A latent empyema or an ill drained persistent cavity may at any time lead to the formation of a broncho-pleural fistula: the most prominent symptom is then likely to be cough with purulent, and sometimes blood-stained, expectoration

All those affected with chronic empyema are prone to suffer from lassitude and other symptoms of poor general health

### *Signs of chronic empyema*

The complexion of a patient with a chronic empyema is frequently pale and "muddy", if the septic process has continued active for more than a month or two. Pleural suppuration almost invariably leads to clubbing of the fingers and toes, sometimes within a few weeks of the onset of the infection. Physical signs in the chest include gross flattening of the affected side, and diminution or absence of movement on respiration. The classical signs of pleural fluid are usually present over the site of the empyema. If the empyema is extensive it is common to find that the trachea and apex beat are displaced towards the affected side, on account of a shift in the position of the whole mediastinum. In cases of empyema necessitatis, a tender soft swelling is found superficial to the ribs, and this usually has an impulse when the patient coughs (Fig 14)

### *Radiological appearances*

A "latent" empyema shows as a dense opacity over which the ribs are approximated. Periosteal new bone on the inner aspect of these ribs is frequently visible and is pathognomonic of chronic pleural disease, it is particularly marked when the infection is caused by actinomycosis. If the empyema has been drained or has discharged spontaneously, or if a broncho-pleural fistula has developed, it is common to see a fluid level in the cavity, in which case it is usually possible to see the thickness of the fibrous tissue lining the cavity above the fluid level. Skiagrams will also show the severity of the scoliosis, and the extent to which the diaphragm has been elevated and the mediastinum shifted as a result of the chronic fibrosis. The films should be examined carefully for evidence of a foreign body within the empyema cavity, but, owing to the density of the opacity caused by the chronic empyema, it is often difficult to see a foreign body even when it is very radio-opaque, as, for example, with a rib sequestrum.

### *Special investigation*

If the empyema is not discharging externally, a sample of the fluid should be obtained by aspiration and sent for bacteriological investigation. The operator may have difficulty in passing the needle through the chest wall, because of narrowing of the intercostal spaces, and he must be prepared to meet with considerable resistance to insertion of the needle, for the fibrous tissue lining the empyema is often tough. The pus is usually thick, but it may be quite thin if the fluid has become sterile.

When there is a sinus present or a tube *in situ*, a sample of the discharge should be collected in a test tube and examined bacteriologically. It is particularly important to search for evidence of tuberculosis, because the treatment of a

tuberculous empyema may differ considerably from that adopted for a chronic pyogenic case. With this in mind, the granulation tissue at the orifice of a sinus should be inspected and, if it is pale and oedematous, which suggests tuberculous infection, a small portion should be removed for histological examination.

The length of an empyema cavity may be estimated in cases in which there is a sinus, by passing a fine gum-elastic bougie into it. If the empyema is still being drained with a tube, the tube should be removed for a few moments and the patient rotated into various positions in order to see whether or not the drainage is adequate; the discharge of a considerable quantity of pus after removal of the tube is clear evidence that the drainage provided is inadequate. In those cases in which there is reason to suspect the presence of intrinsic disease of the lung, such as carcinoma, appropriate investigation should be undertaken before treatment of the empyema is started.

### Treatment

A chronic empyema can be cured permanently only by obliteration of the cavity. This may occur either as the result of re-expansion of the lung or in consequence of operative mobilization of the chest wall; in the latter case, the parietal pleura falls inwards to meet and fuse with the visceral layer where it covers the partially collapsed lung. It is obvious that the functional and cosmetic result is much better when the cavity is obliterated by re-expansion of the lung; plastic operations on the chest wall should therefore be reserved for cases in which complete re-expansion of the lung proves to be impossible.

### Re-drainage

Many cases of chronic empyema will heal by re-expansion of the lung if proper dependent drainage is provided. In those instances in which there is a discharging sinus or a drainage tube *in situ*, 5 millilitres of iodized oil should be introduced into the cavity through the sinus or tube, and skiagrams should be taken in the antero-posterior and lateral positions. From these it is possible to decide the position which a tube should occupy in order to provide dependent drainage. If a sinus leads into the bottom of the empyema, it may be dilated by the introduction of a laminaria tent, and a tube of adequate calibre subsequently fixed in position. A sinus should never be dilated with bougies, for cases have been reported in which this procedure has been followed by the formation of a brain abscess. If the sinus or tube does not lead into the bottom of the cavity, dependent drainage should be provided by the resection of a portion of rib; the site for this operation may be determined from the skiagrams taken after the introduction of iodized oil. Resection of a rib gives an opportunity to obtain a biopsy of the cavity wall for examination for evidence of tuberculosis, and also to inspect the interior of the cavity and thus exclude the presence of a foreign body. Therefore, even in those cases in which there is already an opening into the bottom of the cavity, resection of a portion of rib is usually preferable to dilatation of the sinus with a laminaria tent. The drainage tube is cut flush with the chest wall and allowed to discharge into the dressings, because the mild negative pressure induced by water-seal drainage is insufficient to influence re-expansion of the lung in a chronic case. It is possible to use much higher negative



FIG. 17—Decortication for achronic calcified expectorated empyema. Serial films showed that the calcium in the walls of the cavity was breaking up and being shed into the cavity. This case probably started 7 years previously as a spontaneous haemopneumothorax. (a) 8 months after decortication.



## CHRONIC EMPYEMA

mobilized to allow it to fall on to the visceral wall of the empyema. It is held in this position by a pack placed on its outer surface. The skin and superficial muscles are sutured over this pack, leaving a small opening for drainage. The pack is removed after 10 days and the wound firmly strapped over a large pad of gauze and cotton wool. With moderate and large-sized cavities this operation is preferable to Schede's operation, because it is more likely to be successful, and there is less deformity as well as a quicker convalescence.

### Complications of chronic empyema

**Complications due to chronic toxæmia.**—The most common complications are those which arise as the result of a chronic toxæmia, they include pulmonary osteoarthropathy, amyloid disease and nephritis. The symptoms and joint effusions associated with pulmonary osteoarthropathy usually clear up quickly as soon as the chronic suppuration in the chest has been controlled. It is important to realize that amyloid disease is also reversible, provided that it has not been present over a long period. In many cases the symptoms and signs of amyloid degeneration improve greatly but do not disappear entirely, albuminuria is particularly apt to persist in a mild form. Although nephritis is not common it is of particular importance to the surgeon, because patients with this complication are often bad risks for major surgical operations.

**Brain abscess**—A chronic empyema may also be responsible for pyaemic abscesses, particularly in the brain. A persistent headache with nausea or vomiting should always lead the clinician to suspect a brain abscess, but, even if these suspicions are confirmed, all hope of the patient's recovery need not be abandoned, for numerous authenticated cases have been successfully treated during recent years by appropriate neurosurgery.

## BIBLIOGRAPHY

- Asher, R., and Davies, D. (1951) "Empyema Thoracis Treated with Penicillin and Aspiration Only" *Lancet*, 1, 924
- Blades, B., Hamilton, J. E., and Dugan, D. J. (1945) "Observations on the Treatment of Empyema Thoracis with Penicillin" *Surgery*, 17, 572
- Brook, R. C., and Sellors, T. H. (1947) "Discussion on Treatment of Non tuberculous Empyema" *Proc R Soc Med*, 40, 645
- Brown, B., Ory, E. M., Meads, M., and Finland, M. (1946) "Penicillin Treatment of Empyema: report of 24 cases and review of the literature" *Ann Intern Med*, 24, 343
- Burford, T. H., Parker, E. F., and Samson, P. C. (1945) "Early Pulmonary Decortication in the Treatment of Acute Empyema" *Gaz hop Paris*, 67, 94
- Fatti, L., Florey, M. E., and colleagues (1946) "Acute Infections of Pleura Treated with and without Penicillin" *Lancet*, 1, 257
- — — (1946) "Acutely Infected Pleural Effusions: techniques of Penicillin Treatment" *Lancet*, 1, 295
- Graham, E. A. (1925) *Some Fundamental Considerations in the Treatment of Empyema Thoracis* St Louis, Mosby
- Harley H. R. S. (1949) "Subphrenic Abscess" *Thorax*, 4, 1
- Lawman, T. H., and Dimmler, C. L. (1941) "The Management of Acute Empyema in Children" *Amer J Surg*, 54, 29



## EMPHYEMA

- Mullard K S (1951) Empyema Thoracis *Post grad med J* 27, 104
- Parker E F and Burford T H (1948) Pulmonary Decortication in the Treatment of Early Pyogenic Empyema *Surgery* 24 371
- Roberts J E H (1935) The Surgery of Pleural and Pulmonary Infections *Trans med soc Lond* 58 183
- Schede M (1890) Die Behandlung der empyeme *Verh Kongr inn Med* 9 41
- Sellors T H and Cruickshank G (1951) Chronic Empyema *Brit J Surg* 38 411
- Sherry S Tillett W S and Read C T (1950) The Use of Streptokinase streptodornase in the Treatment of Hemothorax *J Thorac Surg* 20 393
- Tillett W S Sherry S and Read C T (1951) The Use of Streptokinase streptodornase in the Treatment of Postpneumonic Empyema *J Thorac Surg* 21, 275

## CHAPTER 3

### HAEMOTHORAX

KENNETH M A PERRY

#### AETIOLOGY

##### **Traumatic haemothorax**

THE MAJORITY of cases of haemothorax or of haemopneumothorax follow trauma, which is usually associated with perforating wounds, but occasionally with non-perforating blows, even without fractured ribs. This subject was well reviewed by Ogilvie (1950) who found haemothorax in 230 (58 per cent) of his 400 cases of chest injury. Similar figures have been found by other authors. Bradford and Elliott (1915) 156 (92 per cent) out of 170, Kay and Meade (1946) 455 (91 per cent) out of 500, and Thomas and Cleland (1945) 520 (70 per cent) in a series of 750. Traumatic pneumothorax becomes infected in about 22 per cent of cases, the infection being either introduced from without or resulting from inflammatory changes in the lung. A large haemothorax is not so likely to become infected as a small one. *Staphylococcus aureus* is the infecting organism in 50 per cent of cases, but other common invaders include streptococci, *Bacterium coli*, *Clostridium welchii* and other anaerobic organisms.

**Surgical trauma**—Haemothorax is a rare complication of artificial pneumothorax therapy, in which a vessel may be damaged by the needle, 7 cases were reported on by Miller and Rinkel (1947). Haemothorax may also result from thoracoscopy for adhesion section in about 4 per cent of cases, this is reviewed by Harold (1951).

##### **"Spontaneous" haemothorax**

Apart from these accidents there is a rare but definite entity of "spontaneous" haemothorax or haemopneumothorax. In the majority of patients the disease occurs between the ages of 20 and 40 years, and in most of the cases described in the literature the condition has occurred in males. When necropsies have been performed the cause has been found to be the rupture of a pleural adhesion which contained a blood vessel. This accident may occur in a healthy individual or during the course of treatment for pulmonary tuberculosis. Whenever the patient has died, haemorrhage has been the cause of death (Hopkins, 1937; Perry, 1938).

##### ***Haemothorax as a complication of other diseases***

Rupture of either a syphilitic or a dissecting aneurysm of the aorta into the pleural cavity (East, 1939; Post, 1941) may occur as a terminal complication of these diseases.


Endometrioma of the pleura is a rare condition, which occurs in women and will produce spontaneous haemothorax, it is usually associated with endometrioma elsewhere so that ascites may also be present (Nicholson, 1951) (Fig 18).

Apart from these causes of frank haemothorax, blood-stained pleural effusion

# HAEMOTHORAX



FIG 18 —(a) Skia gram of a woman aged 39 years with a large haemothorax and ascites resulting from endometriomas of ovaries colon and pleura (b) after removal of the blood stained effusion






FIG. 19.—Sk agram of a man who developed an internal haemorrhage into left pleural cavity three weeks after fracturing ribs. Six pints of blood were removed at thoracotomy.

occurs as a complication of carcinoma of the lung pulmonary tuberculosis Hodgkin's disease pneumonia cirrhosis of the liver reticulosis scarlet fever smallpox arterial hypertension and mitral stenosis (Vaizey and Perry 1940)

### SYMPTOMATOLOGY

In cases which do not result from trauma the patient usually gives a history of sudden onset of sharp stabbing pain in the side of the chest although sometimes it is located in the upper part of the abdomen. This is followed by shortness of breath. If the bleeding is extensive the patient will be shocked and collapsed. He will rapidly become pale. On examination he may be febrile. There will be a raised pulse and respiration rate. If the haemothorax is rapid he will give the appearance of a patient suffering from internal haemorrhage but if it is less severe and if the lung is torn at the time that the haemorrhage occurs he will have the appearance of a patient with a pleural effusion or a pneumothorax. The trachea is likely to be displaced away from the side of the lesion. On the side involved there will be diminished movement an absence of tactile vocal fremitus impaired percussion notes or stony dullness and distant or absent breath sounds. If there is a haemopneumothorax there will be hyper resonance above the dullness. Sometimes owing to irritation of the diaphragmatic pleura by the blood there may be rigidity of the upper abdomen. This may be very marked and simulate the rigidity of a perforated peptic ulcer. Of the cases of

this disease reported in the literature more than half have been operated on with this mistaken diagnosis

### THE FATE OF BLOOD IN THE PLEURAL CAVITY

The fate of blood in the pleural space has long been a debated subject, but as long ago as 1834 LeBlanc and Trousseau showed that blood injected into the pleural cavity of horses first clotted and then was absorbed, and in 1916 Denny and Minot showed that blood stained fluid in the pleural cavity consisted only of serum and cells

The problem has recently been studied by Ogilvie (1950) and by Harold (1951) and it is now generally accepted that blood in the pleural cavity clots at once, but that this clot is soon broken up and dispersed by the movements of the heart and lungs. The blood is thus defibrinated and the fibrin either is deposited on the pleura or sinks in clots to the bottom of the pleural space. The pleural cavity thus becomes like a blood-cyst lined with fibrin, thicker on the parietal than on the visceral pleura. The blood which would then be aspirated resembles whole blood but, in fact, cannot clot since it has lost its fibrinogen. This blood acts as a foreign body and stimulates the outpouring of fluid, which reaches a maximum within 3 days. The blood is thus diluted and the fibrinogen concentration rises, thus making secondary clotting possible in the second week, this can be prevented by early aspiration. Sellors (1945) gives evidence supporting this theory, by showing that there is a steady fall in haemoglobin concentration in the fluid, and as steady a rise in fibrinogen over the period of 7 days after the haemorrhage. If the blood is not removed an organizing haemothorax results. This process has been described by Samson and Burford (1947). Usually the pleura remains normal though coated with fibrin, but in some cases the scar tissue extends through the pleura into the interstitial tissue of the lung and this may develop more rapidly if an empyema is present. Ogilvie (1950), however, quotes cases in which dense fibrous adhesions have been found at operation in the second week, even in cases of sterile organizing haemothorax.

### PROGNOSIS AND TREATMENT

Ogilvie (1950) showed that the average duration of illness, in cases of sterile traumatic haemothorax, adequately treated by aspiration was about 6 weeks, whereas if the treatment was inadequate this time was doubled. In his whole series of 230 cases, which included infected, clotted and organizing haemothoraces operation was necessary in 63 cases (27 per cent) and that in these complicated cases the duration of illness was much longer and physical and psychological results were less satisfactory, although there was only 1 death.

In cases which occur during artificial pneumothorax therapy treatment, if adequate, is so satisfactory that a useful surviving artificial pneumothorax is the rule. Deaths are rare, as is also the case in spontaneous haemothorax.

Endometrioma may be excised and an artificial menopause induced by radiation therapy. In these cases also the prognosis is good. The prognosis of blood stained effusions depends on the aetiology of the underlying disease.

## PROGNOSIS AND TREATMENT

### Medical treatment

If all the fluid can be removed from the chest and the bleeding stops, aspiration and treatment of the anaemia may be the only treatment required. Aspirations should always be started as soon as possible. They should be continued so that the pleural cavity is kept dry. Intrapleural injection of streptokinase will help to break down fibrin. If the anaemia is not severe, the patient responds quickly to iron, but in some instances blood transfusion will be necessary.

### Surgical treatment

In cases in which blood clot is present, aspiration will not be possible, these cases have been described as clotted haemothorax. Early thoracotomy with removal of the blood clot is then essential. Blood clot left in the pleural cavity causes rapid thickening of the pleura and frequently calcification.

Haemorrhage may be so severe as to cause the death of the patient. In such cases immediate thoracotomy, with removal of the blood clot and tying of the bleeding point, is a life saving operation.

When the blood clot becomes infected or an empyema develops, chemotherapy, systemic and intrapleural, with the drug to which the infecting organism is sensitive is indicated. There is much to be said however, for giving penicillin in all cases as a prophylactic. If an empyema develops it may require drainage.

In cases of clotted or organizing haemothorax, decortication may become necessary, if surgery is needed on these lines it is probably best carried out in the fourth week of the illness.

## BIBLIOGRAPHY

- Bradford J R and Elliot T R (1915) Haemothorax *Brit J Surg* 3, 247  
Denny G P and Minot G R (1916) The Coagulation of Blood in the Pleural Cavity " *Amer J Physiol* 39, 435  
East T (1939) Dissecting Aneurysm of the Aorta. *Lancet* 2, 1071  
Harold J T (1931) Intrapleural Haemorrhage in Artificial Pneumothorax. *Thorax* 6, 162  
Hopkins H U (1937) Spontaneous Haemopneumothorax. Report of 3 cases with a review of the literature. *Amer J med Sci* 193, 763  
Kay E B and Meade R H (1946) War Injuries of the Chest " *Surg Gynec Obstet* 82, 13  
LeP  
Mil  
Nicholson H (1931) Endometrosis of the Pleura. " *Thorax* 6, 75  
Nicholson W F (1946) Penetrating Wounds of the Chest. *Brit J Surg* 33, 257  
Ogilvie A G (1950) Final Results in Traumatic Haemothorax. A report on 230 cases " *Thorax* 5, 116  
Perry K M A (1938) " Spontaneous Haemothorax. *Lancet* 2, 829  
Post F (1941) Haemothorax from Leakage of a Dissecting Aortic Aneurysm " *Lancet* 2, 558  
Samson P C and Burford, T H (1947) Total Pulmonary Decortication. Its evolution and present concepts of indications and operative technique " *J thorac Surg* 16, 127  
Sellers T H (1945) " Haemothorax. *Lancet* 1, 143  
Thomas C P and Cleland, W P (1945) Decortication in Clotted and Infected Haemothoraces " *Lancet* 1, 327  
Vaurey J M and Perry K M A (1940) " Serous Pleural Effusions. *Lancet* 1, 1120

this disease reported in the literature more than half have been operated on with this mistaken diagnosis

### THE FATE OF BLOOD IN THE PLEURAL CAVITY

The fate of blood in the pleural space has long been a debated subject, but as long ago as 1834 LeBlanc and Trousseau showed that blood injected into the pleural cavity of horses first clotted and then was absorbed, and in 1916 Denny and Minot showed that blood stained fluid in the pleural cavity consisted only of serum and cells

The problem has recently been studied by Ogilvie (1950) and by Harold (1951), and it is now generally accepted that blood in the pleural cavity clots at once, but that this clot is soon broken up and dispersed by the movements of the heart and lungs. The blood is thus defibrinated and the fibrin either is deposited on the pleura or sinks in clots to the bottom of the pleural space. The pleural cavity thus becomes like a blood cyst lined with fibrin, thicker on the parietal than on the visceral pleura. The blood which would then be aspirated resembles whole blood but, in fact, cannot clot since it has lost its fibrinogen. This blood acts as a foreign body and stimulates the outpouring of fluid, which reaches a maximum within 3 days. The blood is thus diluted and the fibrinogen concentration rises, thus making secondary clotting possible in the second week, this can be prevented by early aspiration. Sellors (1945) gives evidence supporting this theory, by showing that there is a steady fall in haemoglobin concentration in the fluid, and as steady a rise in fibrinogen over the period of 7 days after the haemorrhage. If the blood is not removed an organizing haemothorax results. This process has been described by Samson and Burford (1947). Usually the pleura remains normal though coated with fibrin but in some cases the scar tissue extends through the pleura into the interstitial tissue of the lung, and this may develop more rapidly if an empyema is present. Ogilvie (1950), however, quotes cases in which dense fibrous adhesions have been found at operation in the second week, even in cases of sterile organizing haemothorax.

### PROGNOSIS AND TREATMENT

Ogilvie (1950) showed that the average duration of illness, in cases of sterile traumatic haemothorax, adequately treated by aspiration, was about 6 weeks, whereas if the treatment was inadequate this time was doubled. In his whole series of 230 cases, which included infected, clotted and organizing haemothoraces, operation was necessary in 63 cases (27 per cent), and that in these complicated cases the duration of illness was much longer and physical and psychological results were less satisfactory, although there was only 1 death.

In cases which occur during artificial pneumothorax therapy, treatment, if adequate, is so satisfactory that a useful surviving artificial pneumothorax is the rule. Deaths are rare, as is also the case in spontaneous haemothorax.

Endometrioma may be excised and an artificial menopause induced by radiation therapy. In these cases also the prognosis is good. The prognosis of blood-stained effusions depends on the aetiology of the underlying disease.





## CHAPTER 4

### SPONTANEOUS PNEUMOTHORAX

KENNETH M A PERRY

#### AETIOLOGY

IN PNEUMOTHORAX the air collects between the layers of the pleura causing it to become a real instead of a potential space. When fluid, pus or blood are also present it is known as a hydropneumothorax, pyopneumothorax, or haemopneumothorax. Air may gain access to the pleural space through the visceral pleura, from the lungs, through the chest wall as a result of penetrating wounds, from the mediastinal organs, through the diaphragm from some hollow viscus, or from gas-producing organisms infecting the pleura.

Aetiologically the causes of spontaneous pneumothorax may be divided into three groups. The first group includes symptomatic pneumothorax which may result from rupture of the parietal pleura as a result of trauma caused by penetrating wounds, when a sucking pneumothorax may occur, or non penetrating injuries which result in laceration of the pleura with or without fractured ribs. The history of injury will be evident. The visceral pleura may be perforated in empyema and also from lung abscess, particularly staphylococcal lung abscesses in children, which may result from bronchogenic infection or staphylococcal septicaemia (Collis and Foster Carter, 1940, Butler and Perry, 1940).

The pleura may also be perforated by carcinoma of the lung or punctured during operations of needling of the chest for paracentesis. Bilateral pneumothorax is a post operative cause of death in thyroidectomy and tracheotomy and the recognition of the condition may be a life-saving measure (Keis, 1934, Barrie, 1940, Ackerman and Bucker, 1941). Oesophageal cancer or accidental perforation of the oesophagus during the passage of an oesophageal bougie or oesophagoscope may give rise to spontaneous pneumothorax. It sometimes complicates sub-phrenic abscess or carcinoma of the stomach and colon and it can result from incision of the parietal pleura during operations on the kidney.

The second group are those cases that develop a spontaneous pneumothorax as a result of perforation of the lung by manifest active pulmonary tuberculosis or as a complication of artificial pneumothorax therapy. Sita-Lumsden (1949) recorded its occurrence as a complication of artificial pneumoperitoneum therapy, the air tracking to the pleural cavity through a persistent pleuroperitoneal canal. A pneumothorax may, of course, be induced accidentally when the lower intercostal route is employed for the induction or maintenance of pneumoperitoneum.

The third group is where it occurs in the apparently healthy. The condition is then known as benign spontaneous pneumothorax. It is most frequent between the ages of 15 and 35 years. It is eight times more common in males than in females.

Rupture of the lung, with resultant pneumothorax, may occur in the

## AETIOLOGY

newborn following a normal birth (Ellison and Carabelli 1940). It used to be thought that benign spontaneous pneumothorax was the result of a sub-pleural tubercle rupturing into the pleura and it is possible that in rare instances this is the cause of the condition. It is quite evident however that if this occurred the pleura would be infected with tubercle bacilli and would immediately give an acute febrile reaction with the outpouring of fluid and raising of the sedimentation rate—a reaction which never occurs in benign spontaneous pneumothorax. Kjaergaard (1932) and Perry (1939) followed groups of cases and showed that it was rare for them subsequently to develop pulmonary tuberculosis which is in marked contrast to patients with pleural effusion. There is no doubt



FIG. 20.—Microphotograph of section of lung showing an intact emphysematous bulla with a ruptured one beside it. This caused a spontaneous pneumothorax.

that a vast majority of the cases result from the rupture of sub-pleural air vesicles. Such an air vesicle may be a result of compensatory emphysema from chronic pulmonary tuberculosis where the disease is quiescent. It is an infrequent complication of the focal emphysema of pneumoconiosis. The vesicles are sometimes of congenital origin but usually result from non-tuberculous scar tissue formation (Fig. 20) or chronic hypertrophic emphysema. The vesicles may be readily seen through a thoracoscope and have been described by Broek (1948). He found that generalized emphysema and localized bullous emphysema were the commonest causes next to these small apical bullae apical scars or tearing of the lung. Some cases resulted from either localized cystic lung or generalized polycystic disease and a few cases appeared to arise from areas on the surface of the lung resembling cuckoo spit. Spontaneous pneumothorax may be

complicated by mediastinal emphysema, which may spread to involve the pericardium and give rise to a pneumopericardium. The air may also spread to cause surgical emphysema of the head and neck and even the arm and chest wall.

### CLINICAL SYMPTOMS

The patient most often complains of pain in the chest on the side affected. The pain is usually of sudden onset so that the patient can record the exact moment and what he was doing at the time, rarely, the onset may be gradual with the patient unable to date it at all. The pain is generally in the side of the chest but sometimes it is situated in the shoulder or the back, and rarely, the patient complains of pain in the upper abdomen. When the pain is abdominal it may easily be mistaken for a perforated peptic ulcer. It most often occurs early in the morning on getting out of bed, but it sometimes occurs later in the day, or after a prolonged rest such as a visit to the cinema. The pain is followed by shortness of breath, especially on exertion, and sometimes the patient may be cyanosed. Among the less common initial symptoms haemoptysis is perhaps the most frequent, a little blood stained mucus being coughed up, probably arising from the lung at the point of rupture. Vomiting may occasionally occur as an early symptom. In a few cases the patient is greatly shocked as a result of the disease.

Physical signs will depend to a great extent on the amount of air in the pleural cavity. There is displacement of the trachea and apical beat away from the affected side of the chest, diminished movement and diminished tactile vocal fremitus, hyper-resonance, diminished or distant breath sounds, and absent voice sounds, and in some cases the coin sound elicited by placing a coin on the chest and tapping with another. The heart and trachea are not likely to be greatly displaced but all degrees of displacement may occur. In a left sided pneumothorax the trachea may lie under the right sternomastoid muscle and the apical beat be found on the right of the sternum. When this occurs the patient is likely to be cyanosed and orthopnoeic. In most cases the patient is afebrile but in those cases of tension pneumothorax which result from a valve like action of the pleural perforation there may be a slight rise of temperature and pulse rate, both of which are immediately relieved by removal of air.

The diagnosis can be established conclusively by x ray examination which will show air in the pleural cavity with usually no disease in the underlying lung. There is often congestion of the opposite lung from increased blood supply, which is always present and should not be mistaken for tuberculous infiltration. Christie and McIntosh (1936) showed that immediately the lung collapses the

As a result  
the blood  
always seen

in the radiographs, and also the rareness of cyanosis in this disease. If there is a tension pneumothorax a mediastinal herniation of pleura to the other side may occur and show itself as a semi circle in the other lung field with the hilum as the centre. Such a shadow is not infrequent in the course of artificial pneumothorax therapy. The sedimentation rate is always normal and in most cases there is no sputum for examination.

## DIAGNOSIS

### DIAGNOSIS

The history of the pain is sometimes misleading and the case arriving at hospital complaining of pain in the side or back is easily labelled rheumatism, lumbago, or fibrositis. When the pain is in the chest it may be labelled pleurisy, if, however, the condition is considered and skiagrams are taken, the diagnosis should not be missed. In cases where there is displacement of the mediastinum the diagnosis is easy because in these cases other signs are well marked. It may




FIG. 21.—Skiagram of a young man showing a single benign spontaneous pneumothorax of medium extent (The arrows indicate the lung periphery.)

however be difficult to differentiate, even radiologically a case of spontaneous pneumothorax from one with a very large emphysematous bulla. Congenital bronchial cysts may grow to a size that will give physical signs and radiological appearances superficially resembling a spontaneous pneumothorax. Such cysts are frequent in infants. A large sub-phrenic abscess containing gas is also liable to be confused with a spontaneous pneumothorax. In cases of bilateral spontaneous pneumothorax it is not possible to compare the signs on the two sides but the patient is usually seriously ill and there is thus no possibility of the condition being missed, skiagrams will establish the diagnosis. In relapsing or alternating pneumothorax the diagnosis is not so difficult as in the first attack, and often the patient makes the diagnosis himself as soon as the symptoms reappear.

## PROGNOSIS

The course of benign spontaneous pneumothorax is favourable in the majority of cases, few necropsies being recorded in the literature. The condition may however become recurrent or chronic, and 71 such cases were described by Brock in 1948. In most cases the lung expands in a period of six weeks, but in a few cases the pneumothorax may persist for months or years. Recurrence occurs in about 15 per cent of cases. When recurrence does occur the patient may have five, six, or even more attacks, and these may be on one or both sides.

## TREATMENT

Puncture wounds of the chest caused by a knife or sharp instrument seal quickly and a closed pneumothorax is then established. As a result, however, of trauma from jagged objects or gunshot wounds, a sucking pneumothorax may develop and become established under atmospheric pressure. This impedes the action of the heart and lungs, and it is most important to close the wound in the chest wall as soon as possible. A thick gauze pad may be fixed over the opening as first aid treatment. But on reaching hospital the wound should be excised and sutured as soon as possible.

The principles of treatment of closed pneumothorax are the same whether the lesion occurs in the apparently healthy, results from trauma or from any other cause. In milder cases the patient frequently does not seek advice immediately and when he does no treatment is required. In more severe cases rest in bed for one week should be sufficient to allow the perforation to heal and the air to start absorbing. Sanatorium treatment is unnecessary except in tuberculous patients. In the benign cases the patient should be warned that there is a certain risk of recurrence of the condition, this being greatest during the first year. Tension pneumothorax usually develops slowly although sometimes as an exacerbation two or three days after the onset of the original pneumothorax. The patient is greatly distressed, orthopnoeic, and cyanosed, but the condition may be immediately relieved by the insertion of a Foster-Carter needle attached to an under water drain. Pneumothorax occurring in cases of severe generalized emphysema or where bilateral pneumothoraces have occurred may also require the insertion of the Foster-Carter needle. Such patients should also receive oxygen therapy, either by means of bilateral nasal catheters, a nasal mask or oxygen tent. The most satisfactory method of treating troublesome chronic or relapsing and recurring pneumothorax is by means of pleurodesis by silver nitrate, 5-10 minims of 10 per cent solution is injected into the pleural cavity. A febrile reaction follows which usually lasts about five days and is accompanied by effusion of varying size. When it has subsided it is advisable to show that the pleural space is obliterated by demonstrating that it is not possible to induce an artificial pneumothorax.

In cases where a lobe is grossly abnormal, usually from congenital cystic formation excisional surgery may be the best form of therapy.

## BIBLIOGRAPHY

- Ackerman L. V. and Bucker E. M. (1941) Mediastinal Emphysema and Bilateral Pneumothorax following Radical Dissection of the Neck " *Arch Surg*, 43, 445

## BIBLIOGRAPHY

- Barrie H J (1940) Interstitial Emphysema and Spontaneous Pneumothorax after Operations on the Neck *Lancet* 1, 996
- Brock R C (1948) Recurrent and Chronic Spontaneous Pneumothorax *Thorax* 3 88
- Butler E C B and Perry K M A (1940) Spontaneous Pneumothorax complicating Staphylococcal Septicaemia *Lancet* 2 619
- Christie R V and McIntosh C A (1936) The Lung Volume and Respiratory Exchange after Pneumothorax *Quart J Med* 5 445
- Collis J L and Foster Carter A F (1940) Spontaneous Pneumothorax and Staphylococcal Lung Abscess in an Infant *Lancet* 1 875
- Ellison R T and Carabelli A A (1940) Tense Pneumothorax in an Infant Successful  
*J* 10 644  
 von Spitzeunarbe-  
 mphysem *Z klin*
- Hayashi J (1914) Ueber todtlichen pneumothorax durch Infarkt und Emphysem *Franz-  
 furter Z Path* 16 1
- Leach J E (1945) Pneumothorax in Young Adult Males Descriptive Statistics in 126  
 Cases *Arch intern Med* 76 264
- Oeschli W R and Miles S H (1934) Simultaneous Bilateral Spontaneous Pneumothorax  
*Amer Rev Tuberc* 30 239
- Ornstein G G and Lercher L (1941) Spontaneous Pneumothorax in Apparently Healthy  
 Individuals Clinical study of 58 cases with discussion of pathogenesis *Quart Bull  
 Seaw Hosp* 7 149
- Perry K M A (1939) On Spontaneous Pneumothorax *Quart J Med* 8 1
- Schneider L and Reissmann I (1945) Idiopathic Spontaneous Pneumothorax History  
 of 100 unselected Cases *Radiology* 44 485
- Sita Lumsden E G (1949) Spontaneous Pneumothorax complicating Artificial Pneumo-  
 peritoneum *Thorax* 4 147

## CHAPTER 5

### CALCIFICATION OF THE PLEURA

I C LODGE PATCH and KENNETH M A PERRY

#### INCIDENCE

SINCE the days of Aristotle pathological calcification of the pleura has been recognized, and in the 18th century Morgagni (1682-1771) drew attention to the condition. Since radiological examination of the chest has become common it has been realized that the condition is not infrequent. If the incidence is estimated by radiography alone it will be found in unselected cases to be less than 0.2 per cent.

In a series of 1,845 cases Ulrich (1930) found a percentage of 0.87, Pritchard (1923) in 7,000 cases found 0.12 per cent, in Taylor and Schwartz's (1937) 10,000 cases there was 0.1 per cent, and 0.18 per cent were found in 15,000 cases by Floyd and Hepburn (1939). In contrast, however, Behrendt (1930), in reviewing ex-servicemen found an incidence of 1.2 per cent in 886 cases. Even allowing for the greater risk of injury to men especially in wartime they show pleural calcification much more commonly than women. Thus Head (1934) found it in 47 men compared with 8 women and Odessky and Khioner (1932) in 11 men and 2 women.

#### AETIOLOGY

Most cases result either from trauma, where there has been haemorrhage into the pleura cavity, or from inflammation of the pleura which may have resulted either in a serous pleurisy or an empyema (Fig 24). It occurs in some cases of spontaneous pneumothorax, probably as a result of bleeding into the pleural space (Fig 22).

TABLE  
CAUSE OF CALCIFICATION IN 70 CASES (HEAD 1934)

Cause	Number of cases	Percentage
Gunshot wounds	17	24.3
Active tuberculosis	8	11.4
Inactive tuberculosis	3	4.3
Pleural effusion	12	17.1
Dry pleurisy	7	10.0
Respiratory disease—		
influenza	1	
pneumonia	1	14.3
pneumonia and empyema	3	
chronic lung disease	4	
empyema	1	
No respiratory disease	13	18.6




FIG. 22—A man aged 50 years with a right chronic spontaneous pneumothorax and calcified pleura

Case (1933) and 24

Case

Case

Vest (1906)

occurs as  
and Tonkin  
and Church

and Ellis (1950) mention cystic changes but there is no recorded case of calcification arising in this condition

Head (1934) collected 70 cases of pleural calcification from the literature. He lists the causes as shown in the Table

### PATHOLOGY

Calcification is commonly unilateral and is related to the aetiological cause: it is only bilateral in the idiopathic cases. The calcium is deposited between the layers of the pleura and while it is often loosely attached to the pleural surface it is in fact intrapleural (Stott and Cotton-Cornwall (1933), Head (1933), Goullioud (1906), Boswell and Dredge (1940)). In extensive calcification it may appear as a crust over the pleura (Hill 1933). The mechanism by which the calcium is deposited is uncertain. Head and Boswell and Dredge (1940) consider that the pleura acts as a dialysing membrane and the calcium is deposited as a superficial crust as the last of the fluid is absorbed. But Pritchard (1923) and Hsu and Wang (1936) suggest that the fluid or blood is encapsulated by thickened pleura and slowly absorbed. The nutrition of the pleura is impaired and hyaline degeneration ensues in this degenerate tissue so that calcium is then deposited



## CALCIFICATION OF THE PLEURA



FIG 23 — Idiopathic calcification of the pleura in a man aged 50 years who had an abscess pointing on the surface




FIG 24 — Calcified pleura resulting from a pleural effusion in childhood

While Pritchard (1923) and Head (1933) did not find local or remote evidence of tuberculosis, Taylor and Schwartz (1937) regarded calcification of the pleura as analogous to the calcification of tuberculosis in any other organ. Haematogenous tuberculosis produces a caseous pleuritis, with or without effusion and subsequent calcification. In three necropsies they found the association of calcification and caseation in the pleural layer. Haematogenous tuberculosis was found elsewhere in all three subjects. It is however quite clear that tuberculosis cannot be the cause in all cases since calcification is found in non tuberculous patients.

The size of the calcium deposits may vary from small spots (Hill 1933) to single plaques (Pritchard, 1923) and multiple plaques (Pallasse and Martinon 1937) and extensive shells (Hill 1933, and West 1906).

Empyema is commonly found in association with calcification of the pleura (Pritchard, 1923) and Taylor and Schwartz (1937) regard this as a common complication when the pleural layers are not adherent. It is probable that the plaques erode the lung tissue and infection passes from a bronchus. Where the pleura is adherent small abscesses may develop which may point on the surface.

### CLINICAL FEATURES

There are usually no symptoms in association with the condition but if it is extensive, shortness of breath may develop and cough is sometimes associated. If the plaques give rise to pus formation there may be fever and an abscess may appear in the chest wall (Fig 23). On examination the chest is often fixed and flattened and shows an impaired percussion note and diminished air entry (Beverley 1937 1942, Hsu and Wang 1936, Passalacqua and Farias 1939, Stott and Cotton Cornwall 1933). There may be scoliosis (Hill 1933).

### TREATMENT

In the absence of symptoms no treatment is indicated. If the calcium plaque or sheet is loose or causing symptoms it may be removed (Head 1933 and Goulloud, 1906). If there is marked shortness of breath and the chest wall is fixed decortication of the lung may be advisable.

### BIBLIOGRAPHY

- Behrendt T (1930) Verkalkungen von Pleuraschwarten. *Brit. Klin. d. Tuberk.* 76 129  
 Beverley S S (1936) Calcareous Pleurisy. *Med. Bull. Vet. Adm'n.* 13 277 278  
 — (1942) *Ibid.* 19, 357-358  
 Boswell J R and Dredge T E (1940) Calcification of the Pleura. Two Cases. *Med. Bull. Vet. Adm'n.* 17, 75 77  
 Church P E and Ellis A R P (1950) "Cystic Pulmonary Fibrosis in Generalised Scleroderma." *Lancet* 1 392  
 Dwight T (1909) Extensive Calcification of Pleuritic Exudation causing Curvature of the Spine. *Boston med. surg. J.* 160  
 Floyd C and Hepburn R H (1939) Calcification of the Pleura. *Amer. rev. Tuberc.* 40 215-223  
 Gandy C and Baize P (1930) Calcifications pleurales. *Pr. mèd.* 38 877 880  
 Goulloud M (1906) Résection d'une large plaque calcaire d'origine pleurale. *Lyon mèd.* 107, 350-353  
 Head J (1934) Calcification of the Pleura. *Ann. Intern. Med.* 7, 1295 1307

# CALCIFICATION OF THE PLEURA

- Hsu, C, and Wang S (1936) "Calcification of the Pleura report of 2 cases" *Chin med J*, 50, 1051-1056
- Lloyd, W E, and Tonkin, A J (1948) "Pulmonary Fibrosis in Generalised Scleroderma Review of the Literature and report of 4 further cases" *Thorax*, 3, 241.
- Odessky, I, and Klioner, I, (1932) "Zur frage der Pleura-verkalkung" *Z Tuberk*, 66, 136-142
- Pallasse, and Martinon (1937) "Calcifications pleurales en Mosaïque" *Lyon méd*, 159, 117-120
- Passalacqua, H A, and Farias, L (1939) "Calcification pleural bilateral y tuberculosis pulmonar" *Prensa med argent*, 26, 1100-1102
- Pritchard J S (1923) "Some Interesting Cases of Calcareous Degeneration found in the Thorax" *Arch intern Med*, 32, 259-282
- Stein H M (1932) "Cholesterol-thorax in Tuberculosis (Cholesterol Pleurisy) Report of a case" *Arch Intern med* 49, 421-428
- Stott L B, and Cotton-Cornwall, V (1933) "Case of Pathological Calcification" *Lancet*, 2, 755
- Taylor, H K, and Schwartz, S (1937) "Calcification of the Pleura" *Quart Bull Seaview Hosp*, 3, 65-75
- Ulrich K (1930) "Über Verkalkungen von Pleuraschwarten Rontgenpraxis" 2, 212
- West, S (1906) "A Case in which the Pleura Contained Several Pints of Calcareous Mortar-like Fluid" *Trans clin Soc Lond*, 39, 42-45
- Young F H (1925) "A Case of Calcification of the Pleura of an Unusual Type" *Lancet* 2, 1224 1225

## CHAPTER 6

### TRANSIENT PULMONARY INFILTRATION WITH EOSINOPHILIA

JAMES L. LIVINGSTONE

#### DEFINITION

TRANSIENT pulmonary infiltration with eosinophilia (Löfller's syndrome) is defined as a group of clinical conditions, varying in duration and severity, in which transitory pulmonary opacities and eosinophilia in the peripheral blood are cardinal features—the histological picture suggests an antigen-antibody tissue reaction.

#### HISTORY

In 1932, Löfller described 5 cases and in 1936 a series of 51 cases with minimal symptoms, transitory pulmonary infiltrations and eosinophilia up to 22 per cent without leucocytosis. Subsequently, Löfller and Maier (1943) found ascaris infestation in a proportion of cases, particularly in soldiers of the Swiss army. Von Meyenberg (1942) reported post mortem examinations of 4 soldiers in whom broncho-pneumonic patches were found with infiltration of eosinophilic cells. Two of these patients had ascaris worms in the intestine.

Since then numerous papers have appeared from all over the world describing groups of cases showing eosinophilia and transient pulmonary infiltrations; these fall into 4 groups: (1) Infestation with parasites, for example *Ascaris*, *Necator americanus*. (2) Toxic reactions from drugs, from serum injections, and sensitivity to foods and plants. (3) Bacterial infections, particularly in patients with asthma or other allergic conditions, in rheumatic fever and brucellosis. (4) Tropical eosinophilia (pulmonary eosinophilosis) occurring chiefly in India and Ceylon. In 1940, Fridolf Møller and Barton reported 175 cases and in 1943 Weingarten reported 81 cases and found that arsenic was specific therapy.

Rich and Gregory (1943) produced polyarteritis nodosa in rabbits by sensitization to horse serum and they found a specific histological picture of fibrinoid necrosis of connective tissue occurring as a result of this antigen-antibody reaction. Harkavy (1943) and Bergstrand (1946) in post-mortem material in asthmatic patients reported arterial lesions, fibrinoid necrosis and rheumatic granulomas essentially similar to the lesions in rabbits produced experimentally by Rich and Gregory.

#### AETIOLOGY

The possible causes of this condition may be divided into 4 groups: (1) Infection with parasites, for example *Ascaris lumbricoides*, *N. americanus*, *Ankylostoma braziliense*, *Fasciola hepatica* and *Trichinella spiralis*. Acari (mites), *Entamoeba histolytica* and microfilaria have been suggested as possible causes. (2) Toxic or allergic reactions to: (a) drugs such as the sulphonamides, organic arsenicals, and thiouracil, (b) injections of foreign sera, (c) foods such as egg and tomato.

and (d) pollens or certain plants such as privet (*Ligustrum*) (3) Bacterial infections of the respiratory tract, particularly in patients with an allergic diathesis (such as asthma, hay fever, and eczema) Acute rheumatism and brucella infections have been reported (4) Tropical eosinophilia (pulmonary eosinophilosis) This appears to be a clinical entity, in that although the cause is unknown, arsenic is a specific remedy

### MORBID ANATOMY

In 1942, von Meyenberg reported post-mortem findings in 4 soldiers, 3 of whom died suddenly as a result of accidents and 1 from tetanus He described consolidations in the lungs without fibrin formation, with eosinophilic infiltration, giant cells in the alveoli and thrombophlebitis and rheumatic granulomas in the interstitial tissue Eosinophilic infiltrations were present in the liver and elsewhere Two of these men had ascaris worms in the intestine Harkavy (1943) in post-mortem examination of four asthmatics found eosinophilic infiltration of the inter-alveolar septa with arterial lesions, varying from simple intimal thickening up to acute necrosis of the arterial wall with thrombosis Bergstrand (1946) described the post-mortem findings in four asthmatics with infection of the respiratory tract, transient lung infiltrations and eosinophilia in the blood Besides polyarteritis nodosa lesions in various stages, he found rheumatic granulomas, as described by Aschoff, in the interstitial tissues of the lungs; there was oedema and swelling of the collagen fibres which split up into fine fibrils, with large fibroblasts which sometimes form multinucleated giant cells eosinophil cells were frequent, and considerable formation of fibrous tissue was present Necrosis in the centre of the granulomas was frequently found, and conglomerate granulomas were seen in the interlobar septa which were visible to the naked eye as greyish white streaks in the lungs These appearances are essentially identical with those seen in rheumatic pneumonia and in subcutaneous rheumatic nodules

### CLINICAL FEATURES

The clinical features of this condition may be considered in 3 groups according to the aetiology

**Helminth and parasitic infestations, serum and drug reactions**

Symptoms may be entirely absent or minimal with lassitude, cough, chest pains, slight fever and occasionally sputum, which is rarely blood stained Physical signs are either absent or confined to a patch of fine râles

Skiagrams of the chest show varying types of nodular opacities which disappear in 10-21 days

The blood shows a moderate leucocytosis with an eosinophilia usually below 20 per cent


A search for parasites or ova should be made and treatment obviously depends upon the cause

**Bacterial infection, especially of the respiratory tract**

In this group there is a history of previous allergic conditions (asthma, eczema,

# CLINICAL FEATURES

Fig 25 — Resolving pneumonia in left mid and lower zones



9 1 45



Fig 26 Further resolution at left base but fresh faint round opacity in right fourth space

20 1 45

TRANSIENT PULMONARY INFILTRATION WITH EOSINOPHILIA





FIG 27 — Extensive opacity in right mid and lower zones shown on the lateral view to be in the anterior segment of the upper lobe and medial basal segment of the lower lobe. Patient had been free of symptoms for three days

5 2 45

FIG 28 — Some clearing of the opacity



13 2 45

# CLINICAL FEATURES

FIG. 29 — Considerable  
increase in the cardiac  
shadow with pleural  
reaction at both bases



FIG. 30 Still gross  
cardiac enlargement  
with clearing of the  
pleurisy at the bases



# TRANSIENT PULMONARY INFILTRATION WITH EOSINOPHILIA




FIG 31—Showing rise and fall in eosinophils and fluctuation of leucocyte count of Case FE

## Case Report FE

Female aged 45 years

No...

ate

attacks gradually  
November 1944  
15 oz (450 ml)  
bed 3 weeks but  
January 15 1945  
Asthma every night  
January 19 1945  
clubbing BP 140/80  
phs Leucocytosis 62  
February 2 1945

Signs of consolidation left lower lobe

Sputum purulent

sputum increased to  
hema persisted in

sputum thick and purulent

ra on 30 Early  
lis mainly eosino

frequent cough  
us epecally from left  
on obtained showed many pus cells  
ls organisms scanty on culture

Cold agglutins normal Car

very fit breathing well and doing

reg on BP  
gns suggested

when  
upper  
numeral  
a few  
Febr  
barson  
Febr  
skipp n  
March 5 1945  
110/98 Feeble  
dilated heart rath

# CLINICAL FEATURES

March 15 1945	Temp 100°	on top 100	Anaemia negative	Liver enlarged	Spleen thick
"	"	"	"	"	d
"	"	"	"	"	t
per cent					
March 26 1945	Improving	good diuresis with Neptal			
March 30 1945	Patient discharged herself from hospital			She remained bed ridden in	
cardiac failure					
July 1945	Died	No necropsy allowed			
<i>Investigations</i>					
Sputum	few organisms	No tubercle bacilli	many Charcot	Leyden crystals and eosinophils	
Pus cells	were numerous				
Stools	no ova or parasites				
Casoni test	negative				
				normality	
				30 c mm	

urticaria and rhinitis) in more than half of the cases, and a family history of allergy is common. The clinical picture is usually that of recurrent attacks of bronchitis with severe asthma. Pleural effusions are common and may be bilateral, with many eosinophil cells in the fluid, the proportion of these cells may differ markedly in the fluid from one side as compared with the other. The heart may dilate, and resume its normal size at a later date. The spleen and liver are often enlarged, the lymph nodes less frequently. Nodules may appear in the subcutaneous tissues, muscles or tendon sheaths, similar to large "rheumatic nodules" in rheumatic fever. Albuminuria with red cells and casts may occur periodically or persistently, and hypertension may supervene with terminal cardiac or renal failure.

There is a leucocytosis from 20,000 to 60,000 per cubic millimetre, with an eosinophilia up to 84 per cent during the acute phases, falling to normal between attacks.

Radiographic examination shows pulmonary opacities, rather large and asymmetrical, which persist for days or weeks and then disappear completely. These opacities do not appear to be areas of segmental collapse, and come and go without definite relation to the acute attacks of infection (Figs 25-30).

The majority of recorded cases have died following an illness of 3-15 months after the onset of the eosinophilia. A number of patients are still alive however, and apparently recovering after observation for 2-4 years. It is obvious that this group, if not identical with polyarteritis nodosa is very closely related to that condition.

There is no specific treatment. Antihistamine drugs may be tried both during the attack and prophylactically.

## Tropical eosinophilia (pulmonary eosinophilosis)

This condition appears to be a clinical entity, although the aetiology is unknown. It occurs chiefly in India and Ceylon, although sporadic cases have been reported in America, North Africa, China, Australia, the Pacific Islands, Egypt and elsewhere.

Viswanathan (1948) reviews the subject and reports the post mortem findings in one case in which there were nodules in the lung showing eosinophilic infiltration and a similar picture to that described by Bergstrand

### *Clinical features*

*Acute type*—The onset is sudden with fever from 102° to 104° F with cough and rapid respirations simulating acute bronchiolitis. The signs are those of acute bronchitis and occasionally of patchy broncho pneumonia. Radiographic examination shows a patchy mottling confined to one part of the lung, or uniformly distributed in all fields. There is a leucocytosis ranging from 12 000 to 80,000 per cubic millimetre with from 20 to 80 per cent of eosinophils.

If untreated, about a quarter of the cases clear up spontaneously in from 3 to 6 weeks, but the remainder become chronic cases.

*Chronic type*—The onset is insidious with malaise, lassitude, anorexia and low grade fever from 99° to 101° F. Cough may be noticed at the onset or more commonly after a week and is unproductive, paroxysmal and often most distressing at night. Small blobs of viscid sputum are expectorated with difficulty and there is marked dyspnoea on coughing or on exertion, rather than typical nocturnal asthma. The spleen is palpable in about half of the cases, and occasionally there is hepatomegaly and enlarged lymph glands. Radiological opacities are seen during the earlier stages of the disease, and occasionally persist for months, coarse mottling may be present throughout the lungs or confined to one part, cavitation may be seen but is uncommon. In the later stages, heavy hilar shadows may be the only abnormality.

The erythrocyte sedimentation rate is usually raised. Viswanathan and

syphilis. The reaction becomes negative after treatment with arsenic.

Weingarten (1943) reported 81 cases and observed, when treating one patient suffering from syphilis with neoarsphenamine, that there was a rapid improvement in the symptoms and in the eosinophilia. He then discovered that arsenic was a specific cure.

The usual course of neoarsphenamine is 0.15 gramme followed by 7 weekly injections of 0.3 gramme. After the first dose there is usually an exacerbation of symptoms with an increase in the eosinophilia. The clinical symptoms and signs quickly disappear but the blood does not return to normal for several weeks after the completion of treatment.

In real asthma or in helminth infestation, organic arsenic does not improve the condition nor is there the preliminary increase in eosinophilia which are points in differential diagnosis.

### BIBLIOGRAPHY

- D. Abrera, V. St. E. and Stork, K. G. (1946) Serological Reactions in Tropical Eosinophilia.  
*Ind. med. Gaz.* 81, 282.

# BIBLIOGRAPHY

- Bergstrand H (1946) Morphological Equivalents in Polyarthritis Rheumatica Periarthritis  
Nodosa Transient Eosinophilic Infiltration of the Lung and Other Allergic Syndromes  
*J Path Bact* 58 399
- Erismodi Möller C and Barton R M (1940) A Pseudo-tuberculous Condition Associated  
with Eosinophilia *Ind med Ga* 75, 607
- Gregory J E and Rich A R (1946) "The Experimental Production of Anaphylactic Pul-  
monary Lesions with the Basic Characteristics of Rheumatic Pneumonitis" *Bull Johns  
Hopk Hosp* 78, 1
- Harkavy J (1943) Vascular Allergy III *J Allergy* 14, 507
- Löffler W (1932) Zur Differential Diagnose der Lungen infiltrierungen I Frühinfiltrate  
unter besonderer Berücksichtigung der Rückbildungszeiten *Beitr klin Tuberk* 79, 365
- (1936) "Die flüchtigen Lungeninfiltrate mit Eosinophilie" *Schweiz med Wschr* 66,  
1069
- Maier C (1943) "Das flüchtige eosinophile Lungeninfiltrat zusammenfassende Ergebnisse  
von über 100 Beobachtungen" *Helv med Acta* 10 95
- Menon I G K (1945) Tropical Eosinophilia *Ind med Ga* 80, 24
- von Meyenberg H (1942) "Das eosinophile Lungeninfiltrat" *Schweiz med Wschr* 72, 809
- Rich A R (1942) The Role of Hypersensitivity in Periarthritis Nodosa *Bull Johns  
Hopk Hosp* 71, 123
- and Gregory J E (1942) "The Experimental Demonstration that Periarthritis Nodosa  
is a Manifestation of Hypersensitivity" *Ibid* 72 65
- (1943) Experimental Evidence that Lesions with the Basic Characteristics of  
Rheumatic Carditis can Result from Anaphylactic Hypersensitivity *Ibid* 73, 230
- Viswanathan R (1948) Pulmonary Eosinophilosis *Quart J Med NS* 17, 257
- and Natarajan B (1945) Cold Agglutination in Tropical Eosinophilia *Lancet*  
I, 148
- Weingarten R J (1943) Tropical Eosinophilia *Lancet*, I 103

## CHAPTER 7

### POLYARTERITIS NODOSA (PERIARTERITIS NODOSA)

JAMES L. LIVINGSTONE

#### DEFINITION

POLYARTERITIS nodosa is a syndrome of unknown aetiology, marked by inflammatory reactions of the collagen connective tissue, particularly in the walls of arteries, and associated with a prolonged febrile course and multiple lesions throughout the body and commonly ending fatally

#### HISTORY

Kussmaul and Maier in 1866 gave the original description of the condition and correlated the clinical symptoms with the lesions found at autopsy. Since then details of some 400 further cases have been published, particularly during the past 15 years. *although the majority have been discovered at autopsy, an increasing number have been diagnosed in life on clinical grounds and confirmed by biopsy of nodules.* Page (1947) records a case in which peritoneoscopy confirmed the clinical diagnosis, the nodules being seen on the arteries of the stomach and colon. During recent years an increasing number of proved cases with recovery have been recorded.

#### MORBID ANATOMY

The essential lesion is an acute fibrinoid necrosis of the collagen tissue, especially in the walls of the smaller arteries and arterioles. There is oedema round the necrosed collagen, with infiltration of endothelioid cells and lymphocytes, and *eosinophils are frequently seen in large numbers.* At a later stage the lesion is identical with an Aschoff's node, still later, fibroblastic proliferation occurs, and when healing is complete a small fibrous scar remains. The arterial lesions start in the media, but there is always a proliferation of the intima at the site, and intravascular thrombosis, rupture of the arterial wall and aneurysm formation are all common. Lesions in different stages of development may be seen at the same time, and have been found in practically every organ and tissue of the body. The kidneys, liver, lungs, spleen, heart, small intestine, pancreas, suprarenal glands and uterus are most frequently affected, and it is relatively uncommon to find nodules in the superficial arteries which are available for clinical diagnosis.

In the kidney the afferent arterioles to the glomeruli are frequently involved, causing hyalinization of the glomeruli, indeed, changes identical with those found in malignant hypertension may be found, and many patients with polyarteritis nodosa die from renal failure with hypertension and uraemia.

In biopsy specimens particularly, it may be difficult to detect the typical arterial involvement unless serial sections are cut, and reticulum stains should be used. The degree of eosinophilic infiltration is very variable (Figs 32-35)

FIG 32 — Acute polyarteritis nodosa (low power) Artery in striated muscle Fibrinoid necrosis and cellular infiltration involving a segment of the vessel wall Haematoxylin and eosin  $\times 105$  (reduced by  $\frac{1}{2}$  on reproduction) (By courtesy of Professor H A Magnus)



FIG 33 — High power of Fig 32 showing acute fibrinoid necrosis of media with cellular infiltration of adventitia and periarterial tissues Haematoxylin and eosin  $\times 370$  (reduced by  $\frac{1}{2}$  on reproduction) (By courtesy of Professor H A Magnus)

*History* Female, aged 41 years. Six months history of loss of weight and swelling of ankles. Admitted to hospital with irregular pyrexia to 102° with purpura, acne rosacea and splenomegaly. Blood pressure 140/105. Haemoglobin 56 per cent with a normal leucocyte count. She developed pain in the chest from a pulmonary infarct and died in coma after sixteen days (Figs 32-33).

### AETIOLOGY

The cause of polyarteritis nodosa is as yet quite unknown. Theories postulating a specific bacterium, virus or chemical toxin have been without scientific proof.


The specific histological picture of an acute reaction in the collagen connective tissue is in many ways similar to the histology in acute rheumatism and is also found in lupus erythematosus disseminatus, dermatomyositis, scleroderma and certain chronic types of transient pulmonary infiltrations with eosinophilia. The

# POLYARTERITIS NODOSA (PERIARTERITIS NODOSA)



FIG 34 — Healed stage of polyarteritis nodosa in small renal artery (low power), showing intimal thickening and segmental fibrosis with aneurysmal bulging of media. Picro-Mallory,  $\times 70$  (reduced by  $\frac{1}{2}$  on reproduction) (By courtesy of Professor H. A. Magnus)

FIG 35 — High power of Fig 34. Picro-Mallory,  $\times 140$  (reduced by  $\frac{1}{2}$  on reproduction) (By courtesy of Professor H. A. Magnus)



History  
infarction  
per cubic  
terminal  
(Figs 34,

theory of an allergic antigen-antibody reaction has been postulated but until recent years there was no known method of producing similar lesions in animals Rich (1942), however, noted a marked increase in the incidence of polyarteritis nodosa in the post mortem department of the Johns Hopkins Hospital from 1935, when sulphonamides were introduced, as compared to the previous 25 years. He found that a high proportion of these patients had suffered from serum sickness (from Felton's anti-pneumococcal serum) or from sulphonamide drug reactions in life.

## CLINICAL MANIFESTATIONS

He then sensitized rabbits by giving them a very large dose of horse serum intravenously (corresponding to 200-300 millilitres in humans) and noted that approximately 75 per cent developed serum sickness 7-10 days later, as evidenced by pyrexia and flushing of the ears. By the fourteenth day all the rabbits were sensitized to horse serum as shown by the dermal skin tests. He then gave these rabbits varying doses of horse serum intravenously, and found that while some animals died immediately of anaphylactic shock, a number survived. These were killed at varying times and showed typical polyarteritis nodosa in different stages of development. The heart muscle and valves showed reactions identical with acute rheumatic myocarditis and endocarditis in man. Rich's work has been substantiated by others.

The inference is that at any rate some cases of polyarteritis nodosa are caused by an antigen-antibody reaction. This does not explain however why polyarteritis nodosa is much more common in males and why it occurs in persons who have never shown any clinical evidence of allergy or toxic reactions to drugs or sera. As a working hypothesis, however, Rich's investigations are of considerable significance.

## CLINICAL MANIFESTATIONS

**General**  
Males are affected at least twice as often as females. Although a case has been reported in an infant, the common age is from 25 to 55 years. The onset may be abrupt or insidious, with muscular aches and pains, irregular pyrexia and tachycardia, with progressive anorexia, loss of weight, and weakness as the most prominent symptoms. Muscular tenderness and splenomegaly are common. Tender nodules may be found in the subcutaneous tissues or muscles, and occasionally a tender thickening is felt in a superficial artery, such as the temporal artery, which of course suggests the diagnosis. The retinal vessels may show localized areas of thickening, but more frequently exudates and retinal detachments are seen.

Remissions of some weeks' duration may occur in the course of the disease, followed by relapses with involvement of other areas. Pathological investigations commonly show a moderate secondary anaemia, with a polymorphonuclear leucocytosis of from 15,000 to 30,000 per cubic millimetre, and moderate eosinophilia is not uncommon. The erythrocyte sedimentation rate is usually high. Albuminuria with persistent red cells in the urine is frequent, and casts of various types are occasionally present. The blood urea may be raised, and this may be very high as a terminal event. Blood cultures are always negative.

### Local

Local manifestations are often widespread and bizarre, with fresh lesions occurring in one region and healing in another.

**Abdominal**—Symptoms vary from minor colicky pains to what is apparently an acute abdominal emergency, when extensive mesenteric thromboses may have occurred. Epigastric discomfort is common, as is anorexia and vomiting. Haematemesis or melaena may be presenting features. There may be transient



abdominal tenderness and the liver and spleen are often enlarged. Occult blood may be found in the stools, without clinical symptoms.

*Pulmonary*—Pleuritic pain, cough and haemoptysis are common, transient physical signs may be found in the chest. Pleural effusion is rare. Skiagrams may show transient opacities suggestive of infarctions.

*Renal*—Attacks of lumbar pain and haematuria are among the most frequent renal symptoms. Transient albuminuria and the passage of red cells in the urine may be observed without obvious cause. Hypertension with uraemia is a common termination of the disease, and a proportion of such cases are wrongly labelled "nephritis".

*Cardiac*—Tachycardia, dyspnoea and praecordial pain are common symptoms, the heart may be enlarged and typical myocardial infarction may occur. Transient pericardial friction may be observed. As renal involvement is so prevalent, the presenting features may be hypertension and cardiac hypertrophy.

At post-mortem examination the heart is frequently involved even when symptoms have been minimal during life.

There is nothing characteristic in the electrocardiographic tracings, although transient minor abnormalities may be seen and gross myocardial infarction is not rare.

*Neurological*—Recurrent severe headaches, painful stiffness of the neck and peripheral neuritis are perhaps the most common neurological symptoms. The cerebral vessels are less often involved than elsewhere, but focal lesions occur in any part of the nervous system. Hypertensive encephalopathy may be a feature in the later stages.

*Locomotor*—Transient arthritis, myalgia and tender nodules here and there about the body are frequent manifestations.

## DIAGNOSIS

The clinical picture of a chronic febrile wasting disease, with disseminated lesions about the body, is characteristic if polyarteritis nodosa is kept in mind as a possibility.

The following conditions have to be considered in the differential diagnosis.

*Subacute bacterial endocarditis*—In this condition the Osler's nodes, petechiae and cardiac murmurs, together with a positive blood culture, are diagnostic. Hypertension is rare in this disease.

*Tuberculosis chronic disseminated or renal*—In this the skiagram of the chest may show infiltration or enlarged mediastinal glands; a polymorphonuclear leucocytosis is uncommon and *Mycobacterium tuberculosis* can generally be recovered by direct smear or culture from gastric secretions, urine or blood.

*Sarcoidosis*—This condition perhaps most closely simulates polyarteritis

in the lung fields, with enlarged lymphatic glands, but there may be no lesion, or failing these, a biopsy of liver or tonsil may show the characteristic histology.

# DIAGNOSIS

FIG 36 — Temporal or giant cell arteritis thickening composed of cellular fibrous tissue with mucoid material between the cells. Segments of the internal elastic lamina are destroyed with fragmentation of remaining portions. The whole circumference of the media is involved in varying degree with areas of coagulative necrosis and heavy cellular infiltration. Multinucleated giant cells of the foreign body and Langhans type are seen in relation to the fragmented internal elastic lamina. The adventitia shows a cellular infiltration of lesser extent. Haematoxylin and eosin  $\times 39$  (reduced by  $\frac{1}{2}$  on reproduction) (By courtesy of Drs B S Cardell and T Hanley and the Editor of J Path Bact\*)



FIG 37 — High power of Fig 36 showing outer zone of intima and media. Cellular infiltration and giant cells. Haematoxylin and eosin  $\times 175$  (reduced by  $\frac{1}{2}$  on reproduction) (By courtesy of Drs B S Cardell and T Hanley and the Editor of J Path Bact\*)



**History** Male aged 68 years. Myocardial infarction when aged 55 years and 61 years. Symptoms began four months before death with generalized pains in neck, shoulders and calves and with anorexia and constipation. After this he developed excruciating pain in the right neck and temporo-parietal region followed in a few days by pain in the left temporal region. After eight weeks he complained of diplopia followed in a few days by blindness of the left eye with occlusion of the central retinal artery. He became completely blind with similar changes in the right eye a few weeks later. He later developed dysphagia and dysarthria and died about sixteen weeks after the onset of symptoms. Presenting clinical features were low grade pyrexia, tender thickening of the temporal arteries and a high erythrocyte sedimentation rate.

## CHAPTER 8

### ASTHMA

E R BOLAND

#### DEFINITION

ASTHMA is the name given to difficulty in breathing of a paroxysmal wheezing type, which is apparently due to *diminution in the bronchial lumen* and may be caused by bronchial obstruction, muscular spasm, vascular engorgement, oedema or increased secretion

It is, therefore, the name for a symptom produced by widely different conditions, and not strictly for a disease. The term is used, for example, in connexion with *cardiac disease*, in which case it would appear to be caused by temporary engorgement of the lungs when the left ventricular output does not keep pace with the right ventricular input. It is also used in connexion with the dyspnoea of advanced renal disease, in which the causes may be partly cardiac and partly metabolic, and it can be used in connexion with the postural dyspnoea produced by the pressure of mediastinal tumours. In this section, however, it is proposed to deal only with what is often termed bronchial asthma, which is paroxysmal in nature and in which, between paroxysms, the patient is subjectively and objectively in good health.

Compilers of text-books of general medicine are often at a loss as to the proper section for allocation to asthma. It is variously allotted to the sections on diseases of allergy, diseases of unknown origin, diseases of metabolism, diseases of the respiratory tract and disorders of the psyche—all with more or less justice. It can, in the absence of any general aetiological certainty, be justly dealt with in a text-book on respiratory diseases, since, in a condition where so much else is uncertain, it is at least incontrovertible that most of the symptoms and complications of asthma are respiratory. Although more tons of paper and more gallons of ink have probably been expended on explaining the aetiology of asthma than have been expended on any other common condition, we still know little of the true causation of asthma. This is a statement which would not be accepted by allergists, who have few doubts either about the cause or the treatment of bronchial asthma. Thus Unger (1945), in his text book, *Bronchial Asthma*, states that, the words bronchial asthma are to be employed to mean Allergic Bronchial Asthma, he also states that therapy is now based on three fundamental principles (1) the discovery, if possible, of the causes of the attacks, (2) the elimination of the cause, or (3) the desensitization of the patient if elimination is impossible.

This simple conception of asthma and its treatment is to be found in many text-books, and is the basis of most of the treatment by allergists, it is believed in by many practitioners and even by many patients, but as a conception it is over-simplified and the treatment is over-rated.

## ALLERGY IN ASTHMA

### ALLERGY IN ASTHMA

Allergy is a vast subject and there is no space here to deal with it in detail, but the term allergy meaning altered reactivity was coined by Von Pirquet in 1906 to describe the changes in tissue reaction occurring in some cases as the result of initial injections described first by von Behring, Richet and others. This term has been used to describe the sensitization reactions of man and it includes hypersensitivity, idiosyncrasy, sensitization and anaphylaxis.

The allergic theory of asthma is that certain individuals are or become sensitized to substances which are not normally harmful to normal individuals. These substances are called allergens and most of them contain protein which is regarded as most probably the active portion of them. It has also been said that some polysaccharides and some lipoids may have the same effect. Some drugs which produce allergic responses do not contain protein. The exact mechanism of the they act by producing antibodies in the allergic subject and that the combination of allergens with fixed antibodies in the tissues produces the allergic response possibly by the release of histamine or some allied substance. The nature of the response depends on the site of the allergen antibody reaction in the skin, urticaria or oedema in the nose, rhinorrhoea or both in the bronchi, broncho-spasm and oedema and hypersecretion by the mucous membranes.

The exact mechanism by which the reactions are brought about is uncertain but the probability is that in the case of bronchial asthma the afferent paths may be from the nose or from the bronchi themselves or possibly from the stomach or other organs and that the action is by vagal stimulation with consequent constriction of the smooth muscle of the bronchi and hypersecretion.

There is plenty of evidence for considering a great number of cases of asthma as being in the allergic category. For example in a considerable percentage of cases there is a family history not only of asthma but of many allied disorders considered to be allergic in origin such as hay fever, urticaria, food sensitivity and less certainly migraine. Furthermore asthmatics are themselves notoriously sensitive to injections of serum or even in the administration of drugs. When proteins are injected into the skin of asthmatic patients they cause local reactions to certain of the substances injected which are quite different from the reactions to others or to the reaction of controls. The skin reaction and sometimes the attacks of asthma are associated with a local or general eosinophilia which is one of the concomitant factors in allergic manifestations as a whole. Clinically cases of asthma are seen in which the patient can be shown to be sensitive to simple substances such as grass pollens which can cause both hay fever and asthma in the same subject, others only get their attacks when exposed to the inhalation of certain dusts at their work and never at other times. Asthma in many cases can be proved to be provoked by exposure to horses, dogs or cats, to feathers, to kapok, to face powder, to house dust and to a whole host of simple and complex substances. To the allergist the whole thing is simple: allergen + antigen = asthma. Without allergens without antigens there can be no asthma.

There is no question about the importance of allergy in asthma, but perhaps too much emphasis has been placed on it through concentration on the anti-allergic methods of treatment which have pre-occupied so many professionally successful allergists, so that what lies behind the allergy has often been ignored

## OTHER AETIOLOGICAL FACTORS

### Psychological factors

There is no doubt whatever of the psychological aspect of asthma, even if one cannot go as far as some psychiatrists, who regard asthma as a psychoneurosis and the allergic manifestations as secondary to the psychoneurosis. Some psychiatrists have claimed to be able to cure their patients by psychoanalysis alone without bothering about the allergens at all. Allergists agree that psychological treatment is important, but only as a secondary factor in a condition in which the allergens play the leading and essential role.

### Other mechanistic theories

There are other mechanistic theories, such as the one which suggests that there are trigger zones in the nasopharynx, which can produce attacks of asthma when stimulated, and that, as a natural consequence of this, the patients can be made insensitive by operation, by cauterization or by local injections, but these have fallen into disrepute. Other authorities state there are biochemical or endogenous toxic causes for asthma, this theory leads to elaborate dietetic therapeutic programmes. There is also an infective theory of the causation of asthma. These and other theories are referred to below.

### Extrinsic and intrinsic causes

Rackeman (1931) neatly classifies the types of asthma into (1) extrinsic asthma, in which he includes all the asthma produced by extrinsic allergens inhaled, ingested or injected, and (2) intrinsic asthma, into which are bundled all the other sorts—infective, toxic, neurotic, reflex and all the rest. Many cases, however, refuse to stay neatly even in these comprehensive pigeon-holes, and many cases that might reasonably be classified as extrinsic are affected or modified by causes that should be labelled intrinsic. However, as the classification has some slight bearing on treatment, it may serve a useful purpose.

### Incidence

Study of groups in Great Britain and in the United States of America and elsewhere suggests that an incidence of about 0.59 per cent is probably correct for both this country and the United States.

**Race**—Little is known about racial susceptibility except that it is probable that the disease is much more common in the "civilized" races than in the others, and that Jews have a relatively higher incidence as a racial group.

*Sex*—There is approximately the same distribution of asthma between the sexes, although some writers state that it is slightly more common in males

*Age*—Asthma makes itself apparent in the first decade of life in approximately 30 per cent of cases; most of the other cases are fairly evenly spread over the next three decades. Excluding cardiac and renal asthma, the disorder is much more rare after the age of 50 years, but it does occur at this age and even later. Those who classify asthma as extrinsic or intrinsic would say that extrinsic asthma is much commoner in younger people, while intrinsic asthma is much commoner in the older group

*Social status*—Asthma is rather more common in the relatively well-to-do, and amongst the workers it is more common in sedentary than in manual workers, even making allowances for the special exposure of these latter in particularly dusty trades, such as flour-millers, cloth workers, rag sorters, furriers, hairdressers, poultry-dealers and the like

*Intelligence*—There is a fairly general idea that a high proportion of asthmatics are to be found amongst the more intelligent section of the community. There may be some truth in this, but the sufferers from asthma are often more aggressive, more self assertive and emotionally less stable than their non asthmatic associates. They sometimes become extremely possessive, selfish and introverted, but most of the characteristics described may occur as the result of their being set apart by their condition and be otherwise unassociated.

*Environment*—Geography plays little part in the aetiology of asthma. No locality is completely immune. Patients often attach great importance to environment, stating that they always get asthma when they go to X and never when they go to Y. Occasionally there may be some particular exposure to allergens at X, which is absent at Y, but more often the attacks at X and the freedom at Y are the results of auto-suggestion. At all events, in a large clinic, some patients will tell you that if they could leave X and be at Y all would be well, but others tell you that if only they could leave Y and go to X they are sure they would be "all right". Taking it by and large, patients are often freer from attacks at high altitudes, provided that they have not developed complications. This may be due to the absence of dust, to the freedom from infection or the general well being induced by altitude, or to just plain suggestion.

*Heredity*—The first book on asthma published in England, that by Sir John Floyer written in 1698, mentions the hereditary tendency in asthma. Since his time there have been many studies, most of which tend to show that the family histories of asthmatics show others suffering from asthma or allergic diseases in approximately 50 per cent of cases, whereas controls give a family history in only about 7 per cent. It would appear that more members of the family are affected when both the parents have an allergic history than when only one has this, and it would appear in the majority of cases that what is transmissible is the asthmatic diathesis and not sensitivity to any particular substance, although there are family histories which appear to show a transmissible hereditary sensitivity to eggs, strawberries or other foods.

In general, asthmatic subjects tend to occur in small families, and genetic studies do not reveal a convincing mendelian pattern.

## CLINICAL FEATURES

### Description of asthmatic attack

*Precipitating causes*—Asthma may occur at any time of the day, but in mild cases it commonly starts in the early morning and wakes the patient up, or else it may become more obvious when the patient wakes at his normal time and is getting out of bed. The stimulus which causes the attacks in the latter case is probably the first breath of cold air when the patient wakes up, associated with the movement and deeper breathing on waking which disturbs the secretion that has been collecting during the night. Any patient with acute or chronic bronchitis starts to cough as soon as he gets up in the morning, and the colder the air he inhales the more prompt is the reaction. In a case of ordinary bronchitis the result is coughing, in the asthmatic the result is bronchial spasm. The probable explanation of the nocturnal attacks is more complicated. Metabolic changes during sleep have been postulated but remain undiscovered. Feather sensitivity, due to inhalation of dust from the pillows, has also been suggested, therefore many an asthmatic has been advised to sleep on kapok or, if he is sensitive to kapok, to sleep without a pillow or on a pillow covered with oiled silk or Cellophane, but the real explanation is probably a mechanical one—the result of respiratory embarrassment in the horizontal position, especially with the slight encroachment of the soft parts and interference with the air-way in deep sleep.

Attacks are often produced by exertion, probably associated with increased ventilation due to exertion, and they may be precipitated by emotional factors. They can, of course, be produced by exposure to specific inhaled allergens, such as, for example, certain pollens and industrial and domestic dusts. Exposure to cold air is mentioned above, exposure to fogs, smokes or chemical fumes may have the same effect in asthmatics, and in such a case it is merely an exaggeration of the normal reaction in non-asthmatics.

### Nature of the attack

The typical attack varies from a slight feeling of constriction behind the sternum or trachea or around the ribs, produced only on attempting deep respiration or on exertion, to the most intense discomfort with violent, noisy inspiratory effort and expiratory distress. During the severe attacks the patient is reluctant to move, he sits bolt upright in bed with his head thrust down and forwards, his mouth open and his alae nasi working, his chest is elevated, his shoulders are raised, with all his accessory muscles being used to move his chest, the veins in his neck are bulging, his sternomastoid muscles are taut, and his diaphragm is heaving. Often he will lean forwards over a table or chair in order to breathe more freely.

All this effort produces much respiratory movement, but it is rather an up-and-down movement than an expansive one. If the attack is a prolonged one or the patient has become emphysematous as a result of repeated attacks, he will become cyanosed. He can talk, if at all, only in gasps, is unable to eat or drink, and may, in some cases, become most upset and emotional. His respiratory embarrassment is audible first as an expiratory wheeze and then by wheezing inspiratory and expiratory sounds, interrupted from time to time by a short, usually ineffective, cough.

*Objective findings*

On examination it will be found that the expansion of the chest is grossly diminished even when the movement is maximal. The rhonchi can be felt, the chest is everywhere hyper-resonant, the cardiac and liver dullness is decreased, and the diaphragm is depressed. Auscultation shows that the breath sounds are diminished but are often obscured by sibilant and coarse rhonchi. In the later stages there are râles of all sorts of pitch, more or less uniformly distributed through the chest.

That the physical signs of hyper-resonance are due to over-distension of the lungs with air, and the rhonchi mainly to bronchial spasm, can be shown by giving the patient an adequate dose of adrenaline. After this medication, hyper-resonance disappears, the normal pulmonary boundaries are restored and the rhonchi disappear, in an early case. In the more chronic cases the originally transient distension of the asthmatic attack results in permanent distension due to emphysema.

*Periodicity of attacks*

Some cases show a periodicity which is of some interest. Certain patients who are sensitive to pollen get asthma only during the periods when the pollen is airborne and not at other times of the year. A large group who suffer from bronchitis, either as the result of, or as the cause of, their asthma, only get attacks during the winter months and not during the summer. Many patients with asthma gain complete relief or improve with puberty, but some cases appear for the first time at that period. Some patients state that their attacks coincide with their menstrual periods, but some women tend to place all unusual phenomena in some relation to their periods, and there is little evidence that there is any real connexion; the menopause does not seem to affect the course of the condition. Many more women are relieved of their attacks during pregnancy. It is not clear whether the factors here are metabolic, mechanical or psychological. Sometimes patients have attacks only at week-ends, or during school holidays or similar periods. Here again the explanations may be multiple: environment, change of habits or routine, topical exposure to allergens, or psychological causes may each or all be operative.

*Conditions allied to asthma*

*Hay fever*—Certain patients are sensitive to grass pollen or, more often, to grass and tree pollens. These cause seasonal rhinorrhoea, sometimes associated with attacks of asthma. The tendency is for the attacks to become less severe as the patient grows older, and they tend eventually to disappear spontaneously. Records of this sensitivity frequently occur in the family history of cases of asthma.

*Allergic rhinitis*—The symptoms of this condition are similar to those of hay fever, namely sneezing, running from the nose, and nasal obstruction, but the disorder is not seasonal in duration and does not appear to be due to pollen. It tends to occur in many people on first rising in the morning and, although it may remain bothersome for years, it tends to die out in the long run. Many explanations have been given for it, varying from sensitivity to dust, ororris face-powder to local nasal conditions such as polypi or sinusitis. It is certainly



## ASTHMA

commoner in women, particularly young ones. Sufferers from this condition not infrequently tend to suffer from asthma sooner or later, and it also is an important finding in the history of asthmatics.

*Food idiosyncrasies*—Some patients get acute abdominal pain after the ingestion of certain foods. All sorts of foods have been implicated, but chocolate, strawberries and eggs appear rather more frequently in case reports of this condition than do other foods.

*Urticaria and angioneurotic oedema*—These conditions are often allergic and are sometimes definitely associated with the ingestion of certain drugs or foods, or with the administration of serum, but they may appear apparently spontaneously or, at all events without manifest cause. These conditions tend to occur rather frequently in asthmatics but it does not seem that the reverse is as common.

*Migraine*—This is often said to be allergic in origin and therefore might be of some importance in history taking, but its exact relation to asthma is rather uncertain.

*Eczema and prurigo (Besnier's)*—Eczema is often found in infants and young children, either associated with asthma or preceding it. It is thought to be allergic in origin. It rarely persists into adult life and it even more rarely appears in adult sufferers.

*Seasonal conjunctivitis*—Conjunctivitis is often associated with hay fever but cases occasionally occur independently of the nasal symptoms.

*Other conditions*—Many other conditions, from Meniere's disease to epilepsy, have been described as allergic and therefore important in an allergic history, but their position in this regard is, unlike the conditions mentioned above, more than dubious.

*Status asthmaticus*—See under Treatment.

### Complications of asthma

*Emphysema*—Probably this is one of the commonest and most important complications. If spasmodic distension of the lung occurs sufficiently often and for sufficiently prolonged periods, over distension and degeneration of the alveoli follow and the emphysema becomes permanent. Most chronic asthmatics show evidence of this condition, and the well known barrel chest, with the hunched shoulders and the forward stoop is the typical appearance of the confirmed asthmatic. When asthma occurs early in life, before the skeleton is fixed, there is often sucking in of the lower intercostal spaces and a pigeon chest develops.

*Loss of vital capacity*—The vital capacity in asthmatics is considerably diminished, at first during the attacks with recovery afterwards, later the diminution becomes permanent, although it is surprising what an improvement in vital capacity may be induced by re-education in breathing even in severe cases.

*Bronchitis*—Bronchitis can be a complication of asthma, or else asthma can be a complication of bronchitis. In most long standing asthmatic cases bron-

## CLINICAL FEATURES

to be secondary to bronchitis and often these cases start with what appears to be a common cold but the infection of the bronchi becomes chronic. In many cases the upper respiratory catarrh is almost certainly a common cold although the possibility of allergic rhinitis must be borne in mind.

*Spontaneous pneumothorax*—This condition is found occasionally when asthmatics are often and routinely submitted to radiography. The probability is that the condition is due to the rupture of emphysematous bullae under the strain of the asthmatic attacks. The symptoms are apt to be submerged in those of the asthmatic attack and the physical signs may be so submerged also so that the fact that the condition is found not infrequently in a casual manner suggests that it occurs more frequently than it is diagnosed. Occasionally it is associated with surgical emphysema.

*Bronchiectasis*—This condition is more often associated with asthma than appears in most text books and there is a generalized bronchial dilatation which gives a rather characteristic radiological picture with marked hilar enlargement and radiating opacities. These cases are often erroneously diagnosed as pulmonary tuberculosis because of a superficial resemblance in the radiological picture. Not infrequently evanescent tension cavities are associated with this condition.

*Pulmonary tuberculosis*—Asthmatics are not immune from pulmonary tuberculosis but when the disease occurs in an asthmatic the diagnosis is apt to be made rather late because all the symptoms and physical signs tend to be attributed to the long standing asthma, emphysema and bronchitis.

*Cardiac complications*—As a late complication chronic anoxic pulmonary heart disease (cor pulmonale) becomes an important and probably an invariable accompaniment of the chronic case in old age. The condition is more likely to develop in cases complicated by the diffuse bronchiectasis described above.

*Massive collapse*—Massive collapse of a lobe or lobes occurs occasionally in asthma when the partially constricted bronchioles are occluded by plugs of impervious mucus. This is most likely after the use of morphine and is the probable cause of death in some cases.

## PROGNOSIS

Considering the apparent severity of the symptoms and the discomfort caused by them it is surprising how little asthma seems to curtail the expectation of life of the sufferers. A great many patients live on to a ripe old age even when the symptoms are continuous. It is sometimes stated that no one ever dies of asthma but this would be an overstatement even if one excluded deaths from the administration of morphine or similar drugs and the occasional accidents of adrenaline since the emphysema which complicates the long standing condition may cause serious cardiac disability and there is always the possibility of broncho pneumonia as a complication. Nevertheless even allowing for these possibilities it is surprising how rarely lethal is a condition at once so spectacular and so crippling.

## DIAGNOSIS

**Clinical diagnosis**

This is made almost entirely on the clinical history—paroxysmal shortness of breath with tightness in the chest and wheezing, but with characteristic freedom from symptoms during the intervals between attacks. The occurrence of the attacks in the early hours of the morning or on waking or at certain seasons of the year, and the relief of symptoms by antispasmodics such as adrenaline or ephedrine, the family history, and the personal history of associated conditions such as hay fever or eczema, all combine to present a picture which helps to differentiate asthma from other conditions somewhat similar in some respects. The most difficult condition to differentiate, however, is cardiac asthma.

**Differential diagnosis**

*From cardiac asthma*—The age of first onset is of some importance in differentiation. Cardiac asthma would be unusual before the age of 45 years, except in the presence of valvular disease of the heart, which would be manifest. It is unfortunately not safe to assume that nocturnal dyspnoea, coming on for the first time over the age of 45 years, is unlikely to be bronchial asthma, because, although asthma usually starts in childhood or adolescence, not a few cases, especially those associated with bronchitis, occur for the first time in the fifth decade or later. The patient with cardiac asthma may have other signs of cardiac decompensation but this is not essential. He will usually suffer from more distress and anxiety than does the patient with asthma from other causes. His lungs will often show marked basal oedema and his sputum may be blood stained. A case showing well-marked cardiac disease is easy to distinguish but there are borderline cases which are much more difficult.

*From constriction or obstruction*—A rare cause of pulmonary wheezing is constriction of the bronchi, produced by enlarged glands or pressure of an intra-thoracic goitre, or by an enlarged thymus gland or a bronchial or mediastinal neoplasm. The wheezing is then often localized or unilateral and presents little difficulty in differential diagnosis.

**Diagnostic aids***Skin tests*

These are performed either by scratching the skin through a drop of a solution containing an extract of various protein substances, or else by the intracutaneous injection of similar solutions. By this means it is possible to test the sensitivity of the patient to a wide range of substances, first by testing for grouped reactions for a range of substances—for example, feathers or moulds—and then, if sensitivity is shown, by making differential separate tests on the various component parts of the group. Some clinicians attach great value to these tests, and some text-books suggest that the whole diagnosis of the cause of a patient's asthma and his consequent treatment depend on the identification of the offending allergens by means of skin tests, and on the desensitization of the patient by the injection of small but increasing doses of the allergen in question. It is a simple conception but, alas, far from the truth, for the skin tests do not tell a simple story.

## TREATMENT

Reactions to skin tests are certainly more likely to be positive in patients who suffer from asthma or allied conditions, but they are often positive in a great many people who do not have these conditions or any family history of them. The reactions are not always quite consistent in the same patient. There is no certain correlation between the protein to which the patient's skin is sensitive and the cause of his asthmatic attack. Thus skin tests form an interesting study and are occasionally useful but we are far from understanding their full significance and any considerable reliance on them for the guidance of treatment is unjustifiable.

### *Eosinophilia*

Abnormal numbers of eosinophil cells may be found in the sputum or in the blood during attacks of asthma and may therefore be of some diagnostic significance, but in the majority of cases they are absent and their appearance seems to be more common in the first attacks than in later ones.

## TREATMENT

Treatment can be divided into (1) preventive treatment (2) symptomatic treatment and (3) specific treatment. Whole books have been devoted to the third division. For reasons which I mention below not much time is devoted to it in this article.

### *Preventive treatment*

#### *General protective measures*

Preventive treatment is any treatment which is designed to make the occurrence of attacks less likely. Patients for example may be advised not to take heavy meals last thing at night since it has been found that a full stomach at bed time makes nocturnal attacks more likely or they may be advised to avoid cold baths in the early morning or to avoid the stimulating effect of frosty air soon after they wake up.

Patients may be given sedatives regularly such as phenobarbitone either during the day time or if their attacks are nocturnal last thing at night in order to make the attacks less likely. They may be given ephedrine regularly for the same reason whether they are having the attacks or not. They may be taught breathing exercises, such as those prescribed by the Asthma Research Council. These improve and maintain the mobility of the chest and the diaphragm diminish the dead space and probably have a psychological effect on the patient as well.

#### *Avoidance of allergens*

Some people consider that the most important part of preventive treatment is the avoidance of allergens when possible. Patients who know that they are sensitive to dogs cats horses hay pollen and the like should take steps to avoid them. Allergists interpret the skin tests as indicating the substances to be eliminated from the patient's diet or from his environment. They sometimes go to great lengths in this direction making the patient wear special clothing and sleep in rooms denuded of all hangings or carpets, and telling him to use pillows covered by Cellophane or with their filling confined to flock or kapok instead of down feathers. Completely dust proof rooms have been constructed for these patients, but extravagant measures based on the uncertain premises of the skin reactions are

not justifiable. It is always worth while to try the effects of eliminating substances to which the patient appears especially sensitive, and some success attends this attempt, especially in early cases in which the sensitizations are simple and few and multiple sensitization has not taken place. As a general rule it is wise for a patient to avoid dusty atmospheres and to keep out of the way when dusty domestic operations are in progress, and the rooms in which he lives should be as clean and dust-free as good housekeeping should ensure.

#### *Occupation and environment*

Advice which involves a patient in changing his occupation should not be given lightly, since an unsettling change may provide more symptoms than it is designed to relieve. Sometimes it is better to keep a patient in an occupation which is theoretically unsuitable but which he enjoys and is good at, than encourage him to transfer to another in which he may be miserable and unsuccessful.

On the same principle, changes of house or environment should not be lightly heartedly recommended. They are rarely successful, and in any large clinic, patients from town X will say that if only they could be in town Y all would be well, and at the same time patients from town Y say that if only they could live in town X all would be well. With the rare exception of a simple sensitization to an allergic contact in one house and not in another, these odysseys are doomed to disappointment. The indications for advising change of employment or environment are usually quite obvious and definite when the sensitization is single, proved and definite, which, in fact, is rarely the case.

#### *Symptomatic treatment*

This is unfortunately, at the present time, in default of any real curative treatment, the most important section of treatment.

#### *Relief of bronchiolar spasm*

The main object of therapy is the relief of spasm of the bronchioles.

**Adrenaline**—This is best achieved in the majority of cases by the use of adrenaline. The drug can be administered in a spray, by the use of nor adrenaline tablets dissolved in the mouth or, most effectively, by the subcutaneous injection of 1 1,000 adrenaline. Responsible adult patients should be trained at an early stage to use a hypodermic syringe and a solution of adrenaline. The correct individual dose has to be arrived at by trial and error, but an approximate can be followed by the injection of approximately 0.5 cc. The correct symptoms are relieved. This is effective in the majority of cases. A large dose of adrenaline is effective in the majority of cases with large doses of adrenaline. The earlier the injection is given during the attack, the less is the quantity necessary and the more complete the relief. Adrenaline should not be given intravenously since it may cause sudden collapse, deaths have followed accidental injection into a subcutaneous vein. It is impossible to over-estimate the value of adrenaline in asthma and fortunately few adrenaline-resistant cases are seen. Patients will sometimes say that adrenaline is of no help to them. This is often found on re trial to be incorrect in fact, and to be due to incorrect administration in the past and the fixed erroneous conceptions

## TREATMENT

to which asthmatic patients seem to be peculiarly liable. Intramuscular injections of 1-1000 adrenaline in oil can also be given. The object of giving it in this form is to produce a more lasting effect but it is rather disappointing in this context.

**Cardophyllin** (syn. aminophylline) — In the rare cases in which adrenaline does not produce relief, Cardophyllin (theophyllin ethylenediamine) given intravenously, is often effective in relieving bronchial spasm. It can be given in doses of 0.25 or 0.50 grammes of Cardophyllin in 20 millilitres of distilled water. Intramuscular injections are ineffective and painful. Cardophyllin can be given effectively *per rectum* in doses of 0.3-0.6 grammes in 15 millilitres of water. When it is given by this route there is a latent period of about an hour but the effect is more lasting than when it is given intravenously. In status asthmaticus Cardophyllin may be given continuously in saline solution in intravenous doses of 0.5-1.0 grammes in 1000 millilitres of glucose saline solution.

**Ephedrine** — Ephedrine sulphate is worth a trial in most cases of asthma. It can be taken either regularly or in advance of the time when attacks appear to occur. It can effectively be taken for example just before the patient goes to bed in order to prevent nocturnal attacks. It is wise in this case to combat the exciting and sleep-preventing effect of the ephedrine by giving  $\frac{1}{2}$  grain of phenobarbitone at the same time. Ephedrine although easy to take is unfortunately ineffective in many cases and a tolerance for the drug is often rapidly built up so that even if it is effective at first it becomes ineffective later. Intervals without ephedrine allow the recovery of its effectiveness in individual cases.

**Histamine antagonists** — Phenergan, Benadryl and the like have proved disappointing in the treatment of asthmatic attacks and no antihistamine drug has yet been found fully effective in the control of asthma.

**Inhalants** — Adrenaline can be given effectively by inhalation either from a simple vaporizing apparatus with a rubber bulb or else in more elaborate vaporizers using the passage of oxygen through the vaporizing fluid. The fluid usually used is adrenaline 1-100 solution. Isupren or Aleudrin an iso-propyl derivative of synthetic adrenaline may be used instead. It has fewer side-effects than has adrenaline. There are many proprietary inhalant fluids on the market which are effective in so far as they contain adrenaline the other constituents having little effect apart from this.

**Hypnotics** — Hypnotics are often useful for asthmatic patients and the one most usually employed is phenobarbitone in doses of  $\frac{1}{2}$  grain. Morphine should never be given to asthmatics for the treatment of their asthma particularly in status asthmaticus.

### Resting in breathing

Breathing exercises are a most effective form of treatment. They can be given to increase the vital capacity and diminish the dead space. In addition they may be successful as a form of psychotherapy. Breathing exercises often aggravate the shortness of breath and produce cough at first but they should be persisted in because ultimately they will give relief. It is advantageous to give inhalations of adrenaline before starting the exercises. The exercises should be taught in the first place by a physiotherapist skilled in their use. The exercises recommended are those published originally by the Asthma Research Council but which are

now published under the title *Advice to Asthmatics* by the Asthma Clinic (1949)  
at Guy's Hospital (London)

*Treatment of status asthmaticus*

Status asthmaticus—a condition in which the attacks become continuous, with little or no remission by day or by night—is the most alarming and most troublesome complication of asthma which the physician is called upon to treat. The consultant physician requires all the confidence born of his knowledge of the favourable prognosis in order to resist the impression produced by the suffering and anxiety of the patient and the terror and desperation of his relatives, as well as of the exhaustion and despondency of the patient's usual medical attendant. In the home, the long continuance of the attacks, in the face of every remedy, has often built up a psychological situation in which any rational approach to the handling of the case has become impossible. The patient is then best admitted to a hospital, preferably to a general ward. Matter of fact reassurance of the patient, in a new atmosphere of apparent competence, induces an attitude of mind in him which is more favourable to suitable therapy. The presence of other patients besides himself who require attention removes him abruptly from the centre of the stage, which he has painfully occupied at home. The effect is often salutary and he begins to improve at once.

If the patient is cyanosed, he may with advantage be placed in an oxygen tent and given oxygen with 5 or 7 per cent carbon dioxide. The tent has the double advantage of supplying the needed oxygen and of screening the patient from others in the ward, who might be unduly sympathetic under the influence of the rather terrifying sight of his respiratory embarrassment, which, however genuine, is unconsciously exaggerated in the presence of an audience.

Adrenaline should be given by the minim-a-minute method and repeated at regular intervals if there is any response. These cases sometimes seem to be resistant to adrenaline, in that case, they are often relieved by intravenous injections of Cardophyllin, 0.25 gramme in 10 millilitres of distilled water, this treatment can be repeated 4 hourly until relief is obtained. Rectal administration of Cardophyllin (0.5 gramme in 20 millilitres of water) can be substituted for or combined with intravenous injections. The effect of intravenous injections should be immediate, the effect of Cardophyllin rectally takes about an hour to develop but the effect is more prolonged. If there is great dehydration, Cardophyllin may be given intravenously in a continuous drip, but patients sometimes associate an intravenous drip with imminent danger of death and the psychological effect may be bad. Patients should be encouraged to drink fluids in the interim, and can usually be persuaded to do so if their fears about their imminent dissolution have been effectively combated. Mild sedatives may be given, such as phenobarbitone, Sodium Amytal or Nembutal in doses of  $\frac{1}{2}$ –1½ grains, but it is not advisable to "push" these or any other sedative drugs.

In cases which fail to respond to other methods of treatment, ACTH or cortisone may be tried. Good results are claimed by Zeller, Randolph and Rollins (1950), Howard and his colleagues (1950) and others. The recommended dosage of ACTH is from 50 to 100 milligrams per day divided into 6-hourly doses given for from 3 to 4 days. Some workers have found cortisone less effective than

## TREATMENT

ACTH but in this case the usual dosage is 200 milligrams per day divided into 6-hourly doses. These substances are given by injection but cortisone has also been given by mouth (Schwartz 1951). This method of treatment is well worth an extended trial. Herxheimer (1952) has reviewed the results claimed by various American authors in 133 cases treated and notes that 53 per cent became symptom free, 19 per cent were partly improved and 28 per cent were unimproved. No similar study of a relatively large group of patients has been reported in this country but the results in individual cases although occasionally encouraging have not been spectacular when objectively studied.

Morphine should never be given to these patients. It is very tempting to give it in order to produce mental relief and rest. Many doctors give it for this purpose and the patient survives in a great number of cases, but sooner or later a doctor who is in the habit of giving it will give it to a patient whom it will kill as certainly as if he were strangled which indeed he is by his own internal secretions. Anyone who has ever seen a patient gasping like a landed fish will realize on what a narrow thread life depends in such a case. Morphine depresses the respiration so that the patient suffocates.

Breathing exercises and postural drainage should be encouraged at an early stage in order to remove the secretions in the bronchial tree. Expectorants are some times given with advantage.

Bronchoscopic aspiration would probably be helpful but bronchoscopy of a possibly terrified patient is extremely difficult.

The prognosis as regards recovery from these attacks, however distressing they are, is good provided that the patient is not elderly and that pulmonary infection does not supervene. This possibility should be guarded against by the use of antibiotics.

### Specific treatment

Most text books and all allergists state that asthma is a readily curable disease, thus Unger stated in 1944. In comparison with other chronic diseases such as chronic arthritis and cardiovascular renal disease the results of treatment in asthma are usually brilliant. In those who do not obtain complete relief there is usually amelioration of symptoms, i.e. the attacks of asthma are milder and less frequent. Failures occur of course but in only a small minority and are often due to negligence on the part of the patient (or parent) or of the Physician. Urbach and Gottlieb in 1949 wrote: 'The large group of allergic asthma including that due to pollen is today readily amenable to treatment. Rackemann (as quoted by Cecil 1947) more conservatively states that the hope of permanent relief depends entirely upon the discovery and the removal of the cause. With that statement I am generally in agreement since he was referring only to extrinsic asthma but how much more easily is this said than done.'

Rackemann does not suggest that it is easy or that it can always be done but hundreds of enthusiastic and successful physicians say that it can. They suggest that the cause in the majority of cases can be discovered by a correct interpretation of the history and of the skin tests. They therefore require that every effort should



be made to avoid or eliminate the cause or allergen and, if that is impossible or unsuccessful, that the patient should be desensitized against the allergen. That is, he should be given repeated small, but increasing, doses of the antigen until an excess of antibodies is produced in the blood, so that further doses of antigen are immediately neutralized before they can have an effect on the tissues. Since this conception would mean the continuous administration of antigen to keep up the antibodies to the desired level, it is further postulated that continual saturation with antigens, or the production of mild anaphylactic shock, eventually leads to permanent "de allergization" through the prevention of the formation of antibodies. Whatever the theory may be, the commonest treatment in vogue for extrinsic asthma is desensitization, specific or non-specific.

#### *Specific desensitization*

Specific desensitization is carried out by the administration of preparations of the allergen or allergens considered to be the cause of the asthma in the particular case. Allergists depend on the history and skin tests for the indications of the appropriate solution of allergens to be used, and the preparation is individually made up from these, but solutions are often prepared from a solution extracted from ordinary house dust if other specific indications are lacking. Some allergists take a tremendous amount of trouble in trying to find an individual solution which is more likely to be effective, others depend on more or less stock solutions with minor additions. Whatever the method used, good results are claimed and, what is more, are claimed by patients.

I am dealing in cavalier fashion with a method of treatment about which many books could be and have been written, but before discussing it further I shall quote one piece of work which shows how difficult it is to be certain of cause and effect in such an uncertain condition as asthma. The results of treatment by desensitization may be judged in the light of it.

A few years before World War II, Witts and Conybeare, at Guy's Hospital in London, started some control experiments on the treatment of asthma, and the work was carried on by myself and Lintott up to the outbreak of war (Boland and Lintott, 1939).

A careful selection of cases was made, and the eventual series of about 50 cases in each group were the product of a careful screening of hundreds of patients. No patient was selected for the series whose attacks were infrequent or were of recent origin, who had a history of long periods of remission for any reason, or who was unreliable in any way. A careful and detailed history was then taken, giving a close account of the attack pattern, its frequency, duration, severity and causation together with the history of treatment, symptomatic and otherwise. Such patients as gave a good history and were considered consistent and reliable were then subjected to routine investigations.

They were advised to eliminate or avoid any allergens which seemed (on the history or, less certainly, on the skin tests) to be possible causes of their attacks. They were given advice about the uses of the various symptomatic drugs, as mentioned above. They were given general advice about eating at bedtime, avoiding dusts and so forth, and then were sent for 2 months' (or more) instruction in

breathing exercises. At the end of this 3 months period a further assessment was made and it was often found that general treatment and advice of this sort had apparently had a good effect. Therefore many cases had to be removed at this stage from the test groups originally selected. However the few survivors were a well tried band whose symptoms still remained and whose pattern was still consistent.

These highly selected persons were then divided into two equal groups. One group was sent to a well known allergist for treatment by protein desensitization and the other group was treated by us with injections of normal saline solution. After 3, 6 and 9 months the cases were re-assessed. Of the number of patients sent for desensitization nearly two thirds had improved either objectively or subjectively. These results considering the severity of the cases chosen for a study were striking but the results in the equivalent group treated with normal saline were equally good or even in our series slightly better.

A further study of the groups showed what to me was the most important finding of the experiment. I mentioned above that many patients were objectively or subjectively improved. Those regarded as objectively improved were the ones whose improvement could be recognized by us from a study of our case records. Those said to be subjectively improved—and these were the majority—were the patients who claimed to be improved although the improvement could not be substantiated from the case records. For example a patient might claim to be much better since starting the treatment stating that he now only had attacks once a week. Reference to the notes taken before the treatment commenced might give the frequency of attacks as being once a week. When this was pointed out to the patient he would express surprise and say that anyway they only lasted about a couple of hours. Reference again to the notes would show that according to the notes the average duration of an attack was about 2 hours. The patient would be momentarily nonplussed when confronted with this history and would then say that at all events he was miles better than he had been. Beyond this it was impossible to go.

Now it is not perhaps really the business of a physician to argue with a patient who thinks he is better and if asthma is regarded as merely a symptom and the patient thinks he is better it could be maintained that he is better. The fact is however that in these experiments normal saline solution proved as effective as were protein solutions intended to desensitize. This gives food for thought but it may be reasonably stated that it is no valid argument against desensitization since the selection of cases and the solutions used may not have been the correct ones. Even if we agree with this point of view though with some mental reservations the most important result of the investigations was the demonstration that the subjective improvement experienced by the patients was not susceptible of objective proof. If effective desensitization were in fact frequently achieved by these methods of treatment improvement would not be a matter of opinion and a subject for dispute.

Similar experiments were carried out in the controlled treatment of rhinorrhoea and the same results were achieved. Just as many cases were relieved by normal saline solution as by protein solutions. Here the question of subjective

improvement did not arise for the only criterion was the stoppage or otherwise of rhinorrhoea. Similar experiments were carried out with autogenous and stock vaccines, again with similar results. The outbreak of the war prevented the extension of these experiments to hay-fever desensitization. Rendered somewhat critical by our findings, we were anxious to see whether this treatment when critically reviewed, was any more effective.

I have discussed these experiments at some length in order to explain why I have not devoted more space to what is a generally accepted treatment for extrinsic asthma, and I shall conclude by giving my views on treatment by desensitization with protein solutions and, incidentally, on the treatment of asthma with vaccines, autogenous or otherwise.

Most treatment of this type is effective only by suggestion and the improvement produced is often more apparent than real. If the effectiveness of the treatment is judged by "cures"—which may be a harsh criterion, but not an unreasonable one—the claims made by allergists are for 20 per cent "cure". The fact that the same figure can be produced by injections of normal saline solution suggests that the "cures" are the result of chance, of the uncertain natural course of asthma and of the powers of suggestion. The theory of desensitization is a sound one but in the vast majority of cases not a practical one, possibly because the basis of asthma is the tendency to become sensitized, and because the majority of patients have become or are capable of being sensitized to so many different unavoidable substances that desensitization is difficult or impossible. I still think that, if a patient can be found to be sensitive to one or two proteins only, it is possible that desensitization may be effective, but this slight residual speculative belief does not seem a firm substance for the treatment of allergic conditions, with which Urbach and Gottlieb, for example, deal in a book of nearly 1,000 pages.

My attitude and theirs may perhaps be considered in the light of a paragraph in their book

"In conclusion, one more important point must be considered. The great majority of physicians are of the opinion that asthma is a dangerous but practically an incurable disease. Although they naturally refrain from expressing this opinion to the patient, the latter very soon senses this pessimistic attitude which very often does immeasurable harm psychologically and consequently physically. The physician who is convinced that many cases can be cured and that almost all can be considerably improved and who is able to communicate this conviction to his patient, possesses the most important pre-requisite for success."

## CONCLUSIONS

My scepticism with regard to the effects of present methods of so-called "specific" therapy may have conveyed the sense of pessimism which Gottlieb and Urbach rightly deplored. I am not in the least pessimistic about the treatment of asthma from a symptomatic point of view. A great deal can be done for most patients suffering from asthma. Alteration of the patient's psychological reaction to his condition—and, as necessary, the alteration of the relatives' reaction to it—is of the utmost importance. This can be done by explanation and reassurance.

## CONCLUSIONS

The use of breathing exercises and the intelligent use of anti-spasmodics have, in most cases, an excellent effect, and the patient gains in confidence when he knows that he has the means of relief at his disposal. If it is possible to detect the precipitating cause of the attacks of asthma, and if this can be avoided, then the prognosis is so much the better. I would not even exclude the possibility of desensitization in an individual case, if a single protein allergen appeared to be specially implicated. I think that its effect is not proved, but that it might be worth a trial, although, in the vast majority of cases, I believe that present methods of desensitization have only a psychological value.

If the attacks of asthma cannot altogether be eliminated, the patient can be taught to live with his asthma, just as a patient can be taught to live who has lost his arm or his leg. He should at all ages be encouraged to lead a life which is normal so far as humanly possible, he should not be encouraged to regard himself as an invalid and, especially, not to regard his asthma as a justification for dodging responsibility or for getting his own way. The power of the spirit is never so beautifully displayed as in asthma. One man with quite mild asthma will mentally surrender and become a complete invalid, another with infinitely more severe asthma will struggle on and lead a useful and effective life in spite of his severe handicap. Sir Arthur Hurst, a life long sufferer from asthma, who had undergone almost every treatment known to man without much effect, almost never allowed his symptoms to interfere with his physical or intellectual life. He would do his ordinary ward rounds on a day and in a condition when other men would have been tucked up in bed, surrounded by anxious relatives and doctors.

It is impossible to over-emphasize the psychological approach to the control and treatment of asthma. The intelligent patient is often well aware that his symptoms are aggravated or produced by difficult psychological conditions. To be aware of this is half-way to being protected against their effects. The unintelligent patient is not aware of it, and may be incapable of being taught a proper psychological approach towards his condition. It is in these cases that the impressive "mumbo jumbo" of some specific treatments is most effective, for a time.

In the course of this article I have referred to various theories of the causation of asthma and have rather tended to accept the differentiation into extrinsic (or allergic) asthma and intrinsic asthma. It would, however, be better to use the word, allergic, in a general sense, meaning an altered reactivity to various stimuli, which may be allergic in the sense of the allergists or may be psychological, reflex, metabolic or mechanical. In that sense all asthma is allergic, and it conforms with Hurst's definition of the asthma diatheses (Hurst, 1943), which involves the inherited or acquired possession of an irritable bronchial system, including the medullary centre, the vagal nerve endings and the bronchial musculature and mucosa, which can be stimulated by blood-borne irritants and by reflex and psychic stimuli.

## BIBLIOGRAPHY

- Asthma Clinic, Guy's Hospital (1949). *Advice to Asthmatics*. London: Headley Brosing. L. (1940). "Asthma. Bronchitis following Chronic Upper Respiratory Infection. A five year study of 235 consecutive cases seen in private practice." *J. Amer. med. Ass.* 115, 1434.

- Boland E R and Lintott G A M (1939) Results of Treatment of Rhinorrhoea by Injection of Protein Extracts *Guy's Hosp Rep* 89 108
- Carey R A Harvey A M Howard J E and Winkenwerder W L (1950) The Effect of Adrenocorticotrophic Hormone and Cortisone on the Course of Chronic Bronchial Asthma *Bull Johns Hopk Hosp* 87, 387
- Cecil R L and Loeb R F (1951) *Textbook of Medicine* Philadelphia Saunders
- Clarke C A (1951) Bronchial Asthma Treated by Bilateral Resection of the Vagus A report of six cases *Lancet* 1 438
- Cohen M B (1944) Bronchial Asthma Classification based on Etiological and Pathological factors *Ann intern Med* 20 590
- Conybeare E T and Wills L J (1936) A Criterion for the Treatment of Asthma *Guy's Hosp Rep* 85 138
- Herxheimer H G J (1952) *Management of Bronchial Asthma* London Butterworth
- Howard J E Harvey A McG Carley R A and Winkenwerder W L (1950) Effects of Pituitary Adrenocorticotrophic Hormone (ACTH) on the Hypersensitive State *J Amer med Ass* 144 1347
- Hurst A F (1921) An Address on Asthma *Lancet* 1 1113
- (1943) Asthma in Childhood *Brit med J* 1 403
- Rackemann F M (1931) *Clinical Allergy Especially Asthma and Hay Fever* New York Macmillan
- (1944) Deaths from Asthma *J Allergy* 15 249
- (1951) *Clinical Allergy Especially Asthma* 8th ed Ed by Cecil R L and Loeb R F Philadelphia Saunders
- Schwartz E (1951) Oral Cortisone in Intractable Bronchial Asthma *J Allergy* 22 1
- Steinberg I and Sorrell A H (1941) Occupational Asthma and Vasomotor Rhinitis An outline of some common industries where these symptoms are initiated *NY State J Med* 41 1649
- Tuft L (1946) Bronchial Asthma A critical review *J Allergy* 17, 27
- Unger L (1944) Annual Critical Survey of Recent Literature on Bronchial Asthma *Ann Allergy* 2 49
- (1945) *Bronchial Asthma* Springfield Ill Thomas
- Urbach C and Gottlieb P N (1946) *Allergy* 2nd ed London Kimpton
- Zeller M Randolph T G and Rollins J P (1950) "Adrenocorticotrophic Hormone (ACTH) gross and histologic effects on skin tests and passive transfer" *J Allergy* 8, 163

## CHAPTER 9

### EMPHYSEMA OF THE LUNGS

RONALD V. CHRISTIE

PULMONARY emphysema has been defined as a condition in which the alveoli of the lungs are dilated and their walls over distended. When used in this wide sense the term includes four clinical entities which are probably quite distinct: (1) chronic obstructive or hypertrophic emphysema, (2) senile or atrophic emphysema, (3) acute vesicular emphysema, and (4) localized or compensatory emphysema. There is also the rare condition of acute interstitial emphysema.

### CHRONIC HYPERTROPHIC OR OBSTRUCTIVE EMPHYSEMA

This is the most common form of emphysema and is the type usually referred to when a diagnosis of emphysema of the lungs is made.

*Synonym:* chronic vesicular emphysema

#### Morbid anatomy

#### Gross changes

The thoracic cage is usually enlarged with a characteristic increase in the antero-posterior diameter giving the so-called barrel shaped chest. There is often a moderate degree of kyphosis involving all the thoracic vertebrae. The vertebral cartilage may be thin and compressed anteriorly where there may be tipping. Partly as the result of kyphosis the ribs run horizontally and are widely spaced and their cartilages frequently show ossification or become elongated and less elastic.

The lungs are voluminous and fail to retract when the thorax is opened. The surface of the lung may present a corrugated effect the result of rib indentation and the visceral pleura is thin, flimsy and atrophic. Bullae due to the protrusion of greatly distended alveoli are frequently seen. The lungs are dry and light but they put on pressure not because they are oedematous but because they have lost their elasticity.

On section many of the alveoli can be seen to be dilated and their walls are thin and in places ruptured so that differentiation of the lobules into alveoli, vena and alveolar ducts is impossible. Because the alveolar walls become thin and disappear the air sacs may lose their honeycomb structure and so be deprived of their supporting framework. This is especially true of the surface of the lung where large bullae may be formed. It has been claimed that the rupture of the alveolar wall begins at the so-called alveolar pores but whether or not this is true seems to be of little significance. Of much greater importance is the effect of this destructive process on the air sacs of the lung and several changes have been described. The normal lung lobule or air sac consists of an alveolar duct opening into the atrium around which are placed the alveoli and in health there is a sudden transition from

narrow alveolar duct to wide atrium, so that the inspired air may be thrown as a jet into the more remote parts of this terminal segment. These lobules have been studied in detail by means of serial sections and of specimens injected with celloidin, and in emphysema it has been shown that the lobule is distorted and sometimes distended (Beitzke, 1925, Kountz and Alexander, 1934). The alveolar duct is dilated and funnel-shaped, so that air is no longer thrown into the atrium as a jet. If it could be shown that this jet of air does, in fact, serve a useful purpose in ventilating the lobule, it would be reasonable to suppose that this function would be impaired in emphysema.

Changes in the lumina of the bronchi and bronchioles also occur, but these are not constant. The terminal bronchioles, as well as the alveolar ducts, are usually dilated, and this dilatation may extend up to the smaller bronchi, producing a picture resembling cylindrical bronchiectasis. The state of the medium and larger bronchi is, however, very variable, and depends rather on the degree of bronchitis than on the degree of emphysema, although some atrophy and dilatation are usual in severe cases.

## Blood vessels

The capillaries may appear to be narrow or obliterated and sclerosis of the pulmonary arterioles is common. There is often hypertrophy of the right ventricle, also enlargement of the pulmonary artery and, in advanced cases, right heart failure. Not infrequently this is associated with hypertrophy of the left ventricle, which can usually, but not always, be ascribed to hypertension.

## Distribution of changes

The distribution of emphysematous changes is characteristic. They are most marked in the superficial parts of the lung, particularly at the apices and along the margins. In the early stages they may be confined to these areas but, as the disease progresses, emphysematous changes may be found throughout the lung.

## Loss of elasticity

Changes in the elasticity of the emphysematous lung were first suggested by William Watson in 1795. When the thoracic cage is opened on the post mortem table, the lungs do not collapse, and this alone would appear to be proof that the normal elastic recoil of the lung had been lost. However, more detailed observations on the pulmonary elasticity, made by measuring the force required to inflate and deflate the lung, have given conflicting results in emphysema (Tendeloo, 1925, Loeschke, 1928, Thies, 1932). Considerable changes in the elastic properties of the lung occur after death, and the significance of post mortem measurements of elasticity has been seriously questioned. The fact that the lungs do not collapse when the thorax is opened has been said to mean only that they are over stretched in the same way as a spring may be overstretched and yet maintain much of its elasticity. Histological study of the lungs gets us no further, as there is the same disagreement among morbid anatomists as to whether the elastic fibres are destroyed or merely overstretched, nor do the morbid appearances indicate whether loss of elasticity is an early or a late manifestation of the disease. That loss of elasticity may be complete in this disease has, however, been shown by measurements of the intrapleural pressure during life (see Figs 38-39). The intrapleural pressure is an

# CHRONIC HYPERTROPHIC OR OBSTRUCTIVE EMPHYSEMA

indirect measure of the tension or elastic recoil of the lung and in emphysema it is not always negative as it is in health even at the end of a full inspiration the intrapleural pressure may be the same as that of the atmosphere and it is not altered when the lung is collapsed by a pneumothorax (see Fig 39). This can only mean that in emphysema the elasticity of the lung may be impaired at a comparatively early stage of the disease (Christie, 1934)

Fig 38 — The elasticity of the normal lung. Synchronous tracing of tidal air (lower curve) and intrapleural pressure (upper curve) in a patient with a minimal tuberculous lesion at the right apex. To make the tracing comparable with that shown in Fig 39 the lung has been partially collapsed by a pneumothorax so that the pleural pressure fluctuates around that of the atmosphere. At A a deep breath is taken, maximal distension of the lung corresponds closely to the time when the intrapleural pressure is most negative. The elasticity of the lung is intact. (From Christie and McIntosh 1934)





Fig. 39 — Reduced elasticity of the emphysematous lung. Synchronous tracing of tidal air (lower curve) and intrapleural pressure (upper curve) in a patient with emphysema and a pneumothorax of 40 cubic centimetres on the right side. The pleural pressure fluctuates around that of the atmosphere. At A when a deep breath is taken the pleural pressure becomes only slightly more negative and at the end of inspiration it returns to that of the atmosphere. The lung does not begin to deflate until the pressure becomes positive. These changes indicate that the elasticity of the lung has been lost. (From Christie 1934)

These observations on lung elasticity are not merely of academic interest, for many of the anatomical and functional changes found in emphysema can be explained on the basis of its loss. It is usually assumed that any loss of elasticity is secondary to over-distension. What is not generally realized is that the converse is more likely to be true — the enlargement of the air sacs and the fact that this enlargement is most conspicuous at the periphery of the lung can be explained as being the direct result of loss of elasticity. That this is so is apparent if it is remembered that in health distension of the lung is maintained by traction of the chest wall and the diaphragm and that this traction is reduced or abolished elastic pull of the lungs. If the elastic recoil of the lungs is reduced or abolished as happens in emphysema the thoracic cage will tend to assume the inspiratory position and the lung will be distended. It is probably this mechanism which is at least partly responsible for the enlargement of the chest and low position of the diaphragm in emphysema.



narrow alveolar duct to wide atrium, so that the inspired air may be thrown as a jet into the more remote parts of this terminal segment. These lobules have been studied in detail by means of serial sections and of specimens injected with celloidin, and in emphysema it has been shown that the lobule is distorted and sometimes distended (Beitzke, 1925, Kountz and Alexander, 1934). The alveolar duct is dilated and funnel shaped, so that air is no longer thrown into the atrium as a jet. If it could be shown that this jet of air does, in fact, serve a useful purpose in ventilating the lobule, it would be reasonable to suppose that this function would be impaired in emphysema.

Changes in the lumina of the bronchi and bronchioles also occur, but these are not constant. The terminal bronchioles, as well as the alveolar ducts, are usually dilated, and this dilatation may extend up to the smaller bronchi, producing a picture resembling cylindrical bronchiectasis. The state of the medium and larger bronchi is, however, very variable, and depends rather on the degree of bronchitis than on the degree of emphysema, although some atrophy and dilatation are usual in severe cases.

#### *Blood vessels*

The capillaries may appear to be narrow or obliterated and sclerosis of the pulmonary arterioles is common. There is often hypertrophy of the right ventricle, also enlargement of the pulmonary artery and, in advanced cases, right heart failure. Not infrequently this is associated with hypertrophy of the left ventricle, which can usually, but not always, be ascribed to hypertension.

#### *Distribution of changes*

The distribution of emphysematous changes is characteristic. They are most marked in the superficial parts of the lung, particularly at the apices and along the margins. In the early stages they may be confined to these areas but, as the disease progresses, emphysematous changes may be found throughout the lung.

#### *Loss of elasticity*

Changes in the elasticity of the emphysematous lung were first suggested by William Watson in 1795. When the thoracic cage is opened on the post mortem table, the lungs do not collapse, and this alone would appear to be proof that the normal elastic recoil of the lung had been lost. However, more detailed observations on the pulmonary elasticity, made by measuring the force required to inflate and deflate the lung, have given conflicting results in emphysema (Tendeloo, 1925, Loeschke, 1928, Thies, 1932). Considerable changes in the elastic properties of the lung occur after death, and the significance of post-mortem measurements of elasticity has been seriously questioned. The fact that the lungs do not collapse when the thorax is opened has been said to mean only that they are over stretched in the same way as a spring may be overstretched and yet maintain much of its

may be complete in this disease has, however, been shown by measurements of the intrapleural pressure during life (see Figs 38-39). The intrapleural pressure is an

indirect measure of the tension or elastic recoil of the lung, and in emphysema it is not always negative as it is in health, even at the end of a full inspiration the intrapleural pressure may be the same as that of the atmosphere, and it is not altered when the lung is collapsed by a pneumothorax (see Fig 39). This can only mean that in emphysema the elasticity of the lung may be impaired at a comparatively early stage of the disease (Christie, 1934)

FIG 38 — The elasticity of the normal lung. Synchronous tracing of tidal air (lower curve) and intrapleural pressure (upper curve) in a patient with a minimal tuberculous lesion at the right apex. To make the tracing comparable with that shown in Fig. 39 the lung has been partially collapsed by a pneumothorax, so that the pleural pressure fluctuates around that of the atmosphere. At A a deep breath is taken: maximal distension of the lung corresponds closely to the time when the intrapleural pressure is most negative. The elasticity of the lung is intact. (From Christie and McIntosh 1934)





FIG 39 — Reduced elasticity of the emphysematous lung. Synchronous tracing of tidal air (lower curve) and intrapleural pressure (upper curve) in a patient with emphysema and a pneumothorax of 40 cubic centimetres on the right side. The pleural pressure fluctuates around that of the atmosphere. At A, when a deep breath is taken the pleural pressure becomes only slightly more negative, and at the end of inspiration it returns to that of the atmosphere. The lung does not begin to deflate until the pressure becomes positive. These changes indicate that the elasticity of the lung has been lost. (From Christie 1934)

These observations on lung elasticity are not merely of academic interest, for many of the anatomical and functional changes found in emphysema can be explained on the basis of its loss. It is usually assumed that any loss of elasticity is secondary to over-distension. What is not generally realized is that the converse is more likely to be true: the enlargement of the air sacs, and the fact that this

chest wall and the diaphragm, and that this traction is counterbalanced by the elastic pull of the lungs: if the elastic recoil of the lungs is reduced or abolished, as happens in emphysema, the thoracic cage will tend to assume the inspiratory position and the lung will be distended. It is probably this mechanism which is at least partly responsible for the enlargement of the chest and low position of the diaphragm in emphysema.

## EMPHYSEMA OF THE LUNGS

narrow alveolar duct to wide atrium, so that the inspired air may be thrown as a jet into the more remote parts of this terminal segment. These lobules have been studied in detail by means of serial sections and of specimens injected with celloidin, and in emphysema it has been shown that the lobule is distorted and sometimes distended (Beitzke, 1925, Kountz and Alexander, 1934). The alveolar duct is dilated and funnel shaped, so that air is no longer thrown into the atrium as a jet. If it could be shown that this jet of air does, in fact, serve a useful purpose in ventilating the lobule, it would be reasonable to suppose that this function would be impaired in emphysema.

Changes in the lumina of the bronchi and bronchioles also occur, but these are not constant. The terminal bronchioles, as well as the alveolar ducts, are usually dilated, and this dilatation may extend up to the smaller bronchi, producing a picture resembling cylindrical bronchiectasis. The state of the medium and larger bronchi is, however, very variable, and depends rather on the degree of bronchitis than on the degree of emphysema, although some atrophy and dilatation are usual in severe cases.

### *Blood vessels*

The capillaries may appear to be narrow or obliterated and sclerosis of the pulmonary arterioles is common. There is often hypertrophy of the right ventricle, also enlargement of the pulmonary artery and, in advanced cases, right heart failure. Not infrequently this is associated with hypertrophy of the left ventricle, which can usually, but not always, be ascribed to hypertension.

### *Distribution of changes*


The distribution of emphysematous changes is characteristic. They are most marked in the superficial parts of the lung, particularly at the apices and along its margins. In the early stages they may be confined to these areas but, as the disease progresses, emphysematous changes may be found throughout the lung.

### *Loss of elasticity*

Changes in the elasticity of the emphysematous lung were first suggested by William Watson in 1795. When the thoracic cage is opened on the post mortem table, the lungs do not collapse, and this alone would appear to be proof that the normal elastic recoil of the lung had been lost. However, more detailed observations on the pulmonary elasticity, made by measuring the force required to inflate and deflate the lung, have given conflicting results in emphysema (Tendeloo, 1925, Loeschke, 1928, Thies, 1932). Considerable changes in the elastic properties of the lung occur after death, and the significance of post mortem measurements of elasticity has been seriously questioned. The fact that the lungs do not collapse when the thorax is opened has been said to mean only that they are over stretched in the same way as a spring may be overstretched and yet maintain much of its elasticity. Histological study of the lungs gets us no further, as there is the same disagreement among morbid anatomists as to whether the elastic fibres are destroyed or merely overstretched, nor do the morbid appearances indicate whether loss of elasticity is an early or a late manifestation of the disease. That loss of elasticity may be complete in this disease has, however, been shown by measurements of the intrapleural pressure during life (see Figs 38-39). The intrapleural pressure is an

indirect measure of the tension or elastic recoil of the lung and in emphysema it is not always negative as it is in health, even at the end of a full inspiration the intrapleural pressure may be the same as that of the atmosphere and it is not altered when the lung is collapsed by a pneumothorax (see Fig 39). This can only mean that in emphysema the elasticity of the lung may be impaired at a comparatively early stage of the disease (Christie, 1934)

FIG 38 — The elasticity of the normal lung  
Synchronous tracing of tidal air (lower



At A a deep breath is taken maximal distension of the lung corresponds closely to the time when the intrapleural pressure is most negative. The elasticity of the lung is intact (From Christie and McIntosh 1934)





FIG 39 — Reduced elasticity of the emphysematous lung. Synchronous tracing of tidal air (lower curve) and intrapleural pressure (upper curve) in a patient with emphysema and a pneumothorax of 40 cub c centimetres on the right side. The pleural pressure fluctuates around that of the atmosphere. At A when a deep breath is taken the pleural pressure becomes only slightly more negative and at the end of inspiration it returns to that of the atmosphere. The lung does not begin to deflate until the pressure becomes positive. These changes indicate that the elasticity of the lung has been lost (From Christie 1934)

These observations on lung elasticity are not merely of academic interest, for many of the anatomical and functional changes found in emphysema can be explained on the basis of its loss. It is usually assumed that any loss of elasticity is secondary to over distension. What is not generally realized is that the converse is more likely to be true: the enlargement of the air sacs, and the fact that this enlargement is most conspicuous at the periphery of the lung, can be explained as being the direct result of loss of elasticity. That this is so is apparent if it is remembered that in health distension of the lung is maintained by traction of the chest wall and the diaphragm and that this traction is counterbalanced by the elastic pull of the lungs. If the elastic recoil of the lungs is reduced or abolished as happens in emphysema the thoracic cage will tend to assume the inspiratory position and the lung will be distended. It is probably this mechanism which is at least partly responsible for the enlargement of the chest and low position of the diaphragm in emphysema.

Loss of elasticity could also explain the over-distension of air sacs, and the bulla formation which occurs on the surface of the lung. With loss of elasticity the expanding force is no longer equally distributed, and equal expansion in different parts of the lung should not be expected. In fact, the greatest expansion should occur where the force is applied, and this is at the surface of the lung. With each inspiration one would expect the superficial air sacs to be strained and stretched to a greater extent than are the air sacs deep in the lung, and, with this process going on for months and years, it is easy to understand the over-distension of air sacs and the bulla formation which occur mainly on the surface of the lung.

### Functional pathology

#### *Haemo-respiratory exchange*

The most important disturbance of function in emphysema is an inability to ventilate the blood, and this is usually apparent many years before the patient reaches the pathologist. It is retention in the blood of carbon dioxide which is believed to be responsible for the dyspnoea on exertion that is an early manifestation of this disease. Cyanosis may appear later, and is due to deficient oxygenation of the blood. Many explanations of this impairment of aeration of the blood in emphysema have been given, but most of them are based only on conjecture and can be shown to be unsound.

*Theory of muscular inadequacy*—It has been suggested that the respiratory musculature is incapable of properly ventilating the lungs, for the same reasons that the vital capacity and chest expansion are diminished, but the burden of blame cannot be placed on the muscles of respiration, since ventilation of the blood is impaired even when the amount of air drawn into the lungs with each breath and during each minute is greater than normal. Inability to hyperventilate on exercise may be a factor in the production of dyspnoea, but the main fault must lie in the lungs, because for some reason the air which is inspired does not properly ventilate the blood.

*Suggested barrier to the transfer of gases*—It is obvious that the defect cannot be due to any barrier to the transfer of gases across the pulmonary epithelium, or to insufficiency of functioning lung, since, in these circumstances, the tension of carbon dioxide would increase in the blood but not in the alveoli. The fact that changes in the blood are associated with proportionate changes in the alveolar air

1947). Although inability to increase adequately the cardiac output on exercise may be a factor in decreasing the exercise tolerance, it cannot be responsible for the impairment of gaseous exchange in the lungs.

*Inadequate ventilation of air sacs*—A much more reasonable explanation, and one which is widely accepted on the Continent, is that the air sacs are not properly ventilated—partly because they are big, but mainly because the alveolar ducts are obstructed (Bastle, 1975; Kountz and Alexander, 1934). It is true that in

has been named the stratification theory, so called because it assumes a gradient of ventilation in the lobule those parts next to the alveolar duct being over ventilated and the parts further away being under ventilated. This sounds reasonable but the validity of this theory of stratification depends on the assumption that in health there must be circulation of air in the air sacs and alveoli and that diffusion alone is insufficient to maintain adequate exchange of gases in the lobules. The diffusion of gases is governed by well defined laws and it can be calculated that even if the lobule is 1 centimetre in diameter diffusion alone would ensure the transport of gases across the alveolar space with a gradient of less than 1 millimetre of mercury. Unequal ventilation within the lobule cannot account for the extreme impairment of gaseous exchange observed in emphysema in which the pressure of carbon dioxide may be raised by 30 or 40 millimetres of mercury (Christie 1939 Rauwerda 1946).

*Increased volume of the lungs*—Increase in the volume of the lungs has been suggested as the cause of impaired ventilation. Provided that ventilation is distributed evenly throughout the lungs an increase in lung volume does not significantly impair haemo respiratory exchange (Bates and Christie 1950).

*Unequal distribution of ventilation*—An increase in the respiratory dead space has been suggested as the cause of impaired haemo respiratory exchange but there is ample evidence that any increase in the anatomical dead space—the space occupied by the respiratory passages down to the bronchioles—is too small to be of practical significance (Hoover and Gammon 1915). I shall not enter into a discussion on the meaning of the physiological dead space. Most physiologists will agree that those who claim an increase in physiological dead space in emphysema are merely stating in a complicated way that the distribution of ventilation is unequal. In the normal individual the distribution of ventilation is remarkably even throughout the lungs and consequently samples of alveolar air taken after expirations of varying depths are more or less uniform in composition. This is not the case in severe emphysema the deeper the patient expires the higher is the carbon dioxide content and the lower is the oxygen content of the alveolar air. These cases show the apparent paradox that although the amount of air inspired is normal or even increased many of the alveoli are under ventilated (Roelsen 1914 and 1939). Furthermore the respiratory quotient of samples of air taken at the end of a forced expiration may be very low and this can only mean that the sample has come from a part of the lung which is grossly under ventilated. There is also ample evidence that the ability to produce a homogeneous mixture of gases in the lung is impaired in emphysema (Cournand and his colleagues 1941 Bates and Christie 1950 Briscoe Blacklake and Rose 1951) this is due to over ventilation of certain parts of the lung at the expense of other parts.

*Pathological respiratory dead space*—The most likely explanation of this inequality in the distribution of ventilation is that much of the inspired air is deflected to the enlarged air sacs and bullae the walls of which are avascular and which are therefore not in contact with the pulmonary circulation (Christie 1939 and 1944 Riley and Cournand 1949). This wastage of ventilation on functionless air sacs cannot be defined as wastage due to increase in physiological dead space for there is nothing physiological about an emphysematous bulla. Nor can

## EMPHYSEMA OF THE LUNGS

it reasonably be called increase in anatomical dead space. If a definition is wanted, I should prefer to give as the cause of impaired haemo respiratory exchange in emphysema the formation of a pathological respiratory dead space, consisting of bullae and other air sacs which are not in intimate contact with the pulmonary circulation. In severe emphysema much of the inspired air is wasted by entering this pathological dead space, so that those areas of the lung which contain alveoli well perfused with blood are under ventilated. The natural result of this abnormal distribution of ventilation in the lungs is imperfect aeration of the blood.

### *Dyspnoea*

The cause of dyspnoea in emphysema is clearly stimulation of the respiratory centre by the carbon dioxide retained in the blood. The chronicity of the disease allows the patient with emphysema full scope for chemical compensatory changes in the blood, and the inability to eliminate carbon dioxide may be efficiently compensated by an increase in the bicarbonate reserve (carbon dioxide combining power). This compensatory mechanism is sufficient to ensure a normal acidity of the blood while the patient is resting, but any increase in metabolism—such as occurs during exercise or during a violent attack of coughing—is immediately associated with an increase in the blood carbon dioxide and stimulation of the respiratory centre.

The emphysematous patient is, indeed in an unfortunate position. Both inspiration and expiration have to be executed by unnatural respiratory efforts, and a considerable proportion of the air which is inspired is wasted by not coming into proper contact with the pulmonary blood. At rest the patient can just make ends meet with the help of compensatory changes in the blood, carbon dioxide elimination and oxygen absorption keep up with the metabolic demands. When the metabolic demands are increased by exercise, this balance breaks down. The patient is unable adequately to increase the ventilation of the blood. Carbon dioxide accumulates and anoxaemia increases, so that true chemical dyspnoea due to acid stimulation of the respiratory centre, appears. In the absence of bronchospasm or heart failure, there is no orthopnoea because there is no pulmonary congestion which is the cause of true orthopnoea.

### *The pulmonary circulation*

Extensive changes in the pulmonary vascular bed can be seen at autopsy, and it is usually assumed that there is sufficient vascular destruction to impair the circulation through the lungs. This belief is not based on any quantitative measurement, because it is not possible to estimate with the microscope even the approximate proportion of capillaries that have been destroyed. It should not be forgotten that the pulmonary vascular bed has an enormous reserve, which is well illustrated in modern thoracic surgery, when the whole of one lung may be removed and yet there may be no evidence of impaired pulmonary blood flow or of strain on the right side of the heart. A demonstration of capillary damage does not in itself, therefore, prove any embarrassment to the pulmonary circulation. Much more suggestive of this are the changes frequently found in the pulmonary artery and right ventricle, as revealed by cardiac catheterization. The pressure within the pulmonary artery is usually increased and this suggests increased resistance to the flow of blood through the lungs. The cardiac output, however,

is also increased even under resting conditions, and the most likely explanation is that, under the stimulus of anoxia, the right ventricle maintains an increased cardiac output in spite of increased resistance in the pulmonary vascular bed (McMichael and Sharpey Schafer, 1944, Richards, 1947). On exercise there is a further large increase in pulmonary arterial pressure but, in spite of this, the increase in cardiac output may be negligible (Hickman and Cargill, 1947, Riley and his co workers, 1948). Inability to increase the cardiac output on exercise may be of considerable functional significance in emphysema, but these observations, which are recent, will have to be confirmed and amplified before they can be applied to the interpretation of symptoms in this disease.

*Electrocardiographic changes*—Much has been written on the electrocardiographic changes in emphysema and in chronic cor pulmonale, there are right axis deviation and usually a prominent P-wave in lead II. Praecordial leads may also show the changes characteristic of right ventricular hypertrophy (Myers, Klein and Stafer, 1948).

*Lymphatic circulation*—The lymphatic circulation in emphysema has received but scant attention. The peripheral lymphatic vessels may show cellular infiltration, and the characteristic pigmentation of the lung may reflect deficient lymphatic drainage.

#### *The blood*

The carbon dioxide combining power is usually increased in patients who are incapacitated by dyspnoea, and may be raised to over 100 volumes per cent. This increase is a response to the retention of carbon dioxide, and represents the mechanism by which the acid-base balance is maintained, although the alkali reserve is increased, these patients are in a state of compensated gaseous acidosis. The oxygen tension and oxygen saturation of the arterial blood are reduced. Less commonly, in patients with severe cyanosis, there is polycythaemia.

Riley and Cournand (1949) and other authors have recently emphasized the value of arterial-blood gas analysis in estimating the impairment of respiratory function in this disease. Both the laboratory procedure and the interpretation of results in terms of "ventilation-perfusion relationship" are complex, to say the least, but the basic principles of approach are simple. The arterial blood

blood is in equilibrium with the carbon dioxide of the alveolar air, then the pressure of oxygen which should exist in the alveolar air and in the arterial blood, if a similar equilibrium existed with regard to oxygen, can be calculated from the respiratory quotient. The difference between this "ideal" oxygen pressure and the oxygen pressure which is actually found in the arterial blood is expressed as the "alveolar-arterial gradient". This gradient is increased in emphysema and is assumed to reflect the passage of unaerated blood through the lungs. It can also be expressed in terms of "venous admixture".

#### *Clinical picture*

##### *History and progress*

With rare exceptions there is a history of chronic bronchitis or asthma of several



it reasonably be called increase in anatomical dead space. If a definition is wanted, I should prefer to give as the cause of impaired haemo-respiratory exchange in emphysema the formation of a pathological respiratory dead space, consisting of bullae and other air sacs which are not in intimate contact with the pulmonary circulation. In severe emphysema much of the inspired air is wasted by entering this pathological dead space, so that those areas of the lung which contain alveoli well perfused with blood are under-ventilated. The natural result of this abnormal distribution of ventilation in the lungs is imperfect aeration of the blood.

### *Dyspnoea*

The cause of dyspnoea in emphysema is clearly stimulation of the respiratory centre by the carbon dioxide retained in the blood. The chronicity of the disease allows the patient with emphysema full scope for chemical compensatory changes in the blood, and the inability to eliminate carbon dioxide may be efficiently compensated by an increase in the bicarbonate reserve (carbon dioxide combining power). This compensatory mechanism is sufficient to ensure a normal acidity of the blood while the patient is resting, but any increase in metabolism—such as occurs during exercise or during a violent attack of coughing—is immediately associated with an increase in the blood carbon dioxide and stimulation of the respiratory centre.

The emphysematous patient is, indeed, in an unfortunate position. Both inspiration and expiration have to be executed by unnatural respiratory efforts, and a considerable proportion of the air which is inspired is wasted by not coming into proper contact with the pulmonary blood. At rest the patient can just make ends meet, with the help of compensatory changes in the blood, carbon dioxide elimination and oxygen absorption keep up with the metabolic demands. When the metabolic demands are increased by exercise, this balance breaks down. The patient is unable adequately to increase the ventilation of the blood. Carbon dioxide accumulates and anoxaemia increases, so that true chemical dyspnoea, due to acid stimulation of the respiratory centre, appears. In the absence of bronchospasm or heart failure, there is no orthopnoea, because there is no pulmonary congestion, which is the cause of true orthopnoea.

### *The pulmonary circulation*

Extensive changes in the pulmonary vascular bed can be seen at autopsy, and it is usually assumed that there is sufficient vascular destruction to impair the circulation through the lungs. This belief is not based on any quantitative measurement, because it is not possible to estimate with the microscope even the approximate proportion of capillaries that have been destroyed. It should not be forgotten that the pulmonary vascular bed has an enormous reserve, which is well illustrated in modern thoracic surgery, when the whole of one lung may be

artery and right ventricle  
within the pulmonary a  
resistance to the flow of

is also increased even under resting conditions and the most likely explanation is that under the stimulus of anoxia the right ventricle maintains an increased cardiac output in spite of increased resistance in the pulmonary vascular bed (McMichael and Sharpey Schafer 1944 Richards 1947). On exercise there is a further large increase in pulmonary arterial pressure but in spite of this the increase in cardiac output may be negligible (Hickman and Cargill 1947 Riley and his co workers 1948). Inability to increase the cardiac output on exercise may be of considerable functional significance in emphysema but these observations which are recent will have to be confirmed and amplified before they can be

the electrocardio  
there are right axis  
rdial leads may also

show the changes characteristic of right ventricular hypertrophy (Myers Klein and Stafer 1948)

*Lymphatic circulation*—The lymphatic circulation in emphysema has received but scant attention. The peripheral lymphatic vessels may show cellular infiltration and the characteristic pigmentation of the lung may reflect deficient lymphatic drainage.

#### *The blood*

The carbon dioxide combining power is usually increased in patients who are incapacitated by dyspnoea and may be raised to over 100 volumes per cent. This increase is a response to the retention of carbon dioxide and represents the mechanism by which the acid base balance is maintained although the alkali reserve is increased these patients are in a state of compensated gaseous acidosis. The oxygen tension and oxygen saturation of the arterial blood are reduced. Less commonly in patients with severe cyanosis there is polycythaemia.

Riley and Courmand (1949) and other authors have recently emphasized the value of arterial blood gas analysis in estimating the impairment of respiratory function in this disease. Both the laboratory procedure and the interpretation of results in terms of ventilation perfusion relationship are complex to say the least but the basis of this new line of approach is as follows. The measurements which have to be made are the pressure of carbon dioxide and of oxygen in the arterial blood and also the respiratory quotient of the expired air. If the assumption be made that in health as well as in disease the carbon dioxide of the arterial blood is in equilibrium with the carbon dioxide of the alveolar air then the pressure of oxygen which should exist in the alveolar air and in the arterial blood if a similar equilibrium existed with regard to oxygen can be calculated from the respiratory quotient. The difference between this ideal oxygen pressure and the oxygen pressure which is actually found in the arterial blood is expressed as the alveolar arterial gradient. This gradient is increased in emphysema and is assumed to reflect the passage of unaerated blood through the lungs. It can also be expressed in terms of venous admixture.

#### *Clinical picture*

##### *History and progress*

With rare exceptions there is a history of chronic bronchitis or asthma of several

years' duration, but the cardinal symptom of emphysema itself is excessive dyspnoea on exertion. The onset is usually slow, but it may be abrupt, in which case there is commonly a history of antecedent acute respiratory infection. Over a period of months or years the patient complains of increasing breathlessness, which may diminish from time to time with any improvement in the associated cough or broncho-spasm. For this reason patients often state that they are better during the summer months.

Cyanosis is usually inconspicuous, even in patients who are very dyspnoeic, but it may be severe, particularly in the terminal stages of the disease, when there may be secondary polycythaemia and clubbing of the fingers. The primary cause of this cyanosis is inability to aerate the blood, but peripheral stasis and polycythaemia may also be important factors.

The progress of the disease is usually insidious, but rarely it is rapid, and the patient may be incapacitated with severe dyspnoea within a year or two of the onset. Usually there is a slow progression of symptoms over many years, and, if the patient does not die of intercurrent infection, the symptoms and signs of cor pulmonale and right sided heart failure may supervene. The danger of right-heart failure is, however, unpredictable, some patients may be incapacitated by dyspnoea for years without any evidence of cardiac impairment, whereas others are first seen in heart failure, and will only admit a brief period of moderate dyspnoea. The most common complication is said to be arterial hypertension. Spontaneous pneumothorax, due to rupture of the flimsy visceral pleura may occur less frequent complications are haemoptysis and interstitial emphysema. Failure of vision due to papilloedema is very rare, but it occasionally occurs, usually in patients who have developed secondary polycythemia.

### Physical signs

Although the significance of physical signs in the diagnosis of emphysema has been disputed, it is nevertheless true that in the advanced stages of the disease certain signs are usually present

**Inspection**—The chest may be barrel shaped, thoracic kyphosis, loss of pulmonary elasticity and expiratory obstruction contribute to this deformity. The relative importance of these factors varies from patient to patient, and it is not surprising that enlargement of the thoracic cage may be marked in some and inconspicuous in others, although they may be incapacitated by the disease. If the enlargement of the thorax is considerable, the intercostal spaces are usually wide and the ribs ter. The supraclavicular Expansion of the cl elevation of the thorax, heaving in character Expiration is often visibly prolonged

These phenomena are usually easily demonstrable, and they reflect the disorganization of breathing which accompanies loss of pulmonary elasticity and expansion of the thoracic cage. The diaphragm, because it is no longer pulled upwards by the lungs, descends and becomes less convex, and its movement on inspiration is greatly diminished. The intercostal muscles will also be unable to perform their proper function, because they are concerned with moderate inspiration and the chest is already expanded when inspiration begins, in order to expand

the chest further, the accessory muscles will have to be used, and these raise the front of the chest as a whole in a "heaving" manner, such as occurs in a normal individual towards the end of a forced inspiration. Thus inspiration has to be performed by means of an unnatural inspiratory effort, and the patient with emphysema will encounter even greater difficulty in deflating his lung. Normally, expiration is largely, if not wholly, a passive act, the thoracic cage is pulled inwards by the elastic recoil of the lung. With loss of elasticity there must be loss of elastic recoil, so that if the lung is to be deflated it has to be squeezed. The respiratory musculature was not built for this task, and the intercostal muscles have to be assisted by the accessory muscles of expiration. The muscles of the abdominal wall can sometimes be felt to contract on expiration, which is prolonged as it is in other conditions, such as asthma and tracheal obstruction, in which the lungs are compressed by an active muscular effort. With so extensive an impairment of both inspiration and expiration, it is not surprising that the vital capacity and chest expansion are reduced.

The apical impulse is seldom visible or palpable, since a considerable thickness of inelastic lung is interposed between it and the chest wall. In contrast, epigastric pulsation is often marked, presumably because of the low position of the diaphragm.

*Palpation*—On palpation the vocal fremitus may be unaltered, but often it is diminished, possibly because bullae and enlarged air sacs are poor conductors of sound.

*Percussion*—On percussion resonance is increased. It is probably the superficial over distension of air sacs and formation of bullae which cause the remarkable hyper-resonance of the percussion note, which is sometimes asymmetrical and which may be more resonant than over a pneumothorax. The limits of lung resonance are increased because the size of the lung is increased, particularly in the region of its margins. The potential gap between the chest wall and the heart becomes filled with emphysematous lung. This space can be clearly seen in the lateral view by radiography, and it explains a variety of physical signs. The absence of the apical impulse, loss of cardiac dullness, and distant heart sounds necessarily follow. Similarly, the loss of liver dullness must in part be caused by expansion of the lung into the pleural sinuses, although the low position of the diaphragm is, of course, another factor.

*Auscultation*—On auscultation the breath sounds are faint, particularly at the bases, but they may be increased in intensity by the concomitant bronchitis or broncho spasm, when coarse or sibilant rhonchi may be heard. The factors leading to suppression of breath sounds are probably the same as those which cause diminution in the conduction of the spoken voice, but another theory has been proposed which may or may not play a part. It is claimed that vesicular breathing in health is caused by the jet of air, which is thrown from the narrow alveolar duct into the wide atrium. In emphysema, the alveolar duct is dilated so that this sudden change in calibre is lost, it is suggested that it is for this reason that vesicular breathing is faint or absent (Beitzke, 1925).

Expiration is prolonged, vocal resonance is usually diminished and the heart sounds are faint for reasons which are described above. The second sound over the pulmonary area may be accentuated or reduplicated.

## Diagnosis

### *Interpretation of physical signs*

The above are the physical signs of emphysema. Although they are usually present in advanced cases, they may be absent in patients suffering from the disease and some are not uncommonly present in patients without emphysema. Considerable confusion exists, even among physicians with wide clinical experience, as to the diagnostic value of these signs, and disagreement is perhaps greatest on the relationship of the barrel shaped chest to emphysema. Some believe, and teach, that the physical signs associated with the barrel shaped chest, if definite, necessarily mean emphysema, although they admit that the lesion may neither progress nor lead to symptoms. Others believe that the "barrel chest phenomenon" may occur quite independently of emphysema, and that in these circumstances it is of no significance. Such a divergence of opinion as to the diagnostic criteria naturally makes for scepticism, and attempts to correlate the clinical diagnosis of emphysema with the facts as revealed by necropsy have led to further doubts. It has been stated that only a small proportion of cases diagnosed at necropsy have been recognized during life and it has also been shown that many subjects with barrel shaped chests furnish no evidence of emphysema when they come to the post mortem table (Cabot, 1927, Davidson, 1936, Roelsen, 1938). For this reason some writers have gone so far as to say that the physical signs of emphysema are meaningless.

Cabot states that of 12 cases diagnosed as emphysema at the Massachusetts General Hospital, only 3 showed emphysema *post mortem*, and that of 153 cases demonstrated *post mortem* only 7 had been recognized during life. My own experience is rather different and is supported by a series of 72 cases diagnosed on the post-mortem table as generalized hypertrophic emphysema (Christie, 1944). All of these came from the medical wards of the London Hospital, but an accurate and detailed case-history was not always available, for some patients were moribund on admission. A history of chronic cough was given in 68 cases and in a quarter of this number it was associated with mild or severe asthma. Dyspnoea on exertion was present in 35 of the 45 case histories in which mention was made of the presence or absence of this symptom. In the majority of cases, however, the physical signs of emphysema were not observed. In 16 cases there was enough evidence to suggest an "emphysematous chest," but in only 6 cases could this evidence be said to be complete.

This series of cases suggests that a more reliable guide to a diagnosis of emphysema than the physical signs is a history of chronic cough or asthma, associated with dyspnoea on exertion. I have therefore investigated another series of 25 patients who gave such a history, and in whom no cause of dyspnoea other than emphysema could be discovered. In 23 of them two or more of the physical signs


of "emphysematous chest"  
no  
ases  
ents

In 9 of them the chest might have been said to be emphysematous and 4 presented the classical physical signs of emphysema

The conclusion that can be drawn is that the signs of emphysema are of significance but are unreliable they may be absent in patients suffering from the disease and are not uncommonly present in patients without emphysema A diagnosis of emphysema based only on physical signs may be fallacious since these depend largely on the barrel chest phenomenon which may occur independently of this disease In my opinion the diagnosis should only be considered to be definite when undue dyspnoea on exertion of insidious onset appears in a patient who has some of the physical signs of emphysema together with chronic bronchitis or asthma The absence either of dyspnoea or of physical signs should cast doubt on the diagnosis but should not exclude it the more physical signs present in a patient with chronic bronchitis or asthma the greater is the likelihood that he has emphysema I must confess however that in the absence of dyspnoea I am always uncertain as to diagnosis and prognosis but when dyspnoea not due to broncho-spasm appears in a patient with chronic bronchitis I am very seldom in doubt A large series of cases will have to be followed up over many years before the necessary criteria for early diagnosis can be established Until this has been done I consider that the terms emphysematous and emphysematous chest should be discarded for it is their use which has been partly responsible for the lack of progress in the understanding of emphysema since its classical description over a century ago by Laennec These terms are quite misleading and should be replaced by some such expression as barrel chest or barrel chest phenomenon reserving the diagnosis of emphysema for those patients who have the symptoms as well as the signs of the disease I believe that if this were done the chances of progress would be increased particularly with regard to the use of special diagnostic procedures

#### *Special diagnostic procedures*

*X ray examination*—The appearances in emphysema are said to be characteristic



vertebrae The movements of the diaphragm on inspiration and expiration as well as those of the sternum are usually diminished The space between the sternum and the heart is almost invariably increased The heart may appear to lie vertically with a prominent pulmonary conus and exaggeration in size of the pulmonary arterial shadows at the lung hila The medium sized vessels tend to stand out more clearly but are smaller in calibre than normal the smaller peripheral vessels may also be more clearly visible than normal The lung fields may show increased translucency particularly at the lung bases Individual bullae may be seen usually at the apices of the lungs

It has been claimed that the radiological picture is a reliable guide to emphysema but attempts to establish the validity of this claim have been unsuccessful In patients with florid emphysema a correct radiological diagnosis based on a postero-anterior film can be made in about 80 per cent of cases but if a control




FIG 40—Radiograph of the chest in a case of chronic obstructive emphysema. The intercostal spaces are wide and there is increased translucency of the lung fields especially at the bases. The hilar shadow is prominent and the heart long and narrow. The diaphragm is low and irregular in outline.

group of elderly individuals be taken, about 20 per cent will be wrongly diagnosed as having emphysema. A more reliable assessment may be made from films taken in the antero-posterior and lateral positions, in both inspiration and ex-

drawn from these observations is that the x-ray film may provide useful additional evidence, but taken alone it cannot exclude or establish a diagnosis of emphysema.

*Vital capacity and maximum breathing capacity.*—The vital capacity is usually reduced in emphysema. Out of 30 patients with a clinical diagnosis of emphysema, I have found that in 25 cases the vital capacity was less than 70 per cent of the normal standard as calculated from the surface area, and in 15 cases it was less than 60 per cent. In a control series of 30 patients of the same age distribution, without evidence of emphysema, the vital capacity was less than 70 per cent of the normal standard in 11, and less than 60 per cent in 3 cases. There are many diseases of the lung other than emphysema which reduce the vital capacity, and it is clear that this test alone can be of little value in diagnosis.

It is more easy to measure the chest expansion than the vital capacity, but unfortunately this also diminishes with age. In my experience, about 60 per cent of patients with emphysema have a chest expansion of less than 1 inch, but I

have seen it reduced to  $\frac{1}{2}$  inch in a man aged 65 years in whom there was no other evidence to suggest pulmonary or cardiovascular disease

The maximal breathing capacity is also reduced in emphysema, but as a diagnostic test it suffers from the same drawbacks as does the vital capacity

*Lung volume and its subdivisions*—An important sequel to the dilatation of the air sacs in emphysema is the increase of the volume of the lungs as a whole which is usually obvious on the post mortem table the large fluffy lungs strongly suggest that during life there was an increase in the amount of air which they contained This amount can be measured, and early in the present century several investigators showed that there is usually an increase in the residual air in emphysema, but the residual air is the amount of air left after a forced expiration, and it is well known that patients with emphysema may have great difficulty in forcibly emptying their lungs A more significant measurement is the amount of air in the lungs at the end of an ordinary quiet expiration—the functional residual air—for this is their true content during life, whereas the residual air is the result of respiratory gymnastics and may have no relation to the normal size of the lungs It has been shown that the quantity of the functional residual air is increased in emphysema, but this increase is not always proportionate to the severity of the disease In some patients it may be inconspicuous, when contrasted with the fairly wide range found in normal individuals In order to escape the need of normal standards which allow for body-size and other factors, it was

If a measurement is required to assist in the diagnosis or investigation of emphysema this ratio is probably the most accurate measurement available, but it should not be forgotten that the ratio is quite an artificial figure and that any increase is due to two quite distinct factors, (1) the enlargement of the lung and (2) the inability of the patient to take a deep breath There has recently been a tendency to assume that an increase in this ratio, because it is a precise measurement, is a more accurate diagnostic sign than is clinical evidence of the disease This has not been shown to be true

*Tolerance to carbon dioxide*—It has for long been known that patients with emphysema react sluggishly to the inhalation of carbon dioxide Although the fundamental cause of this impairment is unknown it is probably related to the high bicarbonate content of the blood and to inefficient ventilation in the lungs Recent improvements in technique allow the measurement of carbon dioxide tolerance to be made more easily than before, and it has been suggested that this reaction may be of value in the diagnosis of emphysema Even if the reliability of this test is established it is unlikely to become a routine diagnostic procedure, because special equipment is required (Donald and Christie, 1949)

*The mixing of gases in the lung*—In emphysema, the efficiency with which gases are mixed in the lung is grossly impaired (Cournand and his colleagues, 1941; Meneely and Kaltreider, 1949; Bates and Christie, 1950) By means of the katharometer accurate measurements of the rate of gaseous mixing are now



possible, and it is likely that tests of this type will contribute to the problem of early diagnosis

### Aetiology

Numerous hypotheses have been advanced to explain the cause of chronic hypertrophic or obstructive emphysema, but most are based on conjecture and are incapable of experimental proof or analysis. Certain facts, however, stand out clearly—almost all patients suffering from this disease are men of middle age or over who give clear evidence of chronic respiratory obstruction due to asthma or chronic bronchitis, or to persistent coughing from pulmonary tuberculosis, bronchiectasis, dusty occupations, or any other cause. It has also been shown, both clinically and experimentally, that tracheal or bronchial obstruction, if sufficiently pronounced and prolonged, usually leads to emphysema. This correlation between respiratory obstruction and emphysema cannot be denied, but the mechanism by which obstruction leads to the changes characteristic of emphysema is in considerable dispute.

### *The expiratory theory*

According to the expiratory theory, respiratory obstruction due to cough or bronchospasm will hinder exit of air from the lungs to a greater extent than entry, since the muscles of inspiration are more powerful than those of expiration. As a result air will accumulate in the alveoli with subsequent distension and rupture of their walls. This theory is simple and appears to be reasonable, but it does not take into account the immunity from emphysema of young persons who suffer from asthma. The theory is also based partly on the false assumption that the lungs as a whole are grossly over-distended. It is true that the amount of air in the emphysematous lung is usually increased, but this increase is not always proportionate to the severity of the disease, nor is the volume of the emphysematous lung at rest as great as the volume of a normal lung in full inspiration; in most patients with emphysema the lung as a whole is not over-distended any more than a normal lung is over-distended when a deep breath is taken. Incomplete emptying of the alveoli due to respiratory obstruction may be a factor in the production of emphysema, but it cannot be the only one.

### *The inspiratory theory*

This theory claims that over-distension is produced because too much air is drawn into the lungs by the increased inspiratory effort which precedes coughing, and which may be associated with asthma, with manual labour and with occupations such as glass-blowing and the blowing of wind instruments. There is no evidence that these occupations or increased inspiratory effort due to any other cause play a part in the production of emphysema (Christie, 1939).

### *Impairment of pulmonary nutrition*

In other diseases there may be more extensive a

## CHRONIC HYPERTROPHIC OR OBSTRUCTIVE EMPHYSEMA

emphysema although the lung does not become emphysematous. For these reasons it is improbable that the vascular changes in the lung are of any aetiological significance.

*Degenerative or atrophic pulmonary changes*

It has been suggested that these changes are secondary and evade the issue is known to be involved.

...count the many aetiological factors

### Alveolar stress

Another theory is that the primary cause is increased stress and strain on the walls of the alveoli. Almost all cases of emphysema have a long standing history of chronic bronchitis or asthma or both and this is also true of emphysema in horses and cattle. The effects of cough and bronchial spasm on the lung as well as their mechanism are therefore of primary importance in any discussion on the aetiology of this disease.

The mechanism of coughing is simple. There is a short inspiratory effort which by no means fills the chest to full capacity; the glottis is then closed and the lungs are forcibly compressed by all the muscles of expiration so that the pressure of air in the alveoli may rise to over 50 millimetres of mercury (Kountz and Alexander 1934; Rasmussen and Adams 1942). The glottis is then opened and the strain on the ribs, the pressure on the pleura and the pressure in the alveoli are suddenly released. In the case of the alveoli there must be a wave of pressure change passing from the bronchi out to the periphery of the lung.

It is presumably this sudden release of pressure which is responsible for the sensation usually experienced that the lung is being distended but it is obvious from the mechanism of coughing that the lung is compressed and not in any sense distended. This building up of pressure followed by its sudden release must mean increased stress and strain on all the structures involved especially on the alveolar walls which are not robust and which carry this wave of pressure change. A chronic bronchitic patient may cough scores of times in a day and it seems reasonable to suppose that this type of stress and strain or wear and tear repeated over many years could in time produce the loss of elasticity and degenerative changes observed in the alveolar walls. In many respects there is an analogy in arterial hypertension in which without any over stretching of the blood vessels there may be loss of elasticity and other degenerative changes and these are ascribed by recent investigators to the chronic stress of raised pressure. It is also possible that wear and tear might occur outside the lungs when cough is chronic. The pleura might suffer for the same reason as do the alveoli and the thoracic cage is also strained. One has only to place a hand over the costal cartilages during coughing to realize that the impact is considerable. In asthma there may be the same degree of change in intra alveolar pressure as in coughing and although the change is not so sudden it is repeated more often (von Neergaard and Wirz 1927; Hartwich 1930). This stress and strain repeated many thousands of times a day could have the same effect as chronic cough—loss of pulmonary elasticity. The mechanism by which this loss may lead to many of the signs and symptoms of



## CHRONIC HYPERTROPHIC OR OBSTRUCTIVE EMPHYSEMA

is loss of elasticity due to the stress and strain of coughing or asthma and a third is the lowered resistance of the lung to wear and tear which varies from person to person and which diminishes with advancing age

### Treatment of chronic hypertrophic emphysema

The treatment of emphysema is essentially symptomatic for elastic tissue cannot regenerate and nothing can restore the structure of the lungs. Until heart failure supervenes the only symptom other than cough is dyspnoea on exertion. Therefore I first emphasize the value of broncho dilator drugs

**Broncho dilator drugs**—Although there may be no evidence of bronchospasm or resistance to respiration the administration of ephedrine or of other broncho dilator drugs not infrequently relieves the dyspnoea of emphysema. A possible explanation of this effect is that the bronchioles leading to the over distended air sacs and bullae are less capable of changes in calibre than are those leading to healthier parts of the lung. bronchospasm although not clinically manifest, might in this case increase the proportion of the inspired air deflected to these useless parts of the lung and the relief of bronchospasm by ephedrine medication would improve the efficiency of ventilation and thus relieve dyspnoea. The broncho dilator drugs which may be used and their method of administration are similar to those recommended in the more chronic phases of bronchial asthma (see p 92)

**Treatment of bronchitis**—Bronchitis should be treated in the usual way (see Volume I p 58) but the results are usually disappointing except when contributory factors such as dusty occupations or smoking can be prevented. The response to antibiotics is variable but is seldom impressive

**Pulmonary deflation**—Several procedures the purpose of which is to deflate the lung have been described and it is said that these increase the efficiency of respiration. Pneumothorax is one and relief of dyspnoea by this procedure both in human beings and in horses suffering from this disease has been claimed. Pneumothorax in emphysema however is a hazardous procedure. More popular in Great Britain and less dangerous are respiratory exercises designed to promote relaxation and to teach the patient to deflate the lung and to increase the use of the diaphragm. A well fitting abdominal belt will also raise the diaphragm by increasing intra abdominal pressure and more recently pneumoperitoneum has been suggested for the same purpose. All these procedures have been recommended and various theories have been put forward to explain the beneficial effects observed. Two effects are common to most of them. First the diaphragm is raised so that its convexity is increased and its efficiency therefore enhanced and it was for this reason that I have suggested the use of the abdominal belt. Secondly the lung is deflated by these procedures so that it contains less air. The decrease in volume is however very small—so small that it can hardly be measured—and the beneficial effects are probably for the most part due to increased efficiency of the respiratory musculature. It is also possible that with greater collapse of the superficial bullae efficiency of ventilation may be increased

**Treatment of heart failure**—When heart failure supervenes the general principles recommended in the treatment of cor pulmonale should be adopted. Oxygen should be given because the added insult of anoxia to a heart that is failing

greatly lessens the chances of recovery but it should be given with care for the sudden relief of anoxia may lead to convulsions and coma. It is believed by some physicians that digitalis should be avoided particularly in an emergency. Venesection has of course been recommended but it is now seldom used. Mercurial diuretics may however be of considerable value. Heart failure is often precipitated by respiratory infection commonly subacute broncho pneumonia and this usually responds to treatment with antibiotics. It is because broncho pneumonia is no longer a lethal complication that recovery from heart failure in emphysema is now a common occurrence.

*Conclusions*—The treatment of emphysema is thus far from being hopeless although it is essentially symptomatic. Bronchitis should be treated and in all cases broncho dilator drugs should be given a trial. Respiratory exercises or an abdominal belt or both may increase considerably the tolerance to exercise.

*Senile emphysema*—The treatment of emphysema is thus far from being hopeless although it is essentially symptomatic. Bronchitis should be treated and in all cases broncho dilator drugs should be given a trial. Respiratory exercises or an abdominal belt or both may increase considerably the tolerance to exercise.

## SENILE EMPHYSEMA

*Synonyms* Atrophic emphysema postural emphysema arteriosclerotic emphysema non obstructive emphysema small lunged emphysema senile atrophy of the lungs

### Clinical picture

As its name implies this is essentially a disease of old age (see Fig 41). The patient often presents a shrivelled and withered appearance. The chest is usually barrel shaped but there is no significant enlargement of the thoracic cage. The bony changes may be more pronounced than in the obstructive type of emphysema.

and in the later stages the



FIG 41—A case of senile emphysema. The wizened appearance, the upper thoracic kyphosis, the bulging of the sternum and the relative increase in the antero-posterior diameter are characteristic. There is no increase in the volume of the thoracic cage.

The appearance of the lung is variable. In some cases the process affects the thoracic cage; in others there is chronic bronchitis and this complicates the picture by introducing the obstructive element. Occasionally this atrophic process is believed to affect the lung itself but this has not been established. On the available evidence no clear description of the lung in senile emphysema can be given. It may be normal or it may show the characteristics found in the obstructive

## LOCALIZED OR FOCAL EMPHYSEMA

type of emphysema, but the lungs are usually small and bullae are seldom conspicuous

The symptoms and signs of senile emphysema bear little relationship to the apparent deformity. Symptoms are often absent, even in the presence of extreme kyphosis. More rarely there may be all the symptoms and signs characteristic of chronic obstructive emphysema.

### Aetiology

This condition is probably but one of the atrophic changes occurring in old age. Changes in the thoracic cage may be ascribed to atrophy of the intervertebral discs, leading to kyphosis, with consequent rotation of the ribs, so that the chest tends to become barrel shaped. Atrophy of the alveolar walls is said to lead to their disintegration but there is often an associated chronic bronchitis which exaggerates the emphysematous changes.

### Diagnosis

*Differentiation from obstructive emphysema*—Senile emphysema differs from the chronic obstructive form in the following respects. It is a disease of old age rather than of middle age. The cough is seldom severe and dyspnoea and cyanosis are usually absent. The kyphosis of the upper thoracic vertebrae is characteristic. Although barrel shaped, the chest is not enlarged. Chest expansion may be impaired, but usually inspiration has not the same heaving quality as in the obstructive type.

### Treatment

Treatment, if any is indicated, is similar to that of chronic obstructive emphysema.

## ACUTE VESICULAR EMPHYSEMA

This is an ill defined entity, it includes several conditions in which transient signs of emphysema appear.

For instance, during an acute attack of bronchial asthma the increased expiratory resistance may lead to distension of the lung, which subsides when the attack is over. Similarly, in acute respiratory infections, diphtheria, whooping cough or any condition associated with transient respiratory obstruction, the lungs may become large. This temporary distension does not rupture the alveolar walls nor, in fact, does it lead to any permanent damage but, as stated above, it may be a contributory factor to the production of obstructive emphysema.

Even less well defined are the changes sometimes observed after congestion of the lungs. Hyper resonance of the lung has been described following acute heart failure, strenuous exercise or septicaemia, it has also been found in the contra lateral lung in pneumothorax and massive pleural effusion, and after residence at high altitudes. To interpret this change as being indicative of emphysema does not appear to be justified.

## LOCALIZED OR FOCAL EMPHYSEMA

**Synonyms** Compensatory emphysema, complementary emphysema, vicarious emphysema.

## Pathogenesis

fails to expand on inspiration, either the surrounding alveoli must become distended to occupy the space, or the chest wall must be drawn in. Both phenomena usually occur if the contraction is sufficient to displace the mediastinum, and signs of emphysema may appear in the contralateral healthy lung.

*Focal emphysema in pneumoconiosis* —In most cases this form of emphysema is a pathological finding rather than a clinical entity, but an important exception is in pneumoconiosis. Recent investigations have shown that the emphysema which occurs in silicosis differs from obstructive or hypertrophic emphysema in its focal distribution. Around the pneumoconiotic nodules the air spaces may be dilated to a diameter of several millimetres, and these foci of emphysema are widely scattered throughout the lung, particularly in the secondary lobules (Gough, 1940 and 1947). The clinical picture is, however, much the same as in chronic obstructive or hypertrophic emphysema, with cough, dyspnoea and rarely cyanosis, and ultimately evidence of cor pulmonale. The aetiology of this type of focal emphysema remains obscure. The relative importance of cough and of contraction due to fibrosis is unknown; it is possible that the condition is one of ordinary obstructive emphysema, but the morbid appearances are modified by fibrosis and scarring, and by the widespread pleural adhesions which are common.

*Lobar obstructive emphysema* —Another type of localized emphysema, often lobar in distribution, may be a sequel to partial obstruction of a bronchus in conditions such as carcinoma of the lung and tuberculous bronchitis. If there is sufficient obstruction to the flow of air through the bronchus the elastic recoil of the lung may be inadequate to deflate the alveoli beyond the obstruction and inflation of this part of the lung will result. This condition has been called lobar obstructive emphysema, and if the obstruction is of long standing, it is claimed that gross over distension with bulla formation may result. Sometimes the obstruction leads to a "ball valve" occlusion of the bronchus which allows the entry of air, but not the exit, and under these circumstances inflation of the lobe may be considerable.

## ACUTE INTERSTITIAL EMPHYSEMA

Interstitial emphysema, or passage of air into the connective-tissue spaces, is usually a sequel to injuries of the chest or lung; it may appear after tracheotomy. It is a rare complication of severe asthma, when the cause is presumably rupture of the lung itself. The air may escape into the interlobular and sub-pleural spaces, or it may be extravasated along the walls of the bronchi and blood vessels to the mediastinum, passing up to the subcutaneous tissues of the neck and face.

There are not usually any symptoms, and the condition is first suspected when the characteristic crepitus is discovered in the subcutaneous tissues of the neck. Sometimes there are cyanosis and alarming dyspnoea, and on auscultation peculiar

crunching and crackling sounds may be heard over the heart Recovery is the rule and very rarely, if ever, is operative intervention indicated to relieve air which is under tension (Rosenberg and Rosenberg, 1938, Wallace, 1946, d Abreu, 1948)

## BIBLIOGRAPHY

- D Abreu A C (1948) Surgical Emphysema\* in *British Surgical Practice* Vol 3 p 382 London Butterworth
- Anthony A (1936) Zur Diagnostik der Lungenerweiterung *Verh Dtsch Ges inn Med*, 48 244
- Bates D V and Christie R V (1950) Intrapulmonary Mixing of Helium in Health and in Emphysema *Clin Sci* 9, 17
- Beitzke H (1925) Zur Mechanik des Gaswechsels beim Lungenemphysem *Dtsch Arch klin Med* 146 91
- Briscoe W A Becklake M R and Rose T F (1951) Intrapulmonary Mixing of Helium in Normal and Emphysematous Subjects *Clin Sci* 10, 37
- Cabot R (1927) *Physical Diagnosis* Baltimore Williams and Wilkins
- Christie R V (1934) The Elastic Properties of the Emphysematous Lung and their Clinical Significance *J clin Invest* 13, 295
- (1936) Pulmonary Congest on Following Artificial Pneumothorax Its Clinical Significance *Quart J Med* 5, 327
- (1939) Chronic Hypertrophic Emphysema Its Aetiology and the Cause of some of its Signs and Symptoms *Ednb med J* 46 463
- (1944) Emphysema of the Lungs *Brit med J* 1 105 and 143
- (1951) Emphysema of the Lungs in *British Encyclopaedia of Medical Practice* 2nd ed Vol 5 p 51 London Butterworth
- and McIntoch C A (1934) The Measurement of the Intrapleural Pressure in Man and its Significance *J clin Invest* 13 291
- Cournand A Baldwin E D Darling R C and Richards D W (1941) Studies on Intrapulmonary Mixture of Gases IV The Significance of the Pulmonary Emptying Rate and a Simplified Open Circuit Measurement of Residual Air *J clin Invest* 20 681
- and Berry F B (1942) Effect of Pneumectomy upon Cardopulmonary Function in Adult Patients *Ann Surg* 116, 532
- Lassen H C A and Richards D W (1937) Distribution of Respiratory Gases in a Closed Breathing Circuit II Pulmonary Fibrosis and Emphysema *J clin Invest* 16 9
- Davidson M (1936) Discussion on Emphysema *Proc R Soc Med* 29 1315
- Donald K W and Christie R V (1949) The Respiratory Response to Carbon Dioxide and Anoxia in Emphysema *Clin Sci* 8, 33
- Gough J (1940) Pneumoconiosis in Coal Trimmers *J Path Bact* 51 277
- Gough J (1947) Pneumoconiosis in Coal Workers in Wales *Occup Med* 4 86
- Hartwich A (1930) Pneumotachographische Untersuchungen über die Atemverhältnisse bei Hyper und Dyspnoischen *Z ges exp Med* 69, 482
- Hekam J B and Cargill W H (1947) Pulmonary Arterial Pressure in Congestive Failure and Emphysema *Amer J Med* 3, 504
- Hoover C F and Gammon J E (1915) The Dead Space in Moderate and Large Respiratory Ventilation *Arch intern Med* 15, 501
- Hurtado A Fray W W and McCann W S (1933) Studies of total Pulmonary Capacity and its Subdivisions IV Preliminary Observations on Cases of Pulmonary Emphysema and of Pneumoconiosis *J clin Invest* 12 833
- Knott J M S and Christie R V (1951) Radiological Diagnosis of Emphysema *Lancet* 1 881
- Kountz W B and Alexander H L (1934) Emphysema *Medicine* 13, 251
- Loeschke H (1928) Das Lungenemphysem *Handb spez path Anat Histol* Berlin 3 612
- Menely G R and Kaltreder N L (1949) The Volume of the Lung Determined by Helium Dilution Description of the Method and Comparison with other Procedures *J clin Invest* 28 129





## CHAPTER 10

### PULMONARY OEDEMA

G W HAYWARD and RONALD V CHRISTIE

#### PHYSIOLOGICAL FACTORS

THE PHYSIOLOGICAL factors responsible for the formation of oedema fluid in the lung are fundamentally the same as those concerned in the production of oedema elsewhere in the body, modified in some respects owing to differences in the pulmonary and systemic circulation. The factors to be considered are (1) hydrostatic pressure relationships, (2) osmotic factors, and (3) permeability of the capillaries.

##### Hydrostatic pressure relationships

Direct measurement of the pulmonary arterial pressure in the unanaesthetized human by means of cardiac catheterization is now possible, and it has been found that the systolic pressure in the pulmonary artery is between 15-30 millimetres of mercury (Cournand, 1947). Direct measurement of the pulmonary pressure in a normal man is not possible but catheterization of the pulmonary veins in patients with an atrial septal defect has given pressure readings of 5-7 millimetres of mercury (Cournand and his colleagues, 1947). The pressure at the arterial end of the pulmonary capillary is about 10 millimetres of mercury, as compared with a pressure of 25-30 millimetres in a systemic capillary. The pressure in the interstitial tissues of the lung is approximately that of the atmosphere so that the effective filtration pressure tending to cause transudation of fluid through the capillary membrane is also about 10 millimetres.

##### Osmotic factors

The osmotic pressure of the blood in the pulmonary vessels is the same as that elsewhere in the body, that is 25-30 millimetres of mercury with the normal plasma protein level of 7 grammes per cent. This is the force which tends to prevent leakage of fluid through the capillary wall.

##### Permeability of the capillary membrane

It is customary to imagine that the capillary wall acts as a semi permeable membrane, allowing crystalloids to pass readily in either direction, but retaining protein in the vessel. Studies of the lymphatic flow in animals show that usually a small amount of protein and fluid escapes from the vessel into the interstitial tissue of the lung and is re- absorbed by the lymphatics. In conditions in which the permeability of the pulmonary capillary is increased, plasma may escape in large amounts and the clinical condition of pulmonary oedema results. The permeability of these capillaries may be affected by a variety of factors. Some humoral, such as anoxia, others neurogenic, such as trauma to the heart or brain, and others toxic.

In animals (L. J. ...)

pulmonary capillaries would be accompanied by increased permeability and the formation of oedema fluid

### *Lymphatic drainage*

The lung is freely supplied with lymphatics which in health are easily able to remove the small quantities of plasma which escape from the capillaries. Changes in lymphatic flow when the permeability of the lung capillaries is increased have been studied by Drinker (1945). As a result of anoxia, or the administration of phosgene or thiourea compounds, the lymphatic flow is considerably increased. As more fluid escapes from the vessel, the lymphatics may be unable to remove all of the fluid, which then escapes into the alveoli causing pulmonary oedema. It is probable that a similar sequence takes place in man.

From these physiological considerations it may be seen that in the normal person the relatively low filtration pressure and high osmotic pressure will reduce the transudation of fluid to a minimum, and any fluid which may escape is efficiently removed by the lymphatics. If, however, the permeability of the capillary wall is increased, either through the influence of anoxia, toxic substances, or nervous stimulation, or if in addition the pulmonary capillary pressure is increased by either an increase in the pulmonary arterial pressure or pulmonary venous pressure, fluid may pass into the interstitial tissue and alveoli of the lung more quickly than it can be removed by the lymphatics, and the condition of pulmonary oedema will result.

## CLASSIFICATION

Pulmonary oedema occurs in such a wide range of clinical conditions that an aetiological classification is unsatisfactory. A clinical classification, based primarily on the type of pulmonary oedema is probably to be preferred.

- (1) Acute pulmonary oedema
  - (a) Associated with heart failure (left ventricular failure, coronary thrombosis, acute cor pulmonale)
  - (b) Associated with damage to or diseases of, the central nervous system (head trauma, cerebral vascular accidents, space occupying lesions of the brain)
  - (c) Following lung irritants such as phosgene
- (2) Subacute
  - (a) Associated with left ventricular strain
    - (i) rheumatic pneumonia
    - (ii) uraemia
    - (iii) polyarteritis nodosa
  - (b) Disturbances of fluid balance
    - (i) desoxycorticosterone acetate poisoning
    - (ii) transfusion
    - (iii) stilboestrol

## ACUTE PULMONARY OEDEMA

- (c) Associated with lung irritants  
Infections (pneumonia, virus infections, industrial irritants)
- (3) Chronic
  - (a) Associated with heart failure—hypostatic congestion
  - (b) Associated with lung irritants—chronic infections

It would be redundant to describe each of these types in detail since much of what applies to one, applies to the others. The attention of clinicians and experimentalists has been mainly focused on acute pulmonary oedema associated with heart failure, and for this reason discussion of the fundamental principles involved will be for the most part confined to this section.

## ACUTE PULMONARY OEDEMA

**Associated with heart failure**  
The syndrome of acute pulmonary oedema associated with heart failure has also been called cardiac asthma, or paroxysmal nocturnal cardiac dyspnoea. These are all manifestations of acute pulmonary congestion any of which may develop into acute pulmonary oedema. In the cases labelled "cardiac asthma" there is in addition a varying degree of bronchospasm, and the term paroxysmal nocturnal cardiac dyspnoea merely indicates two of the well recognized clinical features of the attack. It would be better to adopt the term acute pulmonary congestion, with or without pulmonary oedema, for all cases.

The first clinical description of this condition was made about two centuries ago, and clinicians from the time of Laennec have favoured the simple hypothesis that the cause is acute failure of the left ventricle to expel the blood delivered to it by the right ventricle with consequent engorgement of the pulmonary vascular bed. This purely mechanical concept of the pathogenesis of acute pulmonary oedema has, however, little to support it either clinically or experimentally.

The basis of the mechanical theory of blood accumulating behind a failing left ventricle lies in the experiments of Welch in 1878 on rabbits. He interfered with the function of the left ventricle by crushing it or by ligation of the aorta and produced pulmonary oedema. His results are still quoted, but have been criticized on the grounds that the appearance of pulmonary oedema was inconstant, spontaneous pulmonary oedema frequently occurs in rabbits, the nature of the experiments resulted in gross damage to the animals, and finally that it has been impossible to reproduce the results in other animals. Numerous experiments designed to change mechanically the circulation in the lungs by constriction of the pulmonary veins have shown that it is difficult to produce pulmonary oedema in this way. Experiments with the heart lung preparation have been quoted in support of the theory of left ventricular failure, but it is now agreed that the oedema produced was usually due to chemical changes in the perfused blood. In place of the mechanical theory of the cause of pulmonary oedema, there has been a tendency among physiologists to state that pulmonary oedema is caused by neurogenic vasodilatation of the pulmonary capillaries and resulting increased permeability. The stimulus may arise from direct excitation of nervous centres, as in head injuries, or reflexly from stimuli arising in the heart or large vessels (Luisada, 1940).

The main conclusion that can be drawn from the mass of experimental evidence in the literature is that although pulmonary oedema can be produced by various mechanical and reflex procedures, the only reliable method of producing acute pulmonary oedema in animals is by means of substances which act directly on the pulmonary vascular bed, such as anoxia, phosgene and certain other chemicals, notably the thiourea derivatives. It should not be forgotten that it has so far proved impossible to produce chronic pulmonary congestion or in fact any condition in animals that resembles the clinical picture which is known to predispose to acute pulmonary oedema or which resembles the actual clinical attack of acute pulmonary oedema.

The vast majority of patients with acute pulmonary oedema give a history of chronic left ventricular failure with dyspnoea and orthopnoea, and in these the incidence of attacks is greatly reduced when right heart failure supervenes. This suggests that pulmonary congestion may at least play an important part, and this view is strengthened by the clinical observation that venesection or the application of limb tourniquets which reduce the degree of pulmonary congestion are useful procedures in the treatment of an acute attack.

The experimental and clinical evidence is not, however, wholly contradictory. Although acute pulmonary oedema occurs most commonly in patients with chronic left ventricular failure and pulmonary congestion it may also follow cerebral accidents or right ventricular strain such as occurs in pulmonary infarction, acute cor pulmonale, pneumonia and thrombosis of the coronary vessels supplying the right ventricle. In this group there is nothing to suggest left ventricular failure and yet frank pulmonary oedema may occur. It is possible that in these cases reflex vasodilatation and increased permeability of the pulmonary vessels is the cause, the stimulus arising from the heart, great vessels, or the brain.

The clinical observation that the attacks usually occur during sleep has been the source of much speculation. Increased autonomic activity when the patient is awakened by dreams or coughing may lead to pulmonary vasodilatation, or, in patients who are orthopnoeic, slipping down in bed may lead to increased pulmonary congestion. Another explanation which has some evidence to support it is that a sudden movement while asleep may cause a considerable increase in the venous return to the heart (Christie, 1938). Even in the healthy individual a transitory increase in the heart rate of 30-40 beats a minute may occur during sleep, and with the venous stagnation which occurs in heart failure it seems reasonable to suppose that, after a period of relaxation, a sudden movement during sleep might increase considerably the influx of blood to the heart.

The conclusion which can be drawn from this conflicting evidence is that we do not know the essential cause of acute pulmonary oedema, but that certain contributory factors appear to be of importance. A background of pulmonary congestion favours the onset, and the trigger mechanism which initiates the attack must be increased capillary permeability due either to reflex vasodilatation or to a sudden increase in the return of venous blood to the heart. The reflex stimulus may arise in the heart itself, from a myocardial infarct or sudden pressure changes in any of the heart cavities or great vessels, it may arise from the lungs,

as in pulmonary infarction or lung irritants or from the higher centres from emotion excitement or trauma

## ACUTE PULMONARY OEDEMA

### *Aetiology*

In heart failure acute pulmonary oedema is most commonly associated with hypertension aortic valvular disease particularly aortic regurgitation and coronary artery disease. It may also occur as mentioned above in right heart failure such as occurs in pulmonary or myocardial infarction. In all of these pulmonary congestion is a prominent feature. Attacks of acute pulmonary oedema may occur in patients with tight mitral stenosis. In these patients who often give a history of recurrent haemoptyses there is little enlargement of the heart or left auricle the pulmonary arteries are prominent and the pulmonary artery pressure is raised. Acute pulmonary oedema is rarely seen with auricular fibrillation.

### *Pathology*

The pathological changes in the lung are those of widespread oedema both interstitial intra alveolar and filling the larger air passages so that on section the cut surface exudes frothy fluid. The lymphatics and pulmonary capillaries are widely dilated. As seen at autopsy the whole of both lungs are affected. The changes in the heart are those of the underlying disease.

### *Clinical course*

Acute pulmonary oedema associated with heart failure usually occurs in patients giving a history of increasing cardiac dyspnoea and orthopnoea. With little or no warning and usually at night he suddenly develops intense and terrifying dyspnoea. He sits up or tries to get out of bed struggling for breath. Coughing at the onset is frequent and aggravates the dyspnoea and the onset of bronchospasm may further embarrass his breathing. Substernal tightness is often experienced together with a feeling of suffocation. The attack may stop almost as suddenly as it started or it may increase in severity to a condition of gross pulmonary oedema. The cough becomes more frequent and frothy or watery sputum sometimes blood stained may be brought up.

When seen during the attack the patient appears in obvious respiratory distress. Cyanosis appears early and if the attack continues will become marked. Examination of the lungs in the early stage of the attack may show little or no abnormality. Some rales are usually heard first only in the mid zone of both lungs and later extending and becoming coarser until the breath sounds may be obscured by the added sounds. The heart rhythm is nearly always normal sinus rhythm and as Bedford (1939) has pointed out it is unusual to see attacks of acute pulmonary oedema in patients with established auricular fibrillation. In the latter condition both ventricles are commonly failing and it is well recognized clinically that with the onset of right heart failure attacks of acute pulmonary oedema may cease. There is evidence of left ventricular hypertrophy in most cases the notable exceptions being those patients who have coronary heart disease and myocardial infarction without an associated hypertension. The pulmonary second sound is accentuated and there is frequently

a presystolic triple rhythm—'gallop rhythm'—at the apex. Except in those cases due to myocardial infarction, the blood pressure during an attack is raised or unchanged until the terminal stage of an attack, when it may fall. A significant degree of *pulsus alternans* may often be detected when the blood pressure is being measured.




FIG. 42.—Simultaneous recordings of intrapleural pressure and tidal air in a patient with heart failure showing the increased rigidity of the lung caused by pulmonary congestion. The pleural pressure fluctuation is 27 centimetres of water (10 to -17). Normal fluctuation 2.7 centimetres of water. (From Christie and Meakins, 1934.)

Intense dyspnoea dominates the picture. This is not due to the stimulus of carbon dioxide retention or of anoxia, but is a direct result of pulmonary congestion (Christie, 1938). The lungs become more rigid and less distensible (Fig. 42) and reflex dyspnoea follows through stimulation either of the Hering

## ACUTE PULMONARY OEDEMA

Breuer reflex or of vagal nerve endings in the pulmonary vascular bed (Bulbring and Whitteridge, 1945)

Cyanosis, since it can be relieved by oxygen, must be for the most part pulmonary in origin and is due to the exudate in the alveoli which impedes gaseous exchange. The elimination of carbon dioxide is not significantly impaired, partly because its dissociation curve differs from that of oxygen and partly because it is more soluble and can diffuse more readily through the alveolar exudate.

Bronchospasm is variable in its intensity and is usually ascribed to congestion of the bronchial mucosa through the anastomotic channels which connect the bronchial and pulmonary vessels. This can hardly be called an explanation since the bronchospasm may be relieved by adrenaline and is presumably autonomic in origin.

Why the attack passes off is even more uncertain than why it should start. It may be that a rapidly failing left ventricle and increasing anoxia in the tissues leads to generalized vasodilatation and increasing anoxia in the tissues pulmonary to the systemic vascular bed, with consequent relief of congestion. That this does indeed happen, that the termination of these changes is often imperceptible, that the pulse is often imperceptible, caused by marked left ventricular hypertrophy.

The electrocardiogram, before, during an attack, will often show the changes caused by marked left ventricular hypertrophy. The limb leads, show left axis deviation with flattening or inversion of  $T_1$ . The precordial leads,  $V_1$  and  $V_2$  (over right ventricle) show a small R and deep S, whereas in  $V_3$ - $V_6$  the R wave is tall and the T wave inverted.

### Radiological findings

Radiological changes in the lungs accompany the appearance of clinical signs of oedema in the lungs. The typical early appearance is that of symmetrical areas of increased density spreading out from both hila to give a butterfly wing appearance (Fig 43). At this stage, the periphery of the lung fields and particularly the lung bases may appear unduly clear. Careful examination of the areas of increased density will show a diffuse haziness, obliterating the lung markings, unlike the changes in chronic congestion where the lung markings may be increased and the dilated hilar vessels clearly seen. If the attack is short lived these changes disappear quickly, and all radiological signs may have disappeared in 4-6 hours. If the attack progresses to massive pulmonary oedema, the areas of haziness will spread to involve all of both lungs (Roesler, 1943).

### Other investigations

No accurate measurements of cardiac output or pulmonary arterial pressure during an attack are available. The circulation time is prolonged and the vital capacity diminished. A marked degree of arterial anaemia will be found during the attack, and cyanosis may be exaggerated by the haemoconcentration caused by the loss of a large amount of protein-containing fluid from the circulation.

### Treatment

An attack of acute pulmonary oedema is a medical emergency, and prompt





FIG 43—(a) Skiagram during an attack of acute pulmonary oedema in a patient with hypertensive heart disease (b) sk agram 12 hours later showing considerable resolution

(a)



(b)

treatment in the early stages may abort the attack, whereas if delayed until the later stages, treatment may be difficult

*Oxygen*—The relief of anoxia is important as the pulmonary capillaries themselves may under the influence of anoxia become more permeable with resultant increasing pulmonary oedema and more anoxia. Prompt administration of oxygen will help to break this vicious circle. It should be given either by B L B mask, or if this is not tolerated, by spectacle frame nasal catheters. A flow of 5–8 litres per minute is usually recommended, even if the more efficient B L B mask is used. Because of restlessness and the feeling of suffocation, an oxygen tent is rarely tolerated.

*Sedatives*—Sedatives are necessary in this condition to relieve the patient's distress during the attacks. They may also be helpful because if, as is possible, a nervous reflex mechanism is responsible for initiating the attack, the sedative by depressing reflex action may help to stop it. The drug of choice is morphine which should be given subcutaneously in doses of  $\frac{1}{4}$ – $\frac{1}{2}$  grain, or intravenously in those patients with circulatory collapse. Alternatively, the barbiturates may be used, particularly the rapidly acting ones.

*Procedures acting probably by lowering the venous pressure*—Rapid venesection of 600 millilitres of blood in those patients with a considerable increase in venous pressure will often help to terminate the attack. The use of tourniquets around the thighs, to obstruct venous return, has also been advised for the same purpose. Apart from morphine probably the most useful drug in the treatment of an attack is theophylline with ethylenediamine. In addition to causing a fall in venous and right auricular pressure (Howarth, McMichael and Sharpey Schafer, 1947), it will also relieve bronchospasm. It should be given intravenously in a dose of 0.24–0.48 grammes diluted in 10 millilitres of distilled water. The value of atropine is disputed. It may help to dry up an excess of bronchial gland secretion, but its effect on the pulmonary nerve endings in man is unknown. The place of digitalis in the treatment of acute pulmonary oedema is controversial. It is usually given in a dosage to control the ventricular rate.

*Prevention of attacks*—Any procedure which reduces pulmonary congestion

### *Prognosis*

The occurrence of attacks of acute pulmonary oedema in a patient with heart disease is a sign of serious prognostic importance. The average duration of life after the first attack in a series of patients with left ventricular failure was only 2 years (Bedford, 1939).

### *Pulmonary oedema associated with lesions of the central nervous system*

Severe injuries to the skull or brain not infrequently cause acute pulmonary oedema, and Weisman (1939) found that 70 per cent of patients dying after a fractured skull had signs of oedema of the lungs. It occurs less frequently in

diseases of the central nervous system such as cerebral haemorrhage, and occasionally is associated with brain abscess or tumour

This type of pulmonary oedema which occurs in patients with no disease of the heart, or history of chronic pulmonary congestion, is of particular interest to the experimental physiologist, as it resembles closely the type of acute pulmonary oedema produced in animals by a wide range of experimental procedures. The cause is presumably increased permeability of the pulmonary capillaries, without the added factors of left ventricular failure, and therefore without massive pulmonary congestion. The cause of the increased capillary permeability which follows damage to the brain is unknown, but it is presumably due to reflex dilatation of the pulmonary capillaries and consequent increase in permeability.

The clinical picture of a patient with this type of acute pulmonary oedema differs from that seen in left ventricular failure. Dyspnoea does not dominate the picture—probably because of the absence of gross pulmonary congestion. The respirations are often deep, laboured, and stertorous. The frequent absence of orthopnoea is difficult to evaluate because of the disordered consciousness of the patient. More prominent in this traumatic type of oedema are moist sounds in the lung, with cyanosis, and cough with thin expectoration which may later become frothy. It is usually a terminal condition, and fails to respond to the methods of treatment which have been outlined in the previous section.

### SUBACUTE PULMONARY OEDEMA

#### Associated with severe hypertension

An unusual type of pulmonary oedema which is subacute in its clinical course is seen in several diseases, all of which are associated with left ventricular strain.

The changes in the lungs in rheumatic pneumonia as described by Hadfield (1938) consist of an initial fibrinous alveolitis with later infiltration of the exudate with mononuclear cells and eventual organization when fibroblasts may be seen. The alveolar ducts are lined by a hyaline membrane. The lungs macroscopically are solid or rubbery, particularly in the central position near the hilum. Other changes due to the rheumatic infection are seen, and in all cases carditis is present, often with a fibrinous pericarditis. These changes in the lung were thought by many to represent the specific pulmonary lesion of the rheumatic infection.

The clinical picture of a patient with the 'rheumatic lung' is initially that of a severe rheumatic carditis, with later increasing dyspnoea and often slight cough. The physical signs in the lungs are inconspicuous compared with the extent of the lesions seen radiologically. Slight impairment of percussion note and breath sounds in the mid zone with a few fine crepitations may be detected but not infrequently the lungs show no abnormality on physical examination. The chest skiagram shows a characteristic change, with central cloudy opacities, often confluent, extending peripherally into the lung with the lung bases and apices remaining clear (Fig. 44).

An exactly similar condition is seen in uraemia (Doniach, 1947), in acute nephritis and in polyarteritis nodosa, when there is severe hypertension. It is with those seen

# SUBACUTE PULMONARY OEDEMA

FIG 44 — Sub-acute pulmonary oedema in rheumatic fever (autopsy control)



FIG 45 — Sub-acute pulmonary oedema in malignant hypertension (autopsy control)



It is probable that in all of these diseases the same pathological process is taking place, and that the clinical, radiological and pathological findings are caused by increased capillary permeability, usually in association with left ventricular failure with resulting pulmonary oedema, both interstitial and intra alveolar (Epstein and Greenspan, 1941). The difference between this subacute pulmonary oedema and the acute type described above, presumably lies in the fact that in the former the exudate clots and then becomes organized, with cellular infiltration. In acute pulmonary oedema the exudate is removed before clotting and organization has time to occur. It is of interest that the same histological appearance in the lungs can be produced in animals by various procedures which cause pulmonary oedema, such as the administrations of large doses of thiourea derivatives and in sulphonamide hypersensitivity.

### Subacute pulmonary oedema caused by a disturbance of fluid balance

#### *Intravenous therapy*

In patients with a normal cardiovascular system it is difficult to produce circulatory overloading, even when comparatively large amounts of fluid or blood are given. The increased venous return to the right heart increases the output of the heart, and the additional fluid is accommodated by the opening up of reserve capillaries in the lungs and elsewhere in the body. In patients who are ill, particularly those with anaemia associated with chronic infection, quite small amounts of blood or saline solution given intravenously may cause circulatory overloading and a subacute type of pulmonary oedema may be produced. In such a patient, the earliest signs will be a engorged neck veins, a slight rise in respiratory rate, and a slight dilatation of the lungs at auscultation. If the transfusion is continued, moist sounds will appear at the lung bases and frank pulmonary oedema may result. The x ray findings are increased prominence of the vascular markings at the lung roots with generalized loss of translucency of the lung fields most marked at the lung bases.

° - ' - - - - - ary oedema is presumably

The myocardium in the  
mplication of intravenous

therapy shows the fatty change due to anaemia and toxæmia, and failure occurs when the heart is unable to compensate for the increased venous return and increased circulatory blood volume.

The treatment of this condition is both prophylactic and curative. If the danger of producing pulmonary oedema in patients with chronic anaemia and sepsis is recognized, transfusion should be given slowly, and the use of packed cells instead of whole blood to reduce the bulk of fluid given is advised. If pulmonary oedema does occur during a transfusion, the transfusion should be stopped, possibly temporarily and if gross oedema is present, the immediate administration of a mercurial diuretic, such as mersalyl or Neptal is advisable. The marked diuresis which follows will help to remove the excess of fluid from the body and the pulmonary oedema may disappear.

*Administration of desoxycorticosterone acetate (DOCA) and stilboestrol*

The synthetic suprarenal cortical hormone DOCA differs from the naturally occurring hormone in that a more marked retention of sodium and water occurs during the period of administration of the drug. If given to normal healthy persons, a gain in weight due to the increase in extracellular fluid occurs (Thorn and Emerson, 1940). The same applies to patients with Addison's disease who are given excessive doses of DOCA, and when the sodium intake in the diet is not restricted, pulmonary oedema may be produced. This type of oedema is not associated with marked pulmonary congestion, so that although the respiratory rate may rise, dyspnoea is often not marked. Râles may be heard over both lungs, and the chest skiagram shows the typical appearance of diffuse pulmonary oedema. The condition when recognized usually subsides rapidly when DOCA is discontinued.

Retention of sodium is also known to occur as a result of the administration of stilboestrol and pulmonary oedema has occasionally been noticed during treatment of the metastases of a prostatic carcinoma. The correction of anaemia by transfusion in patients under treatment with either of these drugs should be carried out with caution as this may precipitate an attack of pulmonary oedema.

**Subacute pulmonary oedema associated with lung irritants**

A subacute type of pulmonary oedema is seen in several infections of the lung, notably in the virus infections causing primary atypical pneumonia. The clinical features of this disease are described elsewhere. Radiologically, the soft patchy opacities often centrally placed, resemble the changes described above as typical of subacute pulmonary oedema, and in the few cases which have come to autopsy a similar histological appearance has been noted (Jenson, 1946, Golden, 1944).

Industrial irritants are described in Volume I, Chapter 6.

## CHRONIC PULMONARY OEDEMA

### Hypostatic congestion

In patients with chronic left ventricular failure, a degree of persistent oedema of the lung bases is frequently seen. A similar condition may occur in elderly patients who are confined to bed. Presumably as a result of impaired diaphragmatic movement, slowing of the circulation and anoxia there is increased permeability of the capillaries and oedema at the bases. Treatment is that of the underlying condition and early mobilization of the patient.

## REFERENCES

- Bedford D E (1939) Left Ventricular Failure. *Lancet* **I** 1303  
 Bulbring E and Whitteridge D (1945) The Activity of Vagal Efferent Endings during Congestion in Perfused Lungs. *J Physiol* **103**, 477  
 Christie R V (1938) Dyspnoea—a critical review. *Quart J Med* **7**, 421  
 — and Meakins J C (1934) The Intra pleural Pleural in Congestive Heart Failure and its Clinical Significance. *J clin Invest* **13**, 323

### PULMONARY OEDEMA

- Courmand A (1947) Recent Observations in the Dynamics of the Pulmonary Circulation  
*Bull N Y Acad Med* 23, 27
- Mosley H L Himmelstein A Dresdale D and Baldwin J (1947) Recording of Blood Pressure from Left Auricle and Pulmonary Veins in Human Subjects with Interauricular Septal Defect *Amer J Physiol* 150 267
- Daly I de B (1946) Pulmonary Oedema—physiological consideration "Thorax" 1 182
- Domach I (1947) Uremic Edema of Lung *Amer J Roentgenol* 58 620
- Drinker C K (1945) *Pulmonary Oedema and Inflammation* Cambridge Mass Harvard University Press
- Epstein E Z and Greenspan E B (1941) Rheumatic Pneumonia *Arch intern Med* 68 1047
- " " " " " 187
- Theo-
- Jensen C R (1946) Non suppurative Post streptococcic (Rheumatic) Pneumonitis *Arch intern Med* 77 237
- Luisada A (1940) Pulmonary Oedema *Medicine* 19 475
- Roesler H (1943) *Clinical Roentgenology of the Cardiovascular System* 2nd Ed Springfield Ill Thomas
- Thorn G W and Emerson K (1940) Role of Gonadal and Adrenal Cortical Hormones in Production of Edema *Ann intern Med* 14 757
- Weisman S J (1939) Edema and Congestion of Lungs Resulting from Intracranial Haemorrhage *Surgery* 6 722

## CHAPTER II

### ATELECTASIS

F P LEE LANDER

THE WORD "atelectasis" means "imperfect expansion" (*Oxford English Dictionary*). It was first used in the English literature in 1846, in Swaine's translation of Hasse's *Pathological Anatomy*. Shortly afterwards Rees (1850) wrote a short monograph entitled *Atelectasis Pulmonum* in which he gave the following definition

Imperfect expansion Atelectasis is intended to designate a condition of the lung in which there is simple occlusion or shutting up of the pulmonary air cells without any inflammatory or other morbid deposits

Since that time the term has been in current use, and though a few writers have regarded it as describing collapse of congenital origin only, the majority have not so limited its use. The morbid anatomists of the middle nineteenth century were fully aware of the frequent occurrence of the condition, and advanced many theories as to its formation

Legendre and Bailly (1844) identified broncho-pneumonia as a form of atelectasis, and produced some convincing experimental work to support their theory

Gardner (1851), in his treatise on bronchitis, mentioned the frequency with which the condition occurs, and stated that it was the commonest complication of any bronchitic condition. Furthermore he produced, for the first time, the theory of the mucus plug, and drew a diagram indicating the way in which the plug would be affected by changes of pressure

From the literature it appears that the attention of the morbid anatomists was then distracted from the condition, for in the latter part of the nineteenth century little reference was made to it, and it was left to William Pasteur (1908), in his Bradshaw Lecture, to re-discover it and to connect it with diphtheria and later with post-operative complications. Pasteur distinguished two varieties massive collapse due to diaphragmatic paralysis, which he called active collapse, and lobular collapse due to bronchial obstruction, which he called passive collapse

An answer to many of the questions posed by Pasteur, and a negation of most of his theory is offered by the extensive use since his time of the operation of diaphragmatic paralysis by phrenic nerve interruption. This diaphragmatic paralysis, we know from experience, is not necessarily or even commonly attended by massive collapse, and it seems probable that in the cases described by Pasteur diaphragmatic paralysis was attended by obstruction of the finer bronchi, although it is possible that in some of his cases elevation of the diaphragm and restricted movement secondary to the massive collapse were mistaken for actual paralysis of the diaphragm. Nevertheless, Pasteur's work remains a classic, largely because he described, for the first time, the detailed physical signs and attributed them rightly to lessened movement of the diaphragm



# PULMONARY OEDEMA

- Cournand A (1947) Recent Observations in the Dynamics of the Pulmonary Circulation  
*Bull N Y Acad Med* 23 27
- Motley H L Himmelstein, A Dresdale D and Baldwin J (1947) Recording of  
 Blood Pressure from Left Auricle and Pulmonary Veins in Human Subjects with Inter-  
 auricular Septal Defect *Amer J Physiol* 150 267
- Daly I de B (1946) Pulmonary Oedema—physiological consideration *Thorax* 1 182
- Donach I (1947) Uremic Edema of Lungs *Amer J Roentgenol* 58 620
- Drunker C K (1945) *Pulmonary Oedema and Inflammation* Cambridge Mass Harvard  
 University Press
- Epstein E Z and Greenspan E B (1941) Rheumatic Pneumonia *Arch intern Med*  
 68 1047
- Golden A (1944) Pathological Anatomy of Atypical Pneumonia *Arch Path* 38 187
- Hadfield G (1938) Rheumatic Lung *St Bari's Hosp Rep* 71, 17
- Howarth S McMichael J and Sharpey Schafer E P (1947) Circulatory Action of Theo-  
 phylline Ethylene Diamine *Clin Sci* 6 125
- Jensen C R (1946) Non suppurative Post streptococcic (Rheumatic) Pneumonitis *Arch*  
*intern Med* 77 237
- Luisada A (1940) Pulmonary Oedema *Medicine* 19, 475
- Roesler H (1943) *Clinical Roentgenology of the Cardiovascular System* 2nd Ed Springfield  
 Ill Thomas
- Thorn G W and Emerson K (1940) Role of Gonadal and Adrenal Cortical Hormones  
 in Production of Edema *Ann intern Med* 14 757
- Weisman S J (1939) Edema and Congestion of Lungs Resulting from Intracranial Haemor-  
 rhage *Surgery* 6 722

and no other compensatory mechanism may be called into play. As the child grows, and the thorax enlarges, then the collapse becomes manifest with the development of the other compensatory mechanisms.

Gardner, in 1850, drew three simple diagrams to explain his theory of the production of emphysema. In these diagrams, Fig 46*a*, showing three equal partitions represents three lobes or portions of a lung. Fig 46*b*, demonstrates that when one portion of the lung has collapsed the other two portions expand by drawing in air, to fill the whole space. Fig 46*c* shows that when two portions




FIG 46 — Diagrammatic representation of

space




FIG 47—(a) The same as in Fig 46(a) but including bronchi (b) expansion of bronchus when a lobe collapses

of the lung collapse the remaining portion expands to fill the whole space. In Gardner's original diagrams, he showed an equal expansion of the two remaining portions of the lung. As a result of study of radiographs and lobectomy specimens it is now known that the expansion is not evenly distributed through the thorax, but is more marked nearer the collapsed portion.

So far we have been considering the behaviour of the alveoli. When considering the behaviour of the bronchi exactly the same considerations apply. The bronchi are highly elastic tubes; the part of the bronchus distal to the obstruction (that is cut off from the outside air) will shrink because it is in unobstructed communication with the alveoli, and replenishment of air cannot take place. The part of the bronchus proximal to the obstruction being in free communication with the outside air will therefore expand (Fig 47).

The walls of the thoracic cage are also elastic and expansile and consequently will be affected by any diminution of the volume of any part of the thoracic contents. The area involved will be that which is closest to the shrinking volume of the lung. The mediastinum is readily mobile and displacement of the heart occurs in atelectasis involving the lower parts of the lung on either side, this

Today, whilst bronchial obstruction is regarded as the most important, if not the only, method of producing atelectasis, a great deal of confusion still exists as to the actual mechanism by which it is produced. Furthermore, the compensating mechanisms which are called into play by the shrinkage of the atelectatic area are only just beginning to be studied and understood.

### PHYSIOLOGY

When a bronchus is completely obstructed, alveolar air is absorbed and cannot be renewed. This absorption of air is attended by the shrinkage of the lung and this keeps the pressure of air in the obstructed alveoli constant. If there were no shrinkage, pressure of air would fall and absorption of air would cease when the alveolar air pressure was at the same level as the partial pressure of the gases in the venous blood. The shrinkage is rendered possible by the elasticity of the alveolar walls, which not only allows the obstructed lung to shrink, but also allows the unobstructed lung to expand and fill the space left by the retracting obstructed lung.

This space is filled also by contraction of the walls of the thorax, by a move of the mediastinal structures towards the retracting lung, and also by an elevation of the diaphragm on the affected side. All these factors can be seen and verified, clinically and radiologically. They do not all occur in each case to the same degree. In some cases the unobstructed lung expands to such an extent that no other mechanism is necessary to fill the space. A further effect of shrinkage is a compensation by the bronchi in the unobstructed lung, which dilate to beyond their normal range.

The size of the thorax varies with respiration, and consequently the degree of compensation required is greater in inspiration, when the thoracic volume is at its maximum, than in expiration. This can be watched on the fluoroscopic screen, and demonstrated by inspiratory and expiratory radiograms. Clinically, however, a diminution in the respiratory movements of the affected side can be seen, thus a limitation in the increase in thoracic volume on the affected side is one of the compensatory mechanisms.

Occasionally, in cases of complete bronchial obstruction, as air absorption occurs secretion is poured into the bronchi and thus the gaseous pressure in the obstruction system is maintained at a high level and this continues until all air is absorbed and the state then exists of an airless lung with complete bronchial obstruction and no shrinkage.

All states between the condition just described, through a range of reduced secretion and increased shrinkage, to a completely dry atelectasis, can occur.

If the main bronchial obstruction persists absorption of this secretion may occur, with gradual shrinkage of the obstructed portion of the lung.

In the case of children, a slightly different state exists because the child's thoracic volume is small as compared with the adult's thoracic volume, and the child's lung increases in volume enormously as growth occurs and adult stature is attained. The increase in size occurs largely by an increase in size of each individual alveolus. Consequently, when atelectasis occurs in a child's lung, compensation takes place simply by enlargement of the adjacent alveoli,

## CAUSATION

In certain cases a new growth of a main bronchus may not, by itself, produce complete obstruction. The bronchial lumen however, may be narrowed so much that the obstruction is made complete by thick, tenacious mucus. As a result, air is absorbed in the usual way behind the obstruction and the mucus is sucked down the bronchial tubes. If a mass of mucus is small, an airway may be re-established and the condition of the lung return to normal (Fig 48). However, if the mucus is of sufficient volume the obstruction may be complete (Fig 49).




FIG 48—(a) Small mass of mucus in bronchus, (b) airway re-established




FIG 49—(a) Large mass of mucus in bronchus (b) the obstruction is complete

This mucus plug will be sucked down the bronchus splitting up at each division until finally the original plug is split into a thousand fragments obstructing a thousand terminal bronchi. In this way lobar collapse may be produced by

complicating cases of tuberculous tracheo bronchitis

Obstruction of finer divisions of the bronchi leads to segmental atelectasis and obstruction of terminal bronchi leads to lobular atelectasis. The nature of the obstruction in the finer bronchi is not certain as direct observation is impossible. Spasm of the bronchial muscle, oedema of the bronchial mucous membrane, and fine plugs of mucus probably all play a part. These fine plugs of mucus have been demonstrated in autopsy specimens in post operative cases. Whooping-cough, asthma and bronchitis are commonly complicated by this degree of atelectasis, and tuberculous tracheo bronchitis is a less common cause

## SYMPTOMS

Massive atelectasis is usually attended by a dramatic onset. Sudden urgent dyspnoea, pain in the chest and palpitations are frequent. Occasionally cyanosis also occurs.

Lobar collapse is usually less obtrusive in its onset. A pain in the chest or tightness may be noticed. dyspnoea is sometimes present, and palpitations may occur.

## ATELECTASIS

displacement being usually accentuated in inspiration. The trachea is displaced by atelectasis of the upper lobe on either side, and the pulmonary arteries are also pulled up and distorted.

Atelectasis of one complete lung produces the maximal effect on the thoracic boundaries. It has been shown that the atelectatic lung lies in the paravertebral sulcus behind the hilum, the anterior part of the affected side being occupied by the heart and by the lung from the other side. The diaphragm is involved in all these cases of lobar collapse. In complete collapse of one lung, elevation of the diaphragm is extreme, and the diaphragm is immobile. In lower lobe collapse, elevation and flattening of the diaphragm occurs, with restriction of the diaphragmatic movement. In upper lobe collapse, elevation may not be so marked, but restriction of movement occurs. In lobular or segmental collapse the effects are more limited, and distortion of the mediastinum occurs as well as localized effects on the diaphragm.

## CAUSATION

Any obstruction of a bronchus, if complete both in the inspiratory and expiratory positions, is attended by atelectasis of the corresponding bronchi and alveoli. The obstruction may be situated anywhere between the epiglottis and the finest contributory bronchus. When obstruction of the trachea takes place, death from suffocation occurs before the lungs are completely airless. When obstruction of one main bronchus takes place, atelectasis of the corresponding whole lung occurs. When obstruction of a primary division of a main bronchus takes place, collapse of one lobe of the lung occurs, and so on, until finally obstruction of one terminal bronchus leads to collapse of a small group of alveoli.

Atelectasis may be classified, according to the extent of lung involvement, as (a) massive (one whole lung) and lobar atelectasis and (b) lobular and segmental atelectasis.

This classification is convenient as it corresponds roughly to the different aetiological causes. Atelectasis of a whole lung is usually seen in the condition of carcinoma of the bronchus. Occasionally post operative atelectasis may also involve the whole of one lung. Inhalation of a foreign body may also produce this condition. Trauma to the chest, especially crush injuries, may sever a main bronchus and produce a traumatic stricture leading to a permanent atelectasis of that part of the lung. In childhood the adherence of a tuberculous gland to a main bronchus, with erosion of the wall of the bronchus and the production of an intense oedema of the bronchial mucous membrane, can cause complete obstruction and consequently collapse of a whole lung. This condition may also occur in adult life. Occasionally obstruction of a main bronchus by glandular enlargement due to one of the reticuloses may produce blockage of a primary lobar bronchus, and consequently lobar collapse. Obstructions in the main bronchi or their primary branches are readily visible through the bronchoscope.

Tuberculous tracheo-bronchitis sometimes produces obstruction, either of a main bronchus leading to massive atelectasis, or of a primary (lobar) bronchus leading to lobular atelectasis.





FIG. 50.—The mucus plug. (a) diagrammatic

obstruction (e) a further stage showing the terminal bronchus involved (f) no further movement of the mucus plug but

the affected lung elevation of the diaphragm with displacement of the gastric hyper resonance up into the axilla may lessen the area of dullness and lead to some confusion. The air entry is variable. In the early stage there may be complete absence of breath sounds especially in cases of main bronchial obstruction such as new growth or inhaled foreign body. In other cases loud breath

# ATELECTASIS



Segmental or lobular collapse usually passes unnoticed and the only symptom may be a localized pain over the area


## PHYSICAL SIGNS

When a whole lung is atelectatic the signs are well marked. Flattening and immobility of the affected side of the chest are present. The trachea is deviated towards the affected side and the apex beat is widely displaced towards the same side. The percussion note is dull over the affected lung though if the left is

FIG 51 — (a) Atelectasis of left lower lobe. The heart is slightly displaced to the left and there is an area of emphysema in left costo-phrenic angle. The shadow of the collapsed lobe cannot be seen. (b) crowding together of the lower lobe bronchi with a small amount of dilatation and the lingula bronchi which have come down to occupy the space left by collapsing lobe



(a)



(b)



## ATELECTASIS

sounds of a tubular character may be present from the beginning. Added sounds are not usually present.

In lobar collapse, similar signs to a lesser degree may be present. Thus, the trachea and heart are rarely both displaced. In upper lobe collapse, the trachea is displaced towards the affected side and the heart is not disturbed, the opposite being the case in lower lobe collapse when the heart is displaced towards the affected side and the trachea remains central. Dullness is present over the collapsed lobe which in the case of the lower lobe lies against the spine posteriorly when the area of dullness is triangular in shape and does not extend further forward than the mid axilla. In the case of the upper lobe, the dullness is confined to the anterior part of the upper chest, and is not usually extended below the third rib. The signs produced by collapse of the middle lobe are so slight that they are difficult to elicit. Dullness over the front of the chest, from the fourth rib down wards, may be present, extending to the anterior axillary line. The breath sounds over the collapsed lobe are usually absent, but may be tubular and loud.

## RADIOLOGICAL SIGNS

The radiological signs are characteristic, but vary with the duration of the collapse. In early cases, a homogeneous opacity of characteristic shape (the character varying with the affected part of the lung) will be seen. This opacity is usually attended by the signs of the compensatory mechanisms already outlined—a localized area of compensatory emphysema, deviation of the heart and trachea, and elevation of the whole or part of one diaphragm.

In adult cases, one or all of these radiological abnormalities will be present. In children, especially in cases in which one lobe or less is involved, the compensatory mechanisms may not be present and may only be established if the atelectasis persists with the growth in stature of the child.

## INVESTIGATION AND TREATMENT

In massive or lobar collapse, the possibility of new growth of the lung or foreign body should always be borne in mind and bronchoscopy undertaken. The possibility of tuberculosis should also be investigated by examination of several specimens of sputum for tubercle bacilli. In the more extensive cases, this possibility may be confirmed by the finding of ulcerative lesions in the bronchi.

When the diagnosis is made within a short time of the onset of symptoms then postural drainage with percussion over the affected part of the chest may dislodge the foreign body or mucus plug. In cases of lesser degree an anti-spasmodic, such as ephedrine,  $\frac{1}{2}$  grain by mouth or tincture of stramonium, 15 minims, is worth administering and repeating at 4 hourly intervals. Penicillin should be given to all patients in whom there are signs of infection in the atelectatic lung. The intramuscular route is the method of choice, as the intrabronchial obstruction may prevent the penicillin reaching the area infected if it is administered by inhalation.

## Infection

## COMPLICATIONS

If virulent organisms are trapped behind a bronchial obstruction, an acute febrile




(a)



FIG 53 — (a) Atelectasis of whole of left lung. The trachea is not apparently deviated but no heart shadow is seen to the right of sternum. (b) complete filling of all bronchi with very marked increase in size of left main bronchus and also lower part of trachea. (c) the whole lung is crowded together in to a small area against the posterior chest wall

(b)



(a)

FIG. 52 (a) Collapsed middle lobe. There is a sharp upper demarcating line falling off into normal lung markings in the lower lobe. There is no shift of the mediastinum. (b) the typical shadow of a collapsed middle lobe.



(b)



(a)



FIG. 54 — (a) Complete atelectasis of left lung with right lung being pushed over and filling up part of the space left by the retracted lung. (b) the whole of the atelectatic lung is lying against posterior chest wall.

(b)

## ATELECTASIS



FIG 53(c)

illness is likely to be set up resembling lobar pneumonia and leading possibly to break-down of lung tissue with the formation of a lung abscess. In cases with a less virulent infection the febrile illness will be less dramatic in its onset and course.

### Rupture of an emphysematous bulla

In a number of cases the compensating emphysema which occurs is so localized and so marked that rupture of one of the bullae occurs and the condition of spontaneous pneumothorax is produced. This complication is one that occurs at the same time as the atelectasis and persists after the lung has re-aerated.

## POST OPERATIVE ATELECTASIS OF THE LUNG

This variety of atelectasis is particularly favourable for study and treatment. It occurs most frequently after operations on the mouth, nose and throat but also complicates all types of operations. The variety of anaesthetic used does not influence its incidence to any great extent but intubation of the larynx appears to increase the frequency of its occurrence.

Atelectasis occurs within 48 hours of the operation and may have a dramatic onset resembling that of a lobar pneumonia.

The patient may have a sudden pain in the chest with breathlessness and a rapid rise of temperature up to 104°F. With this type of onset it is usual to find large



(a)



FIG. 54 — (a) Complete atelectasis of left lung with right lung being pulled over and filling up part of the space left by the receding lung. (b) the whole of the atelectatic lung is lying against posterior chest wall.

(b)

# ATELECTASIS



FIG 55 — Atelectasis right lower lobe with heart and mediastinum pulled across towards the same side





FIG 56 — (a) Honeycombing in upper zone of right lung. The trachea is central with no apparent deviation of mediastinum. (b) bronchi of upper lobe are crowded together. (c) complete filling of upper lobe which occupies a very much smaller area than normally with marked crowding together of bronchi and with sacculated dilatation



(b)



(c)



# ATELECTASIS



(a)

FIG 57 — (a) Opacity due to a complete obstruction of left upper lobe bronchus with consolidation of upper lobe and little shrinkage. (b) and (c) bronchograms showing the extent of the upper lobe. Bronchoscopy in this case showed the obstruction to be complete and due to a carcinoma of the bronchus. (d) air has been induced into pleura showing a marked shrinkage of upper lobe



(b)



(c)



(d)

areas of lung involved and consequently physical signs are found. Dullness to percussion over the affected area with deviation of the cardiac impulse (and possibly the trachea) to the affected side are found. The auscultatory sounds are variable. At first there is a complete silence over the dull area but later loud bronchial breathing may be heard and later still numerous moist sounds.

It is more usual to have a quieter onset, in which a low fever is noted on the temperature chart, and on questioning the patient admits to a slight cough dating from the operation. The physical signs also are inconspicuous but definite, small areas, at the bases of lungs of moist sounds are heard with bronchial breathing in the same areas.

The time incidence of the temperature coupled with the physical signs in the chest establish the diagnosis of atelectasis.

The treatment depends on the extent of the lesion and the earliness of the diagnosis.

If a large area of lung is involved and the condition is recognized within 24 hours of the onset then bronchoscopy should be performed and attempts made to aspirate them.

With smaller areas or a later recognition this procedure is valueless, as the mucus is then in the finer bronchi, out of range of the bronchoscope.

Postural drainage should be used and vigorous percussion over the affected area of lung applied in an attempt to dislodge fine plugs and enable the patient to cough the plugs out.

Discretion in the use of these methods is necessary, frail old age after extensive resection operations being treated more gently than robust youth after an appendectomy.

Prophylaxis plays an important part in the treatment of this condition. Pre-operative breathing exercises are valuable but the chief measure is to avoid an operation in the presence of an upper respiratory infection or bronchitis.

In operations on the upper respiratory passage attention to bleeding points and the disposal of small pieces of tonsil, teeth and mucous membranes is of great importance.

Frequent change of posture of the semiconscious patient immediately after operation and for the next 24 hours can also help to minimize the effects of mucus in the air passages.

### THE FATE OF THE ATELECTATIC LUNG

For how long a lobe can remain atelectatic and then become re-aerated is a matter of considerable speculation. When the obstruction is permanent, such as that caused by a carcinoma of the bronchus, a bronchial stricture secondary to a tuberculous process, or an undisturbed foreign body, the collapse is permanent. When the obstruction is temporary, such as that caused by a plug of mucus or oedema of the bronchial mucous membrane, then the atelectasis is as temporary as is the obstruction, and may last for any period from a few minutes to a few weeks. In post-operative cases, in which it is possible to assess the moment of onset, clearing has frequently been found to be complete in 1 week, but has also persisted for 2-3 weeks, and yet ended in complete re-aeration of the affected lung. In cases which persist for longer than 3 weeks, it is safe to say that

although some re-aeration may occur, large areas of atelectasis are likely to persist

The majority of cases do re expand completely, yet such is the frequency of atelectasis that many patients are seen in whom the atelectasis has persisted, often only recognized when one of the late results calls attention to the condition

## BIBLIOGRAPHY

- Gairdner, W T (1850) "The Pathological Anatomy of Bronchitis and the Diseases of the Lung connected with Bronchial Obstruction" *Mon J med sci Edinb*
- Hasse, C E (1846) *An Anatomical Description of the Diseases of the Organs of Circulation and Respiration* Trans by W E Swaine London
- Legendre and Bailly (1844) *Archives generales de medicine journal complementaire des Sciences Medicales* Paris Vol 4
- Pasteur, W (1908) "The Bradshaw Lecture on Massive Collapse of the Lung" *Lancet*, 12, 1351
- Rees, G A (1850) *Atelectasis Pulmonum* London

## CHAPTER 12

### BRONCHIECTASIS

F P LEE LANDER

#### HISTORY

THE WORD "bronchiectasis" first appeared in the English literature in 1846, when the Sydenham Society published Swaine's translation of Hasse's *Pathological Anatomy*. The word is derived from two Greek words—*bronchos* meaning a tube, and *ektasia* to stretch.

In 1821 Laennec had described for the first time a condition of dilatation of the bronchial tubes, his attention having been drawn to the condition by his pupil Andral who had noted it at post-mortem examination. Laennec's description in the first edition of his great work *Médiate Auscultation* was brief, but in the fourth edition the description was enlarged to include the clinical findings of cases which later came to post-mortem examination. He described the condition as occurring in the upper lobes of the lungs, and the clinical differentiation from pulmonary tuberculosis was made mainly on the length of the history and on the appearance of one or two added sounds, to which today we attach no importance. He did not describe clubbing of the fingers as occurring in the condition, nor did he lay any stress on displacement of the heart and trachea. His theory of the pathogenesis was that the dilatation of the bronchial tubes was caused by a copious secretion of sputum distending the bronchi, and the copious secretion of sputum he considered to be the primary disease.

Andral (1831) mentioned the condition, but did not advance any theory as to causation, though he did remark that while generally the dilatation occurred in those who had suffered from chronic cough in children who, previous to the occurrence as a complication of whooping cough, had had no cough at all.

Stokes (1837) drew a parallel between aneurysm of the aorta and dilatation of the bronchi. He suggested that the elastic tissue in the non muscular part of the bronchial wall was destroyed, that there was a paralysis of the circular muscles due to inflammation, and that the cilia lost their power of movement. These three causes, he considered, contributed to the dilatation of the bronchial tubes and the stasis of their contents.

The early theorists regarded the disease as primarily one of the bronchial wall, and considered that the changes occurring in the surrounding lung tissue were secondary to the bronchial change. The first person to challenge this concept was Corrigan, who, in 1836, maintained that the lung changes were primary and that as a result of the contraction in the lung, bronchial dilatation was produced. His theory was that a fibrosis occurred in the lung, secondary to some chronic disease, and that the contraction of this fibrous tissue produced dilatation of the bronchial tubes.

Hasse subscribed to the theory that bronchial dilatation was part of a compensating mechanism, secondary to the collapse of the lung, and stated that bronchitis was one of the major causes. On the other hand, Gairdner, in 1850, came to the conclusion that the so-called bronchial dilatation was in reality ulcerative excavation of the lung, which became refined by bronchial mucous membrane.

Since that time the majority of theories advanced have been in favour of the primary disease being one of the bronchial wall, and much of the research has been directed into proving this theory, showing that disease in the bronchial wall results in the destruction of the elastic tissue, so that as a result of the increased pressure, either of a forced inspiration following a cough or because of the raised intratracheal pressure due to the forced expiration against the closed glottis, a dilatation of the bronchi is produced which becomes permanent. Of recent years, however, there has been a swing in thought to the ideas put forward by Corrigan (1836), the only difference being that whereas he postulated a fibrosis or "cirrhosis" of the lungs secondary to chronic infection, the current theory is that the shrinkage of the lung occurs largely as an acute process following peripheral bronchial obstruction and atelectasis.

Until 1922 the actual proof of the presence of the dilatation was found only at post mortem examination. In 1922, however, Sicard and Forestier introduced the use of opaque oil in radiography for the study of the hollow viscera of the body, thus enabling bronchiectasis to be diagnosed, with certainty, in life. The immediate result of this outstanding contribution to medical science was the stimulation of interest in bronchiectasis and a consequent outpouring of articles. Amongst these was one by Wali and Hoyle (1933), in which the concept of the condition was enlarged from the confines of the heavily infected extensive cases to the almost symptomless cases of dilatation.

Articles by Finlay (1935) and Warner (1934) drew attention to the association between dilatation of the bronchi and collapse of the lobes of the lung. Smith (1930) demonstrated the appearance of spirochaetes in the sputum of bronchiectatic patients, and attributed the bronchiectasis to this organism. Since then however, the presence of spirochaetes has been demonstrated, times without number, in the sputum of bronchitic patients, and those suffering from every type of respiratory disease, with or without bronchial dilatation.

Jennings (1937) published a case in which well marked dilatation of the bronchi was shown later to have disappeared, and the name of "reversible bronchiectasis" has been given to the condition. Some writers state that this condition has nothing to do with permanent bronchiectasis and consequently have labelled it "pseudo-bronchiectasis". Modern American writers in discussing the reversible changes in the bronchi, state that the appearances of the dilatation in a collapsed lobe are possibly due to a shortening of the bronchi, which, at the same time, produces a widening of the calibre. Against this theory is the fact demonstrated by Maclellin (1929) that with expiration the bronchi narrow and shorten. Shortening of the bronchi is not necessarily attended by dilatation. There are cases on record in which collapse of the lung is attended

by shortening of the bronchi with no alteration in calibre This can be demonstrated also in cases of artificial pneumothorax

Andrus (1937) examined in detail all the physical forces in the lung occurring with various conditions, and came to the conclusion that the forces arising as a result of pulmonary atelectasis provide the most satisfactory explanation of the disease

### EXPERIMENTAL WORK

Tannenberg and Pinner (1942), in their experiments, produced a main bronchial obstruction in dogs, and the final state of the bronchi seen at post-mortem examination was found to depend on the presence or absence of infection Where infection has been added the bronchi were found to be dilated, whereas where no infection had been added the bronchi were collapsed and the walls touching

Adams and Escudero (1938) also did a series of experiments on dogs, but they were only able to produce bronchiectasis when the obstruction was incomplete and when contamination of the bronchial tree had occurred In none of their cases with complete obstruction, even in the presence of infection, were they able to demonstrate bronchial dilatation The demonstration of its presence or absence was again made at post-mortem examination

Weinberg (1937) produced bronchial obstruction in rabbits by means of a sand burr Every possible attempt was made to exclude infection, and although this was not successful in each case, bronchiectasis was the universal finding, also demonstrated at post-mortem examination

Davidson and Lee Lander (1938) produced bronchiectasis in cats by the introduction of gum acacia In this case the demonstration of the dilatation was made by means of a bronchogram in the living animal

These varying, and apparently contradictory results must be made to fit into some all-embracing theory of the production of the dilatation In interpreting any experimental work the first essential is to note in each case whether the bronchiectasis was demonstrated by bronchography in life or at post mortem examination This is an important point because, if modern theories are correct, the dilatation is, in the first place, dependent on the normal physical conditions existing in the intact thorax, and removal of the lungs from the thorax *post mortem* destroys these physical conditions Corrigan (1836) was the first to point out the importance of the rigid walls of the thorax for the production of the condition This was also clearly illustrated in one case seen at the Brompton Hospital in the early days of lobectomy, when a lobe was removed from the thorax for the condition of bronchiectasis and subsequent examination of the lobe showed it to be dense and airless but with completely normal bronchi The pre operative bronchograms in this case clearly illustrated the presence of a fusiform bronchiectasis

The second point to note is the position of the obstruction, either central or peripheral All air spaces distal to the obstruction are subject to the same forces, whether the obstruction is central or peripheral, and similarly conditions proximal to the obstruction are identical thus bronchi peripheral to the obstruction will collapse The main difference is that in the case of a central obstruction the bronchial walls are strengthened by cartilaginous rings and are consequently

## EXPERIMENTAL WORK

less easily distensible or collapsible. Once this point is realized a great many of the apparently divergent results in the experimental work will fall into line. The third point to note is the extent of the obstruction in the main bronchus. If the obstruction is incomplete, collapse will not follow unless at the site of the narrowing of the main bronchus mucus forms and renders the obstruction complete.

The theory first propounded by Gairdner (1851), that the bronchial dilatations are produced by an actual destruction of the bronchial wall, with a complete loss of all the normal structures of the wall and the resulting cavity then becoming lined by bronchial mucous membrane still has some adherents (Lisa and Rosenblatt, 1943). In support of this theory the histological appearances of cystic dilatations have been quoted. The point that is overlooked is that the structure of the bronchial wall is not uniform throughout its length. In the region of the lung hilum the bronchial wall is a complicated structure, containing glands, muscle, elastic tissue, fibrous tissue, and cartilage, whereas in the periphery of the lung it is simple and consists of bronchial mucous membrane and a thin layer of muscle and elastic tissue. Consequently, the interpretation of the histological appearances of any cavity associated with the bronchus must be correlated with the position of the cavity in its relation to the bronchus.

The possibility of a congenital origin for bronchiectasis has been raised, and there is a certain group of cases still referred to as congenital cystic bronchiectasis. This theory is an extremely difficult one either to prove or discount. The frequent association between bronchiectasis and other congenital abnormalities, such as transposition of viscera (Adams and Churchill, 1937) or cystic conditions of the pancreas (Anderson, 1938; Baylin, 1944), kidney, and liver, have been used as an argument in favour of the congenital origin. Some writers have suggested that congenital bronchiectasis arises as a maldevelopment or non-development of parts of the primitive lung bud. Others have suggested that obstruction of the bronchial tree occurs during intra-uterine life, probably with amniotic fluid, and that this obstruction prevents expansion of parts of the lung. Neither of these possibilities can be completely ruled out in any case occurring within the first few weeks of life. Many of the cases showing the so-called typical appearance of congenital cystic lung can be shown to occur in children whose lungs have been known previously to be healthy. The proof of the presence or absence of a developmental abnormality must rest finally on the demonstration of cystic spaces in the still-born lung.

It has already been mentioned that peripheral bronchial obstruction is the important factor in the production of bronchiectasis. Whether the patient suffers from the clinical syndrome of long-standing cough, foul sputum, and haemoptyses depends on the second factor, which is the presence of infection.

## AETIOLOGY

Patchy atelectasis occurs in many conditions in childhood. The two most frequent causes of this condition are whooping-cough and measles. These two diseases consequently are the commonest causes of bronchiectasis. Enlarged



tuberculous glands in the hilar region of the lungs may produce lobar collapse and account for a number of cases of bronchiectasis (Brock, Cann and Dickinson, 1937) Brock (1950) drew attention to this as a cause of haemorrhagic bronchiectasis of the right middle lobe.

Inhaled foreign bodies are responsible for a small number of cases, and post-operative atelectasis, more especially following operations in the upper respiratory tract, account for a further number. Later in life there are many cases of atelectasis of obscure origin, which may persist and give rise to permanent collapse and bronchiectasis. Later still carcinoma of the bronchus plays an increasing part in the production of the condition.

Lobar pneumonia has always been ascribed an important role in the aetiology of bronchiectasis, and in cases in which resolution is delayed or incomplete, bronchiectasis does follow. These cases of unresolved pneumonia are probably due to retention of secretion in the consolidated lung leading to a permanent atelectasis. It has been stated that the occurrence of bronchiectasis in cases with asthma is rare, and that atelectasis also does not occur in spite of the bronchial oedema and spasm. A thorough search by bronchography in cases of asthma does not bear out this statement, both atelectasis and bronchiectasis being frequent complications of asthma.

Many children with bronchiectasis give a history of a tracheotomy during an attack of diphtheria and a tracheotomy scar is found on examination of the neck. There may be no connexion between these two facts. Pasteur, however, pointed out that atelectasis was a frequent occurrence in severe cases of diphtheria. It is also possible that the tracheotomy was performed on the mistaken diagnosis of laryngeal obstruction, the obstruction in fact being lower down the respiratory passages.

#### *Case report*

J.T. a small boy aged 3 years was seized with intense dyspnoea and cyanosis. Marked stridor was present and a tracheotomy was performed as an emergency measure. No relief was obtained and the child was admitted to hospital. On admission dyspnoea and cyanosis were present and on examination of his chest complete dullness and absent air entry on the left side were found.

Bronchoscopy was performed. His larynx and trachea were found to be normal. The mucous membrane of the left main bronchus was oedematous and intensely inflamed and the oedema completely obstructed the lumen of the bronchus. The right main bronchus was normal. No cause for this was found. The left lung remained massively collapsed and bronchiectasis was demonstrated in the collapsed lung (see Fig. 58).


Many other factors have been put forward as the cause of bronchiectasis. Infection of the nasal sinuses has been ascribed a major role, and the frequency with which the two conditions coexist has been used as a proof of their relationship. It has been shown that inspiration of opaque oil into the nose at night has produced widespread dissemination of the oil in the lungs the following day. This shows that it is possible for mucus dropping down the back of the nose to be inspired, and possibly produce obstruction and collapse.

Fig. 58 — (a) Massive bronchectasis of left lower lobe 1936. Few bronchi on the right outlined are within normal limits. (b) 1946 complete collapse of left lung with bronchectasis involving all bronchi on left side. Bronchi on middle lobe have already come across the midline of the left thorax but all bronchi outlined are normal.

(a)



(b)



Fibrosis of the lung is not a primary condition, and probably plays no part in the production of bronchiectasis. In cases of silicosis the widespread fine fibrosis is an essential part of the disease, and bronchography has demonstrated normal bronchi in the majority of cases.

Bronchial dilatation is a frequent complication of tuberculous conditions of the lung, more especially in those cases in which atelectasis has played a major role, and in conditions of tuberculous tracheobronchitis, bronchiectasis is an invariable accompaniment.

The histology of bronchiectasis, obtained from post mortem specimens, has nearly always shown the presence of gross infection, and this fact has been used to give infection a dominating role in the production of the condition. Histology of lobectomy specimens on the other hand, when the lobectomy has been performed in early cases or cases in which there were few symptoms, has shown little, if any, infection.

Bacteriology of the sputum is extremely varied. It is unusual to find one organism as the only inhabitant of bronchiectatic sputum, and it is much more usual to find 5 or 6 varieties of organisms, none of them predominating. The bacteriology closely resembles that of bronchitis.

## DIAGNOSIS

### Symptoms and signs

The diagnosis is based on the presence of dilated bronchi, but on infection, if virulent, and on the presence of a chain of signs and symptoms which have come to be regarded as pathognomic of bronchiectasis. Dilated bronchi surrounded by airless lung may lead to recurrent attacks of pneumonia, but whether this is due to a lighting up of a dormant infection or to a superadded infection cannot be stated with any certainty.

A low-grade infection may produce excessive granulations in the dilated bronchi and these granulations may, on occasion, bleed, giving rise to repeated haemoptyses. Finally bronchiectasis being part of a widespread compensatory mechanism, of which displacement of the heart and emphysema are the other manifestations, dyspnoea may be the outstanding symptom. This is especially so in the case of the older patient.

The symptoms of the moderately severe infection leading to chronic ill health are striking. Repeated respiratory crises in childhood are followed by chronic cough and copious sputum, which varies in amount from 1 ounce to 1 pint per day. The character of the sputum is muco-purulent as a rule, and in certain cases is offensive, a peculiar sweet offensiveness, said to be characteristic of the disease. With this productive cough is a series of haemoptyses, as a rule not great in amount but increasing in severity. In a case such as that just described, dyspnoea is always a marked feature, although not the primary complaint. Further inquiry into the incidents in the past may reveal a rib resection for

## DIAGNOSIS

empyema, a tracheotomy for supposed laryngeal obstruction, or an operation for tuberculous glands in the neck. Examination of the patient usually reveals a thin, wasted, slightly cyanosed person, who is dyspnoeic and whose conversation is punctuated by a productive cough. Clubbing of the fingers and cyanosis of the finger-tips are often present, and on examination of the chest there is wasting with recession of the intercostal spaces, a poor chest movement (as a rule the apex beat is neither visible or palpable) and the trachea is central. Percussion note is hyper-resonant at the apices, and the note at the bases is variable. Patchy dullness may be present, though hyper-resonance is just as likely.

On auscultation the first, and most noticeable, finding is the presence of râles of all calibres over all areas of both lungs, but more marked at the bases. Should the patient get rid of a good deal of sputum during the examination the air entry, which previously may have been poor, becomes loud and bronchial in character. Further examination of such a patient will show the clubbing of the toes and possibly the presence of some signs of amyloid disease, albuminuria or enlargement of the liver.

X-ray examination of the lungs will show widespread changes throughout the whole of both lungs, with mottled opacities which are more markedly present at the bases. Bronchography as a rule is not very satisfactory, the patient's tolerance to intratracheal interference being low and any Lipiodol introduced is so diluted by the sputum that no clear picture of the degree of the bronchial obstruction is obtained. Post-mortem examination, however, reveals the widespread nature of the complaint.

Cases of this nature usually originate in severe attacks of whooping-cough. Cases of less marked degree, and especially in which one lower lobe (usually the left) is affected are more common, and some of them exhibit a syndrome closely allied to that already outlined. In such cases, however, the physical signs are more localized and mediastinal displacement is obvious. The heart is displaced to the left. There is a small area of dullness close to the spine posteriorly, with a hyper-resonant note outside this area. Breath sounds are weak in the axilla, and may be absent over the dull area. Coarse râles are present, confined to the area of dullness. If such a patient is then tipped head downwards he will produce, without much effort, 3-4 ounces of thick pus, which may or may not be offensive. The air entry over the dull area will then be loud, and the breathing cavernous in character.

It is rare to find the dilatations confined to one lobe. When the left lower lobe is bronchiectatic, the lingular division of the left upper lobe is usually involved and frequently there is some involvement of the right lower lobe. Occasionally cases are seen in which the middle lobe bears the brunt of the disease and then the corresponding bronchi on the left side may be involved as well. Sometimes the only symptoms are those of bronchitis. These cases should not be regarded as cases of infected bronchiectasis. The cases of infected bronchiectasis are those in which the infection has spread through the bronchial wall and is located in the pielectatic lung tissue. It is this infection that gives rise to the toxæmia which is characteristic of the disease. This smouldering infection may occasionally burst into flame, giving rise to an illness closely resembling lobar pneumonia.

Patients are also encountered who complain of cough and sputum in damp and foggy weather, but who are symptomless in spells of fine weather. This is such a common occurrence in patients with normal bronchi living in large cities that it alone should not be regarded as being directly connected with the bronchial dilatation.

Occasionally patients with a history of productive cough of some years' standing will say that they have had 4 or 5 attacks of pneumonia. This history is of great significance, especially if the same lobe of the lung has been involved in each attack of pneumonia.

In some of the series of observed cases of bronchiectasis, sputum has dated from a recent acute upper respiratory infection whilst it has been known that the bronchial dilatation has been present for a great many years but hitherto has been virtually symptomless.

Cases are also seen in which the only complaint is that of recurring haemoptysis. As a rule clinical examination of these patients does not reveal any gross abnormality though an area of bronchial breathing and a few râles may be discovered. Radiological examination, however, may indicate a localized abnormality sufficient to warrant bronchography.

Other cases are detected by the routine investigation of certain clinical and radiological abnormalities. These may consist of cases of unresolved pneumonia, if completely investigated. Numbers of cases of asthma, in which, after treatment, are revealed, again reveal dilated bronchi. Post-operative lung conditions, if persistent, will often show a similar state of affairs.

### Radiological appearances

The important radiological signs are (a) those of the other compensatory mechanisms secondary to collapse, and (b) the patchy opacities associated with infection in the atelectatic lung.


In the early stages, and when the collapse is of recent origin, the atelectatic area may throw a shadow on the x-ray film, and the shape of the shadow is characteristic for the area involved. In cases of longer standing, no shadow of the atelectatic area will be seen, and the radiological appearances will be those of the compensatory mechanisms, that is, elevation of the corresponding half of the diaphragm with limitation of movement, a localized area of emphysema, and displacement of the mediastinum to the affected side (and this displacement of the mediastinum includes such changes as elevation of the pulmonary artery and peaking of the pericardium either on the right or left). Occasionally the condition is associated with infection in the lungs.

If opaque areas are seen in the lungs, these will be found to be crowded together. The condition may be manifest, and that is the great increase in the size of the bronchi with inspiration, and the return to a basic dilated level with expiration.

Associated with these changes there may also be a movement of the diaphragm. The diaphragm moves to a higher level with inspiration and returns to its normal position with expiration (see Figs 59-60).

FIG. 59 —Bronchectasis of long standing. Bronchograms showing (a) the size of the bronchi in expiration and (b) the size of the same bronch in inspiration.

(a)



(b)







FIG 60 — (a) Inspiratory and (b) expiratory films in a patient with bronchiectasis of at least 17 years standing showing increase in calibre with inspiration



It has been customary to classify bronchiectasis according to the bronchographic appearances, and cylindrical, fusiform, and saccular bronchiectasis have all been described. These distinctions, although entirely artificial, are of some use clinically, as it will be noted that the cases of saccular dilatation as a rule are the cases of more widespread involvement of the lung, and in which dyspnoea is the outstanding symptom, whereas cylindrical dilatation is as a rule localized.

The diagnosis of the condition of bronchiectasis depends on the demonstration of dilated bronchi by means of opaque oil in the bronchial tree. The method of introduction of the oil is a matter for the operator's preference. Pleas have been advanced for methods where the oil is introduced over the tongue or through the nose, as involving less trauma and less distress to the patient. In a large series of cases in which repeated bronchograms were performed by the author and various methods used, the experience was that the patients preferred injection through the crico-thyroid membrane, and the only explanation put forward for this is that less manipulation of the tongue and cocaineization of the throat is necessary by this method, and that it takes less time.

The most important part of any method is the posturing of the patient, so that either the desired area of lung is filled with the radio opaque oil, or preferably, the whole of the bronchial tree is filled on the one side. In order to get the most satisfactory and easily interpreted results two separate manipulations are necessary. At the first, the diseased part is completely filled, and after an interval, of days or weeks depending on the amount of residual Lipiodol the other side is filled. The modern technique is to fill one side completely and then go on immediately to fill the other side, and take oblique x-ray photographs. These pictures are somewhat difficult to interpret, as overlapping of the bronchi must occur sometimes, giving the appearance of bronchiectasis where none may be present.

The amount of iodized oil used is about 15-20 millilitres. This should be sufficient to fill one side of the lung completely, and it is better if used cold (as instructed by the manufacturers). Alveolar filling is annoying, as it obscures the final pictures. It is unpredictable and perhaps unavoidable, but the author found it to be less frequent when cold heavy deposit contrast oil was used.

Bronchography carried out on children is apt to be disappointing, and alveolar filling is extremely common. The explanation of this is probably that the oil is put in under a general anaesthetic, through a bronchoscope, and that the pressure applied through the syringe and catheter causes the oil to be squirted with some force into the finer bronchi.

Bronchoscopy is a useful adjunct to bronchography in the diagnosis, and is an essential preliminary to the surgical treatment of bronchiectasis. It enables pus to be seen welling out of the bronchi and may help the decision as to which is the most heavily infected of the bronchiectatic areas. A fibrous stricture suggests a tuberculous or traumatic origin for the dilatation. Unsuspected foreign bodies are sometimes found and can be removed.



As part of the preliminary investigations tipping of the patient into the various positions required for drainage of the affected bronchi is essential. In the more grossly infected cases, as a result of tipping, 1 or 2 pieces of thick, purulent sputum may be obtained. The non-production of sputum by this method does not necessarily rule out the presence of infection, and this is especially true in the case of children. The children may be held in the appropriate position for minutes at a time and produce no sputum at all, whereas subsequent bronchoscopy or other manipulation may result in a copious production of foul, purulent sputum.

Other investigations which are helpful are a white cell count and possibly sedimentation rate. In the majority of cases in which infection exists, a leucocytosis is present, and in these cases the erythrocyte sedimentation rate is usually raised. As a rule bronchiectasis is not associated with fever except in the acute incidents.

### COURSE OF THE DISEASE

As already indicated, in some cases the infection is so overwhelming that death occurs within a short period of the establishment of the bronchial dilatation. These cases more closely resemble an acute lung abscess; they were always

acute exacerbation of the lung condition (see Figs 61-67). This type of case may develop at any stage a cerebral abscess which in itself may prove fatal.


These cases of bilateral involvement have been heavily infected from the beginning. They have also been bilateral from the beginning. The spread of bronchiectasis from one part of the lung to another is extremely rare, and when it does occur, it happens as a result of a plugging of further bronchi, with consequent atelectasis. The spread of bronchiectasis by infiltration, such as happens in pulmonary tuberculosis, does not occur.

The incidence of infection occurring in a previously collapsed uninfected lung is extremely hard to determine. In a series of over 100 cases followed by the author for 13 years, this did not happen in a single instance. In the same series, repeated bronchograms were performed, and spread occurred in one case only, and then finally disappeared.

vast majority of patients seen and diagnosed in Great Britain. In the patients followed by the author, 53 per cent married and 31 per cent had children, a sequence of events not likely to occur if the patients were as repellent as has been suggested. It was found that the pregnancy did not affect the course of the disease, and the disease did not adversely affect the pregnancy.



(a)



(b)

FIG. 61 — (a) Complete collapse of left lower lobe (1933) with markedly dilated bronchi present in collapsed lobe. Both divisions of the Ingula are also involved. Filling on the right is poor but there is possibly some dilatation of the bronchus seen close to the vertebral border. (b) 1948 no appreciable change, no further bronchiectasis in left upper lobe, and all bronchi outlined on right side are within normal limits except possibly the one seen close to the vertebral border.




Fig 62 (a)



Fig 62 (b)

FIG. 62 — (a) Outline of a collapsed left lower lobe with some abnormally heavy shadowing in the right lower zone (1934), (b) poor filling but confirming the collapse of the left lower lobe and showing dilated bronchi in it (1934). Bronchi elsewhere insufficiently filled, (c) collapsed left lower lobe containing dilated bronchi (1948). The filling of the lingula bronchi is poor but shows what is possibly some bronchiectasis. The other bronchi on the left side are normal. Bronchi on the right are dilated in the middle lobe but otherwise normal.



Figures in the past have shown an extremely heavy mortality in medically treated cases. Most of these studies were in the early years of the use of Lipiodol, and it can only be assumed that the heavily infected cases predominated.

Bradshaw, Putney and Clerf (1941) in a series of 171 cases of bronchiectasis followed from 1925 to 1935 had 59 deaths (34.5 per cent) during the period under observation.

Perry and King (1940) in a detailed analysis of 144 cases found that 41 per cent died within 5 years of onset and only 15 per cent remained alive for 20 years or longer.

Roles and Todd (1933), out of a total of 45 patients treated medically, found that 23 were dead within a short period.

### COMPLICATIONS

The main complication of bronchiectasis is the recurrence of attacks of infection in the atelectatic lung and in the dilated bronchi. The incidence of this complication has never been seriously worked out, but it is undoubtedly high. The severity of infection varies from case to case and may be in the nature of a severe pneumonia, or of a trivial febrile attack associated with an increase in cough and sputum.

Cerebral abscess is listed as one of the most frequent complications, and occurs mainly in the heavily infected patient. No case of cerebral abscess in an uninfected patient has been recorded.

# BRONCHIECTASIS



FIG. 63 — (a) Complete collapse and bronchiectasis of the left lower lobe (1932). The lingula filling is poor but suggests bronchiectasis. The bronchi filled on the right side are within normal limits. (b) collapse of the left lower lobe with ectatic bronchi (1948) and bronchiectasis of the lingula. The bronchi filled on the right side are within normal limits.

Amyloid disease, although it occurs, is not nearly so common as in the corresponding stages of pulmonary tuberculosis

Rheumatism, including infective arthritis, is a rare complication

Nephritis has been reported on several occasions

## TREATMENT

The aim of treatment in bronchiectasis must be to rid the bronchi of infection. Re-establishment of normal calibre is not possible, except in a small number of patients in whom reversion to normal takes place without medical aid. The eradication of infection depends on the drainage of the affected bronchi and the administration in one form or another, of drugs designed to kill the organisms prevailing in the sputum.

### Postural drainage (Fig. 68)

The patient must adopt a posture in which the affected bronchi are at a higher level than the bifurcation of the trachea, so that the sputum may run down to this point, whence it can be voided by the patient. Thus, when the ventral (or anterior) bronchi are affected, the patient lies on his back, and when the dorsal (or posterior) bronchi are affected, he lies on his face. Drainage is also assisted by tilting the patient, so that when the lower bronchi (those below the bifurcation of the trachea) are involved the patient lies with the foot of the bed raised, and when the upper bronchi are involved drainage takes place naturally as the patient is standing. As already mentioned, it is unusual to find a case of infected upper lobe bronchiectasis.

The drainage is further helped by inclining the patient so that he lies with his uninfected side uppermost. Therefore in the case of right dorsal lobe involvement the patient should lie partially on the left side, face uppermost, and in right middle lobe involvement he should lie partially on the left side, face downwards.

These postures are easily maintained for a considerable period of time, and consequently are the ones to be recommended in the initial treatment. More drastic drainage can be intermittently supplied by making the patient hang over the edge of the bed, with the body inclined vertically downwards—this, of course, for the patient with lower lobe infection. This posture cannot be maintained for a long time, and usually produces a severe headache. Postural beds and frames have been designed so that the maintenance of these postures can be more easily carried out, and they play a useful part, particularly in cases of

hung over the top of the frame

Occasionally, the combination of bronchi involved is such that more than one posture is necessary to produce effective drainage. The length of time allowed for each posture is governed by the extent of the infection in various areas, but it can be stated generally that the lower lobe bronchi, with possibly the middle lobe and lingula bronchi, are more heavily infected than the bronchi of the upper lobe.



(a)



(b)

FIG 63 — (a) Complete collapse and bronchiectasis of the left lower lobe (1932). The lingula filling is poor but suggests bronchiectasis. The bronchi filled on the right side are within normal limits. (b) collapse of the left lower lobe with ectatic bronchi (1948) and bronchiectasis of the lingula. The bronchi filled on the right side are within normal limits.

Amyloid disease, although it occurs, is not nearly so common as in the corresponding stages of pulmonary tuberculosis

Rheumatism, including infective arthritis, is a rare complication

Nephritis has been reported on several occasions

## TREATMENT

The aim of treatment in bronchiectasis must be to rid the bronchi of infection. Re-establishment of normal calibre is not possible, except in a small number of patients in whom reversion to normal takes place without medical aid. The eradication of infection depends on the drainage of the affected bronchi and the administration, in one form or another, of drugs designed to kill the organisms prevailing in the sputum.

### Postural drainage (Fig. 68)

The patient must adopt a posture in which the affected bronchi are at a higher level than the bifurcation of the trachea, so that the sputum may run down to this point, whence it can be voided by the patient. Thus, when the ventral (or anterior) bronchi are affected, the patient lies on his back, and when the dorsal (or posterior) bronchi are affected, he lies on his face. Drainage is also assisted by tilting the patient, so that when the lower bronchi (those below the bifurcation of the trachea) are involved the patient lies with the foot of the bed raised, and when the upper bronchi are involved drainage takes place naturally as the patient is standing. As already mentioned, it is unusual to find a case of infected upper lobe bronchiectasis.

The drainage is further helped by inclining the patient so that he lies with his uninfected side uppermost. Therefore in the case of right dorsal lobe involvement the patient should lie partially on the left side, face uppermost, and in right middle lobe involvement he should lie partially on the left side, face downwards.

These postures are easily maintained for a considerable period of time, and consequently are the ones to be recommended in the initial treatment. More drastic drainage can be intermittently supplied by making the patient hang over the edge of the bed with the body inclined vertically downwards—this, of course, for the patient with lower lobe infection. This posture cannot be maintained for a long time, and usually produces a severe headache. Postural beds and frames have been designed so that the maintenance of these postures can be more easily carried out, and they play a useful part, particularly in cases of bilateral infection. If this is of the lower lobes the patient should have the frame at an angle of 100 degrees and should lie over the frame on his face, learning if possible, to sleep in this position. If the infection is of the lingula and middle lobe the same frame is used but the patient should lie on his back with his knees hung over the top of the frame.

Occasionally, the combination of bronchi involved is such that more than one posture is necessary to produce effective drainage. The length of time allowed for each posture is governed by the extent of the infection in various areas, but it can be stated generally that the lower lobe bronchi, with possibly the middle lobe and lingula bronchi, are more heavily infected than the bronchi of the upper lobe.



# BRONCHIECTASIS



(a)

FIG. 63 — (a) Complete collapse and bronchiectasis of the left lower lobe (1932). The lingula filling is poor but suggests bronchiectasis. The bronchi filled on the right side are within normal limits. (b) collapse of the left lower lobe with ectatic bronchi (1948) and bronchiectasis of the lingula. The bronchi filled on the right side are within normal limits.



(b)

## TREATMENT

Amyloid disease, although it occurs, is not nearly so common as in the corresponding stages of pulmonary tuberculosis

Rheumatism, including infective arthritis, is a rare complication

Nephritis has been reported on several occasions

## TREATMENT

The aim of treatment in bronchiectasis must be to rid the bronchi of infection. Re-establishment of normal calibre is not possible, except in a small number of patients in whom reversion to normal takes place without medical aid. The eradication of infection depends on the drainage of the affected bronchi and the administration, in one form or another, of drugs designed to kill the organisms prevailing in the sputum.

### Postural drainage (Fig. 68)

The patient must adopt a posture in which the affected bronchi are at a higher level than the bifurcation of the trachea, so that the sputum may run down to this point, whence it can be voided by the patient. Thus, when the ventral (or anterior) bronchi are affected, the patient lies on his back, and when the dorsal (or posterior) bronchi are affected, he lies on his face. Drainage is also assisted by tilting the patient, so that when the lower bronchi (those below the bifurcation of the trachea) are involved the patient lies with the foot of the bed raised, and when the upper bronchi are involved drainage takes place naturally as the patient is standing. As already mentioned, it is unusual to find a case of infected upper lobe bronchiectasis.

The drainage is further helped by inclining the patient so that he lies with his

These postures are easily maintained for a considerable period of time,

for a long time, and usually produces a severe headache. Postural beds and frames have been designed so that the maintenance of these postures can be more easily carried out, and they play a useful part, particularly in cases of

hung over the top of the frame

Occasionally, the combination of bronchi involved is such that more than one posture is necessary to produce effective drainage. The length of time allowed for each posture is governed by the extent of the infection in various areas, but it can be stated generally that the lower lobe bronchi, with possibly the middle lobe and lingula bronchi, are more heavily infected than the bronchi of the upper lobe.



FIG. 64 — (a) Bronchiectasis of left lower lobe and some ectatic bronchi in left upper lobe including the lingula and probably the pectoral division (1930). The very few bronchi on the right that are filled are within normal limits. (b) 1936 similar condition on left side, right side is not filled. (c) 1950 no extension of disease. The marked spinal curvature is secondary to an attack of poliomyelitis.



(b)



(c)

The second method of procuring drainage is to insert a bronchoscope and apply suction. This has been done as frequently as twice a week, as a routine, for years, but is a poor substitute for well applied postural drainage.

One of the early operations on the lung was designed to drain bronchiectatic cavities through the chest wall. This operation has been abandoned for some years, but occasionally it has been performed as a result of a mistaken diagnosis of lung abscess. Bronchiectatic cavities are multiple; it is therefore ineffective.

It is sometimes necessary to augment postural drainage by percussion therapy carried out while the patient is in the proper posture, and accomplished by sharply tapping the chest over the affected bronchi for a period of about one minute.

#### Introduction of penicillin and the sulphonamides

In diseases of the lung the two most effective methods of applying penicillin are by intramuscular injection and by inhalation. There is still some doubt as to which of these two methods is the more effective, and it is really a matter for individual choice. Intramuscular injection produces sputum devoid of penicillin sensitive organisms in 4-5 days. Inhalation of penicillin produces the same results, and is much more impressive to the patient and consequently has a marked psychological effect.

Penicillin by injection should be given in doses of 250,000 units 4 times a day, and the course should last for at least 2 weeks. If ineffective in this time it is

# BRONCHIECTASIS





FIG 65 — (a) Bronchiectasis of right lower lobe with some dilated bronchi in lower zone of left lung (1927). Filling is too poor to identify these bronchi or to state the condition of lingula. (b) 1937 similar distribution of bronchiectasis also showing that upper lobes on both right and left side are perfectly normal. (c) shows that the bronchiectasis on right side is situated in lower lobe, the middle lobe is normal. (d) 1945 no gross abnormality apart from a high d a phragm on right side


(a)



(b)



(c)




(d)



FIG 66—(a) Localized emphysema at left base and shadow of collapsed lobe behind heart (1927) (b) 1931 confirming this and showing bronchiectasis of left lower lobe and lingula with healthy upper lobe (c) 1937 identical condition the bronchi on right side are normal, (d) 1948 similar condition though with rather a poor filling

(a)



(b)

# TREATMENT

(c)



(d)






FIG 67 — (a) Bronchiectasis of left lower lobe and of the bronchi in the region of right middle lobe (1930) (b) 1948 filling is incomplete but all bronchi of upper lobe appear normal (c) the right middle lobe appears normal in a right posterior oblique view but there is one ectatic bronchus in the dorsal lobe on right side and also one or two in pectoral division of upper lobe. These may be the bronchi outlined in (a)

(a)



(b)



(c)

unlikely to be of any continued benefit but examination of the sputum at the end of the fortnight must be made to find whether the organisms present are insensitive to this form of treatment

If after 2 weeks treatment there has been little decrease in the amount of sputum and the organisms present are found to be Gram negative a short course of streptomycin 0.5 gramme twice a day for 2 days may help to clear the infection

Penicillin by inhalation is given in short spells of about 10 minutes duration during which time 200 000 units of penicillin in 10 millilitres of solution are atomized and inhaled. Any form of atomizer is effective though for hospital treatment one of the machines in which the current of air is produced by an electric motor is probably the best. These machines can be hired for use in the home but there are also some effective hand spray instruments which are inexpensive and can be used and maintained by the patient without any great skill or care.

Chloromycetin can be used instead of streptomycin 250 milligrams 4-hourly by mouth for 5 days. This antibiotic can be used in combination with penicillin therapy in all cases which show a mixed flora in the sputum.

Occasionally the sputum is found to contain streptococci or Gram positive organisms which are insensitive to penicillin but sensitive to the sulphonamides. 1 or such cases sulphamerazine 2 grammes 4-hourly for 2 days and then 1 gramme 3 times a day for 2 weeks may be effective.

These methods are usually adequate to deodorize the offensive sputum and

# BRONCHIECTASIS




FIG 68 — Positions to be adopted for efficient drainage of the bronchi

also reduce it considerably in bulk. However, should the offensive odour persist creosote capsules, 3 grains, 3 times a day by mouth are effective.

As a result of these combined forms of treatment, a remarkable improvement in individual cases is recorded. The penicillin and sulphonamides are, as stated, usually maintained for 2 weeks, but postural drainage may have to be undertaken for much longer, and indeed in many cases, has to be maintained for life.

### Indications for surgery

No case should be presented for surgery until a prolonged period of observation and postural drainage has been tried. After what length of time bronchiectasis becomes irreversible is not certain and on this score alone 1 year should be allowed to elapse from the date of the first demonstration of the condition before a decision as to surgery is made. In that year the case may with postural drainage and antibiotics have become symptomless and even if the dilated bronchi are still present operative treatment is not justifiable.

The practicability of the removal of a lung rests with the surgeon, and with the surgeon only. The amount of lung which the surgeon is willing to remove

it must be undertaken as a life saving measure. The removal of segments of the lung has opened up a wider field, and has rendered operation possible in cases previously judged to be inoperable.

In the treatment of bronchiectasis there are several indications for surgery. There are certain patients who are so weighed down by the presence of the symptoms even though these symptoms may not be marked that surgery may have to be advised on psychological grounds as this type of patient resents any abnormality which distinguishes him from his fellow men and having to undergo prolonged postural drainage is as repugnant to him as is the 4-5 ounces of sputum.

Haemoptysis of gradually increasing severity and of increasing frequency is another reasonable ground for suggesting lobectomy or pneumonectomy.

Repeated major infections in the bronchiectatic area constitute another reason for surgery. Infection in the bronchi does not except in the case of children, give sufficient trouble to justify lobectomy as the infection can be controlled and probably eradicated by the means already outlined.

Certain cases fail to respond to the medical treatment outlined and the infection at the end of the preliminary period is as heavy as it was beforehand. Such cases should be re-considered after a period of observation, and if the symptoms persist and there is evidence of septic absorption operation should be advised.

Surgery has been advised on several occasions on prophylactic grounds, the prophylaxis being the prevention of spread and of infection in previously uninfected dilated bronchi. These grounds cannot be justified as it can be shown that infection and spread are neither inevitable nor frequent.

### BIBLIOGRAPHY

- Adams R. and Churchill E. D. (1937) "Situs Inversus Sinistis and Bronchiectasis" *J. thorac. Surg.* 7: 206.  
 Adams W. E. and Escudero L. (1939) "The Aetiology of Bronchiectasis" *Tubercle, Lond.* 19: 351.

# BRONCHIECTASIS

- Anderson D H (1933) Cystic Fibrosis of the Pancreas and its Relation to Coeliac Disease *Amer J Dis Child* 56, 344
- Andral G (1831) *A Treatise on Pathological Anatomy* Vol 2 Trans by R Townsend and W West Dublin
- Andrus P M (1937) Bronch ectasis An Analysis of its Causes *Amer Rev Tuberc* 36 46
- Baylin G J (1944) Pulmonary Changes in Chronic Cystic Pancreatic Disease *Amer J Roentgenol* 52 303
- Bradshaw H H Putney F J and Clerf L H (1941) The Fate of Patients with Untreated Bronchiectasis *J Amer Med Ass* 116 2561
- Brock R C (1950) Post tuberculous Broncho stenosis and Bronch ectasis of the Middle Lobe *Torax* 5 5
- Cann R J and Dickinson J R (1937) Tuberculous Mediastinal Lymphadenitis in Childhood Secondary Effects on the Lungs *Guy's Hosp Rep* 87, 295
- Corrigan D J (1836) On Cirrhosis of the Lungs *Dublin med J* 13, 266
- Davidson M and Lander, F P Lee (1938) The Aetiology of Bronch ectasis *Brit J Radol* 11 65
- Findlay L (1935) Atelectatic or Compensatory Bronchiectasis *Arch Dis Childh* 10 61
- Gardner W T (1851) On the Pathological States of the Lung connected with Bronchitis and Bronchial Obstruction *Mon J Med Sci Ednb* 13 238
- Hasse C E (1846) *An Anatomical Description of the Diseases of the Organs of Circulation and Respiration* Chap 8 Trans by W E Swaine London
- Jennings G H (1937) Re-expansion of the Atelectatic Lower Lobe and Disappearance of Bronchiectasis *Brit Med J* 2, 963
- Laennec R T H (1834) *A Treatise on the Diseases of the Chest and on Mediate Auscultation* 4th ed Trans by John Forbes London
- Lisa J R and Rosenblatt M B (1943) *Bronch ectasis Pathogenesis Pathology and Treatment* London Oxford University Press
- Macklin, C C (1929) The Musculature of the Bronchi and Lungs *Physiol Rev* 9 1
- Perry K M A and King D S (1940) Bronch ectasis study of prognosis based on a follow up of 400 patients *Amer Rev Tuberc* 41 531
- Roles F C and Todd, G S (1933) Bronchiectasis *Brit med J* 2 639
- Sicard, J A and Forestier J (1922) Methode Generale d'Exploration Radiologique par l'Huile Iodee (Lipiodol) *Bull Soc med Hôp Paris* 10 463
- (1926) Radiological Exploration with Iodised Oil *Brit J Radol* 31, 239
- Smith D T (1930) Etiology of Primary Bronchiectasis *Arch Surg Chicago* 21, 1173
- Stokes W (1837) *A Treatise on the Diagnosis and Treatment of Diseases of the Chest* Dublin
- Tannenberg J and Pinner M (1942) Atelectasis and Bronchiectasis *J thorac Surg* 11, 571
- Wall C and Hoyle J C (1933) Observations on Dry Bronchiectasis *Brit Med J* 1 597.
- Warner W P (1934) Massive Atelectatic Bronch ectasis *Quart J Med* 3 401
- Weinberg J (1937) Experimental Production of Bronchiectasis *J thorac Surg* 6, 402

## CHAPTER 13

### INTRA-THORACIC MANIFESTATIONS OF SARCOIDOSIS

J G SCADDING

#### DEFINITION

HISTORICALLY the concept of sarcoidosis as a systemic disease has arisen from correlation of partial descriptions each referring only to manifestations in one organ, sometimes with passing reference to possible changes in one or more of organs. The syndrome has received a confusing variety of names, including sarcoidosis, non caseating tuberculosis, pseudo tuberculosis, lymphogranulomatosis benigna, and eponyms such as Boeck's disease, Besnier-Boeck's disease, and Schaumann's disease. The word sarcoidosis has the advantage that it implies no etiological claim and that it may be interpreted flexibly. In the absence of agreement about aetiology, the best criterion of definition is histology. Sarcoidosis may thus be defined as a disease characterized by the presence in

ceeding in the older lesions to conversion into a peculiar hyaline type of fibrous connective tissue. The clinical observations which have led to the concept of a systemic disorder characterized by this histological picture have been well summarized by Schaumann (1936), Pinner (1938), Snapper and Pompen (1938) and Longcope (1941).

The possible manifestations and combinations of manifestations of sarcoidosis are numerous. The skin may be involved in several ways including sarcoids of Boeck and the lupus pernio of Besnier. There may be localized or generalized enlargement of lymph nodes, and enlargement of the spleen. The liver usually shows microscopic lesions, but these relatively rarely give rise to clinically evident enlargement or to symptoms. The eyes may show a chronic irido-cyclitis which may be associated with enlargement of the salivary glands giving rise to Heerfordt's syndrome or uveo-parotid fever. The small bones of the hands and feet may show the changes of Jungling's osteitis tuberculiformis or multiplex cystoides. Rarer manifestations include endocrine changes, notably diabetes insipidus from lesions of the pituitary; and various forms of involvement of the central nervous system, of the kidneys, of the heart and of the gastro-intestinal tract. The lungs and intra thoracic lymph nodes are involved in a



not been found during life, they have been demonstrated, albeit with difficulty, at necropsy. There is no clear distinguishing characteristic between chronic miliary tuberculosis and diffuse pulmonary infiltrations of sarcoid type; every gradation between the two extreme types occurs. Many reports of post-mortem examinations have been published in which typical sarcoid lesions were present in some organs, typical caseating tuberculous lesions with bacilli present in other organs, and intermediate types of lesions elsewhere. The evidence thus appears

definition of sarcoidosis, has up to now been proved to be capable of causing lesions of sarcoid type, this obviously does not exclude the possibility that some such agent may be found in future.

## DIAGNOSIS AND DIFFERENTIAL DIAGNOSIS

Where there are other manifestations of sarcoidosis, particularly skin lesions, generalized lymph adenopathy, and irido-cyclitis with or without salivary gland involvement, the diagnosis of accompanying intra thoracic changes usually presents little difficulty. Difficulty is most frequently met in those cases in which the presenting manifestation is a pulmonary infiltration.

In such cases the conditions which most frequently have to be considered in differential diagnosis are tuberculosis, either of chronic miliary type in the diffuse infiltrations, or of old burnt out fibrotic type in the later stages of pulmonary sarcoidosis, industrial lung disease, notably that due to exposure to beryllium compounds, diffuse nodular fibrosis of the lungs of obscure aetiology, and occasionally fungus diseases of the lung. Histology is the only objective method of supporting a clinical diagnosis of sarcoidosis, but since the definition here adopted demands that the characteristic changes should be present in all affected tissues, the discovery of such changes in single biopsy specimens is not absolute proof of the diagnosis, but is evidence to be added to clinical, radiological and other observations. In all cases a careful search should be made for superficial lymph nodes, in cases of intra thoracic sarcoidosis lymph nodes above the clavicles, especially just above and behind their inner ends, and in the axillae are often found to be palpable on careful search. It is always worthwhile obtaining biopsy of any clinically enlarged lymph node from any site in a suspected case of sarcoidosis. When no palpable lymph nodes can be found, aspiration liver biopsy, even though there is no clinical evidence of involvement of that organ, often shows characteristic lesions.

In differential diagnosis from caseating tuberculosis, histological evidence is not incontrovertible, since it is well recognized that lymph nodes at a distance from a focus of caseating tuberculosis may show non-caseating tuberculous lesions indistinguishable from those of sarcoidosis. The points which suggest a diagnosis of sarcoidosis are a general condition better than would be expected in a patient






FIG 69 — Enlarged hilar lymph nodes in the early stages of pulmonary sarcoidosis. There are also some rather indefinite abnormal shadows in the lungs.



FIG 70 — Fine diffuse mottling due to pulmonary sarcoidosis with slight enlargement of hilar lymph nodes.




FIG. 71 — Coarser diffuse mottling with out obvious enlargement of hilar lymph nodes due to pulmonary sarcoidosis




FIG. 72 — Late stage pulmonary sarcoidosis with fibrosis in middle zones of lungs and emphysematous changes elsewhere in the lungs. In the original film the residue of diffuse pulmonary mottling was evident (same case as in fig. 69 five years later)

## INTRA-THORACIC MANIFESTATIONS OF SARCOIDOSIS

Adopting a purely histological definition of sarcoidosis, it is found that a high proportion of cases have negative Mantoux reactions, often to as much as 10 milligrams of old tuberculin, on the other hand moderate sensitivity, say to 1,000 old tuberculin, cannot be regarded as incompatible with the diagnosis. It is often the benign course of the disease which determines the label to be applied, so that the application of the term sarcoidosis might be regarded as indicating a prognosis rather than a diagnosis.

In all cases showing diffuse pulmonary infiltrations a careful and complete occupational history must be taken. Difficult diagnostic problems arise in cases with a possible exposure to noxious dust and radiographic changes in the lungs which could be interpreted as due to sarcoidosis. In such instances all available evidence must be examined, and it is sometimes only after a period of observation that an opinion can be given. Special difficulty arises in the case of beryllium granulomatosis of the lungs, since in this condition the histological changes as well as the radiographic appearances and symptoms are similar to those of sarcoidosis, in fact on a histological definition of sarcoidosis it would be logical to refer to these cases as beryllium sarcoidosis. Estimation of the beryllium content of the urine (which is a specialized procedure) may be helpful in such cases.

### Mantoux-negative erythema nodosum

An interesting group of cases of erythema nodosum with bilateral hilar lymph-node enlargement and negative Mantoux reaction appears to have a relation to sarcoidosis. This group is clearly demarcated from the more usual type of erythema nodosum associated with primary tuberculous infection, in which there may be hilar lymph-node enlargement, but this is characteristically unilateral, and the Mantoux reaction is invariably strongly positive. In cases of Mantoux-negative erythema nodosum with bilateral hilar lymph-node enlargement, diffuse pulmonary infiltrations, which generally prove to be transient, are frequently seen, and in a few cases biopsy of enlarged superficial lymph nodes has shown histological changes of sarcoidosis.

## PROGNOSIS OF THE LUNG LESIONS

In the early stage of diffuse pulmonary infiltration it is impossible to give an accurate forecast of the probable course in the individual case. It certainly depends to some extent upon the patient's race, in Negroes a progressive course possibly culminating in frank tuberculosis is more frequent than in Europeans. A survey of published figures suggests that where intra-thoracic manifestations are a leading feature, regression of the lesions amounting to complete clearing may occur in about half, sometimes after the lesions have been apparently stationary for a considerable time. Of the remaining half some will show very little change after relatively prolonged periods of observation, but some will show increase in the extent and severity of the lesion with eventual development of fibrosis and secondary emphysema, and some of this last group will eventually produce tubercle bacilli in the sputum. Death in such cases may occur after prolonged illness from right heart failure or from intercurrent respiratory infection. It is obvious that if a patient first comes under observation with the late fibrotic and emphysematous changes, no permanent improvement can be expected.

## TREATMENT

No form of treatment can be relied upon to influence the course of the disease. In the more acute types approximating to chronic miliary tuberculosis it may appear reasonable to try the effect of streptomycin. However, streptomycin is most effective in the acute exudative types of tuberculosis and relatively little, if at all, effective in the most productive types, and the lesions of sarcoidosis are the perfect example of the productive non exudative reaction. The relatively little information at present available about the effect of streptomycin in sarcoidosis is in accordance with expectation based on these considerations. Calciferol has been used in an attempt to modify the course of some of the progressive types of nodular infiltrations. Since in these forms of pulmonary sarcoidosis spontaneous regression is frequent results are difficult to interpret, and evidence is as yet inconclusive about the value of this form of treatment. Many patients with sarcoidosis are peculiarly susceptible to the toxic effects of calciferol and may suffer serious biochemical and symptomatic disturbances with evidence of renal damage after the uncontrolled administration of quite small doses. At present, therefore calciferol should not be given to patients who are symptom free and whose lesions do not seem to be progressive, its use may possibly be justified in those patients who have diffuse pulmonary infiltrations which are increasing in density under observation. In such cases, after preliminary investigation of the levels of serum calcium and blood urea and of renal function, 50 000 units daily by mouth should be given at first. This may be increased progressively if there is neither symptomatic nor serious biochemical disturbance up to a maximum of 150 000 units daily and the tolerated dosage maintained for 4 months. Minor upsets during the course of treatment may necessitate adjustment of dosage, and some patients cannot tolerate even 50 000 units daily. Radiographic improvement usually is not evident until after the end of this course of treatment.

For the later stages of pulmonary sarcoidosis with permanent fibrotic and emphysematous changes no curative treatment is possible, and the treatment is similar to that of a case of pulmonary emphysema and fibrosis of any other cause.

## BIBLIOGRAPHY

- Longcope W T (1941) "Sarcoidosis or Besnier Boeck Schaumann Disease" *J Amer med Ass* 117, 1321
- Pinner M (1919) "Non-caseating Tuberculosis" *Amer Rev Tuberc* 37, 690
- Scadding J G (1950) "Sarcoidosis with Special Reference to the Lung Changes" *Brit med J* 1, 745
- Schaumann J (1936) "Lymphogranulomatosis Benigna in the Light of Prolonged Clinical Observations and Autopsy Findings" *Brit J Derm* 48, 399
- Snapper I and Pompen A W N (1938) *Pseudotuberculosis in Man* Haarlem

*Resultant or associated pulmonary disease*—Broncho-pneumonia is usual in the terminal stages of all reticuloses. Pulmonary tuberculosis is not infrequently seen in association with chronic leukaemia or lymphadenoma and difficulty may be experienced in deciding which disease came first. Pulmonary cavitation occurring during the course of a reticulosis may be due to tuberculosis, to a lung abscess or, occasionally, to the breakdown of a mass of lymphadenomatous glands.

### LYMPHADENOMA AND LYMPHOSARCOMA

In lymphadenoma, abnormal radiological changes are seen in rather more than 50 per cent of the chest radiographs. They consist of enlarged lymphatic glands or pleural effusion. In rare cases the disease may be infiltrated in one or other of the ways described above (Peirce, Jacob and Hildreth, 1936, Gail and Mallory, 1942). The rate of progression is slow and the results of radiotherapy, other than for enlarged mediastinal glands, is disappointing. A sudden deterioration of health, with rapid increase in the lesions, suggests sarcomatous degeneration (see Figs 75-76).

In lymphosarcoma the radiographic changes are similar to lymphadenoma but occur in only about one case in five. They progress rapidly and usually show a favourable initial response to radiotherapy (Fig 77).

### MYCOSIS FUNGOIDES

The disease may be limited to the skin for many years and then rapidly extend to other parts of the body, the lungs may show large roundish masses or a coarse mottling, as a result of haematogenous dissemination (see Fig 78).

### THE LEUKAEMIAS

#### Acute leukaemia

In the acute leukaemias, which run a febrile course and terminate fatally within a few months, pulmonary infarcts and venous thromboses are common, an aspiration pneumonia, perhaps with an acute lung abscess, is likely to occur particularly in those patients, comprising about one-third of the total number, who have an associated acute ulcerative infection of the mouth or throat. The respiratory symptoms of these complications are often surprisingly slight and may be overlooked. In children with acute lymphatic leukaemia massive mediastinal glandular enlargement may develop (see Figs 79-80).

#### Chronic leukaemia

In the chronic leukaemias about one case in five shows an abnormal radiograph of the chest, in chronic lymphatic leukaemia the changes are indistinguishable from those of lymphadenoma, enlarged mediastinal glands being the commonest feature (see Fig 81), in chronic myeloid leukaemia the radiological signs are usually slight and inconclusive (see Kirklin and Hefke, 1931, Falconer and Leonard, 1938).

### OTHER DISEASES OF THE RETICULO ENDOTHELIAL SYSTEM

*Gaucher's disease*—This rarely extends to the lungs. Merklen, Wartz and Watter (1933) were able to recover Gaucher's cells from the sputum of an advanced

FIG. 75 — Lymphadenoma with enlarged mediastinal glands and rounded deposits in the lungs




FIG. 76 — Lymphadenoma with extensive involvement of the lungs and no radiological evidence of enlarged mediastinal glands. The invasion had been progressive during the previous year and failed to respond to radiotherapy



FIG 77 — Lymphosarcoma with massive glandular enlargement. Following radiotherapy the mass disappeared radiologically to recur 3 years later.

FIG 78 — Mycosis fungoides. Section of the whole of the right lung shows many deposits. An extensive eruption of the skin had been present for 6 years and a radiograph of the chest had been normal 3 months before death.



FIG. 79 — Acute lymphatic leukaemia in a boy aged 13 years showing massive mediastinal glandular enlargement. He died after an illness of two months.



FIG. 80 — A lateral view of the case shown in Fig. 79. The characteristic enlargement of the glands into the anterior mediastinum is seen.







FIG 81 — Chronic lymphatic leukaemia with enlarged mediastinal glands and a left pleural effusion

case, and Myers (1937) found considerable involvement of the lungs and mediastinal glands in another

*Niemann Pick's disease* —In this disease the lungs may be involved in the general process of congestion and oedema and the alveoli may contain many of the characteristic large pale cells (Canmann, 1944)

*Other syndromes* —Hand Schüller Christian disease Letterer Siwe's disease and eosinophilic granuloma of bone are described separately under the heading of Honeycomb Lungs

### LEUKAEMOID REACTIONS IN PULMONARY DISEASE

Leukaemoid reactions are occasionally seen in diseases of the lungs. They consist in a leukocytosis with immature cells and need to be differentiated from leukaemia. Forkner (1938) states that in leukaemia there is almost invariably a thrombocytopenia and that the immature forms are of a uniform type and comprise at least 70 per cent of the white blood cells. Neither of these features is seen in a leukaemoid reaction. In whooping cough white cell counts of 100 000 per cubic millimetre and more have been recorded usually in cases complicated by broncho pneumonia, nearly all the cells being lymphocytes. Pneumonia in children may provoke counts of 50 000–100 000 white cells per cubic millimetre, often including many immature forms, the counts rapidly return to normal during convalescence.

*The reaction in miliary tuberculosis* —A leukaemoid reaction is fairly common in miliary tuberculosis and should be suspected when there is enlargement of the liver,

spleen or lymph glands, the abnormal blood cells are nearly always of the myelogenous series. At autopsy the affected organs show miliary tubercles but none of the changes typical of a true leukaemia. The reaction is rare in the more chronic forms of tuberculosis.

## BIBLIOGRAPHY

## BIBLIOGRAPHY

- Canman M F (1944) Niemann Pick Disease (Acute Idiopathic Xanthomatosis, Phosphatid Lipoidosis Lipoid Cell Splenomegaly Nemann Type) *J Pediat*, 24, 335
- Falconer E H and Leonard M E (1938) "Pulmonary Involvement in Lymphosarcoma and Lymphatic Leukaemia" *Amer J Med Sci* 195 294
- Fieissinger N and Fauvet J (1941) "Le Poumon leucémique" *Pr Méd* 49, 449
- Forkner C E (1938) *Leukaemia and Allied Disorders* New York
- Gall E A and Mallory T B (1942) Malignant Lymphoma Clinico-pathologic Survey of 618 Cases *Amer J Path* 18 381
- Joachim H and Loewe L (1927) "Atypical Acute Myeloid Leukaemia with Unusual Pulmonary Manifestations Case Report with necropsy findings" *Amer J Med Sci.* 174 215
- Kirkin B R and Hefke H W (1931) Roentgenologic Study of Intrathoracic Lymphoblastoma" *Amer J Roentgenol* 26 681
- Merklen P Waitz R and Watter J (1933) Maladie de Gaucher diagnostiquée par la ponction de la rate lésions vertébrales étendues présence de cellules de Gaucher dans l'expectoration *Ann Med* 33, 97
- Myers B (1937) Gaucher's Disease of the Lungs" *Brit med J* 1, 1029
- Oswald N (1950) Pulmonary Changes in the Reticuloses *Proc R Soc Med.* 43, 208
- Pearce C B Jacob H W and Hildreth R C (1936) "Roentgenologic Considerations of Lymphoblastoma I Roentgen pulmonary pathology of the Hodgkins type" *Amer J Roentgenol* 36 145

## CHAPTER 15

### HONEYCOMB LUNGS

NEVILLE OSWALD

#### AETIOLOGY

AMONGST the many varieties of pulmonary cysts, instances are occasionally seen of lungs which on cross section resemble a honeycomb or sponge, as a result of cysts uniformly distributed throughout both lungs and measuring up to about 1 centimetre in diameter. The cysts may be due to a *primary pulmonary disease* or be part of a general medical disorder, in either case the clinical features are similar, spontaneous pneumothoraces being common and right heart failure almost invariably the cause of death. The condition has been reviewed in an article by Oswald and Parkinson (1949).

*Pathogenesis*—There is no doubt that in a minority of instances a developmental defect is the causative factor. It is equally certain that diffusely infiltrating

occasionally be the cause

#### DEVELOPMENTAL GROUP

*The developmental group is in many ways the most typical and the most easily* . . . . . The . . . . . walled, tend to be rather large and give . . . . . Is in otherwise translucent lung fields . . . . . in childhood or early adult life; they may be recurrent or bilateral and are sometimes fatal. Other causes of death include right heart failure, broncho-pneumonia and associated developmental defects. Few patients live for more than 30 years. Congenital heart disease is a not infrequent concomitant (see Figs 82-83).

*The mode of production of the cysts is unknown.* Pulmonary cysts of any kind are excessively rare in still-born foetuses and for this reason, apart from any others, the term "congenital cystic disease" should be abandoned. Doubtless there is a developmental factor, which may not necessarily operate until some months or years after birth. The problem as it presents is to account, often, for a child with well developed cysts and no previous history of respiratory disease, in the light of the knowledge that such cysts have hardly ever been observed at birth. In the present state of our knowledge it is impossible to state whether the cysts are truly congenital or whether they have been caused by a trivial respiratory infection after birth in lungs of poor quality. The characteristically thin walls of the cysts, the paucity of reaction in the connective tissues of . . . . . developmental abnormalities elsewhere . . . . . entity and to justify their being label . . . . .




FIG. 82 — Developmental type showing fine linear striations through both lungs. A man aged 23 years who had several spontaneous pneumothoraces from the last of which he suddenly died.

## TUBEROUS SCLEROSIS

### Classification

Tuberous sclerosis, in its classical form, is seen as a familial disease of children, mental deficiency, epilepsy and adenoma sebaceum of the face being its commonest manifestations. There may in addition be myomas of the heart and kidneys, sub-ungual fibromas and tumours of the retinae. Such children rarely survive until adult life and none have so far shown any pulmonary lesions. In recent years some half-a-dozen adults, all females and some with a family history of tuberous sclerosis, have been found to have a diffuse leiomyomatous infiltration of both lungs with honeycomb cysts (see Fig 84), none has had mental deficiency or epilepsy, but all except one have had adenoma sebaceum and sub-ungual fibromas and two showed the typical involvement of the cerebral cortex and kidneys at autopsy.

There is thus a small group of patients with somewhat atypical tuberous sclerosis whose symptoms are predominantly respiratory. Closely related to these are others with identical pulmonary changes and with evidence of extra-pulmonary mesodermal dysplasia, but with none of the characteristic clinical signs of tuberous sclerosis. They would appear to belong to a range of diseases having tuberous sclerosis at one end and the angiomatoses of Sturge-Weber and Hippel-Lindau at the other, they may conveniently be regarded as cases of multifocal mesodermal dysplasia. A third group comprises cases with a vast increase of muscular tissue in the lungs and with honeycomb formation, but with no extra-pulmonary lesions.

# HONEYCOMB LUNGS




Fig. 1 (a)

(b)

In the past these have been called muscular cirrhosis of the lungs but it is probable that they belong to this group of mesodermal defects and represent a purely pulmonary form.

The pathological process in the whole group is similar and consists of smooth muscle growing out from the smaller bronchi and blood vessels often forming irregular masses in the lungs or within the lumina of the bronchi and sometimes directly infiltrating the alveolar walls. The mechanism of production of the cysts

is probably by obstruction or strangulation of the smaller radicles of the bronchial tree

### Clinical picture and diagnosis

The symptoms derive from spontaneous pneumothoraces and right heart failure, which rarely occur before adult life. The radiological appearances are of two kinds. Usually a diffuse reticulation appears throughout both lungs, which increases in density over a period of years, the cysts may be difficult to see in the early stages, but they become more apparent as the general density of the lungs increases. In the other variety an infiltrating process starts in one or both lungs and gradually spreads, making the differential diagnosis from pulmonary tuberculosis extremely difficult, unless there are extra pulmonary manifestations of the disease or a piece of lung is removed for microscopical examination (Fig. 85).

## XANTHOMATOSES

### Classification

Amongst the xanthomatoses there are three varieties which are closely related and often present in a mixed form, the lungs may be involved in any or in any combination of them and show a diffuse granuloma with honeycombing. The three varieties comprise (1) Letterer Siwe's disease, (2) Hand Schüller Christian disease, and (3) eosinophilic granuloma of bone.

Letterer Siwe's disease is typically an acute aleukaemic reticulosis of infants, having a non lipid-containing mononuclear as its characteristic cell, and running an acute febrile course of a few weeks or months, with enlarged liver, spleen and lymph glands, and terminating fatally.

Hand Schüller Christian disease tends to develop in early childhood. It frequently shows its triad of exophthalmos, diabetes insipidus and erosions of the skull, as a result of invasion by cholesterol-containing mononuclear cells (Fig. 86).

Eosinophilic granuloma may occur in childhood or adult life, and is diagnosed by the finding of cystic spaces, usually in the shaft of a long bone, filled with eosinophil cells.

Microscopical examination of sections taken in any of these diseases may show that the lungs have become diffusely infiltrated with a granuloma. The characteristic cells may, however, be sparse or difficult to see because of secondary infection. There may or may not be honeycombing of the lungs, and it seems extremely probable that the cyst formation represents a late phase.

### Clinical features

The respiratory symptoms depend largely upon the age of the patient and the clinical course of the extra pulmonary features. In infants and young children, bronchitis and broncho-pneumonia are common and although the immediate cause of death may be right heart failure, prolonged fever and inanition are major contributory factors.

In a certain number of cases, the xanthomatosis appears to become arrested, and many years later evidence of pulmonary or pituitary insufficiency may be apparent (Oechsli and Miles, 1934). One such case, a man in the late twenties, has been under observation for 5 years with well marked honeycomb lungs which, apart

# HONEYCOMB LUNGS





Fig 84 — (a) Tuberosclerosis with diffuse mottling of lungs and a honeycomb appearance becoming visible in the mid zones in a woman aged 31 years. She also has adenoma sebaceum of the face, subungual fibromas of the toes and a filling defect in the left kidney which probably represents a myoma. During 4 years' observation she has continued to lead a busy life as a shop assistant troubled only by slight breathlessness. (b) Higher magnification showing myomatous infiltration of the lung.

(a)



(b)

FIG. 85 — (a) Irregular type of infiltration which may be seen in tuberculous sclerosis in a woman aged 36 years. A spontaneous pneumothorax had occurred 5 years previously and the infiltration starting in left upper zone was seen to spread radiologically during 3 years. She also had adenoma sebaceum of the face. The diagnosis of leiomyomatous infiltration with honeycomb formation was confirmed by lung biopsy. (b) Higher magnification



(a)



(b)



# HONEYCOMB LUNGS




FIG 86 — (a) Hand Schuller Christian's disease in a baby aged 9 months. The skull and spine were also affected. The wide upper mediastinal shadow represents an enlarged thymus gland confirmed at autopsy. There was considerable secondary infection of the lungs at the time of the radiograph. (b) section of lung showing the cystic spaces



(b)

from a patient with

insipidus for 8 years and his honeycomb lungs have been radiologically unchanged for 4 years, in recent months progressive pituitary failure has developed. The combination of honeycomb lungs and diabetes insipidus (see Fig 89), of which there are some half-dozen examples in the literature, is almost certain evidence of a former xanthomatosis.

The radiological changes in infancy consist in a mixture of rather ill defined cysts, reticulation, and signs of pulmonary congestion and broncho pneumonia, the

cysts when present are usually sufficiently obvious to enable a diagnosis to be made. In adults the density of the reticulation varies within very wide limits; in some cases the connective-tissue reaction is so slight that the appearances are very similar to those seen in the developmental type of honeycomb lungs, in others the reticulation is gross and coarse. The cysts can usually be seen without difficulty especially during the phases of spontaneous pneumothorax.

### BILIARY CIRRHOSIS OF INFANTS

Biliary cirrhosis of infants, a condition of which little is known, consists in a granulomatous infiltration of the biliary tracts, which gives rise to jaundice and is usually fatal; the pulmonary changes are indistinguishable from those of xanthomatosis.

### HONEYCOMB LUNGS OF UNCERTAIN AETIOLOGY

In addition to the above examples of honeycomb lungs occurring as part of various general medical disorders, there are others, numerically representing about 50 per cent of the total, in which the disease is limited to the lungs and the aetiology of which is by no means certain. The group comprises adults, mostly males, ranging in age from 20 to 60 years, with spontaneous pneumothoraces and right heart failure as the outstanding clinical features of the disease and having diffuse radiological changes similar (see Fig 90) to those described above as occurring in adults. It is likely that there are several different aetiological factors within the group, but, whatever their nature, they must affect both lungs equally. Pathologically, the features are somewhat similar in all, with varying degrees of inflammation and fibrosis of the connective tissue, and universal cyst formation. The following possible aetiological factors should be considered.

#### Diffuse cystic granuloma of the lungs

the cysts arise by a process of strangulation of the smaller branches of the bronchial tree, and that their walls become infiltrated by the same process. They are uncertain of the aetiology of the condition, but it is likely that

xanthomatosis lies in the paucity of the characteristic cells, largely as a result of secondary infection. As things stand, in those cases in which there is a diffuse

#### Emphysema

Whether or not universal emphysema should be regarded as a variety of honeycomb lungs is a matter of opinion. Emphysema normally derives from sustained





FIG. 86 — (a) Hand Schuller Christian's disease in a baby aged 9 months. The skull and spine were also affected. The wide upper mediastinal shadow represents an enlarged thymus gland confirmed at autopsy. There was considerable secondary infection of the lungs at the time of the radiograph. (b) section of lung showing the cystic spaces.



(b)

from passing through a phase of spontaneous pneumothoraces, have remained unchanged radiologically, in his case severe diabetes insipidus has recently developed (Fig 87). Another man (see Fig 88) in his fifties, had a pathological fracture of his femur as a result of an eosinophilic granuloma 10 years ago, he has had diabetes insipidus for 8 years and his honeycomb lungs have been radiologically unchanged for 4 years, in recent months progressive pituitary failure has developed. The combination of honeycomb lungs and diabetes insipidus (see Fig 89) of which there are some half dozen examples in the literature, is almost certain evidence of a former xanthomatosis.

The radiological changes in infancy consist in a mixture of rather ill defined cysts, reticulation, and signs of pulmonary congestion and broncho pneumonia, the

## HONEYCOMB LUNGS OF UNCERTAIN AETIOLOGY

cysts when present are usually sufficiently obvious to enable a diagnosis to be made. In adults the density of the reticulation varies within very wide limits, in some cases the connective-tissue reaction is so slight that the appearances are very similar to those seen in the developmental type of honeycomb lungs, in others the reticulation is gross and coarse. The cysts can usually be seen without difficulty especially during the phases of spontaneous pneumothorax.

## BILIARY CIRRHOSIS OF INFANTS

Biliary cirrhosis of infants, a condition of which little is known, consists in a granulomatous infiltration of the biliary tracts, which gives rise to jaundice and is usually fatal; the pulmonary changes are indistinguishable from those of xanthomatosis.

## HONEYCOMB LUNGS OF UNCERTAIN AETIOLOGY

In addition to the above examples of honeycomb lungs occurring as part of various general medical disorders, there are others, numerically representing about 50 per cent of the total, in which the disease is limited to the lungs and the aetiology of which is by no means certain. The group comprises adults, mostly males, ranging in age from 20 to 60 years, with spontaneous pneumothoraces and right heart failure as the outstanding clinical features of the disease and having diffuse radiological changes similar (see Fig 90) to those described above as occurring in adults. It is likely that there are several different aetiological factors within the group, but, whatever their nature, they must affect both lungs equally. Pathologically, the features are somewhat similar in all, with varying degrees of inflammation and fibrosis of the connective tissue, and universal cyst formation. The following possible aetiological factors should be considered.

### Diffuse cystic granuloma of the lungs

In a recent pathological study, Cunningham and Parkinson (1950) have observed gradations in the amount of granulation tissue and fibrosis, which would account well for the range of densities seen in radiographs of the chest. They consider that the cysts arise by a process of strangulation of the smaller branches of the bronchial tree, and that their walls become infiltrated by the same process. They are uncertain of the aetiology of the condition, to which they apply the rather non-committal term, diffuse cystic granuloma of the lungs.

There is no doubt that the changes are very similar to, and may be indistinguishable from, those seen in xanthomatosis—a fact which raises the question of the difference between this condition and the "eosinophilic xanthomatous granuloma" of Thannhauser (1947). The difficulty in the diagnosis of solitary pulmonary xanthomatosis lies in the paucity of the characteristic cells, largely as a result of secondary infection. As things stand, in those cases in which there is a diffuse granuloma of the lungs without evidence of a xanthoma elsewhere in the body a non-committal term such as diffuse cystic granuloma should be used, in order to permit of an alternative aetiology to xanthomatosis, if necessary.

### Emphysema

Whether or not universal emphysema should be regarded as a variety of honeycomb lungs is a matter of opinion. Emphysema normally derives from sustained




FIG 88 — (a) Coarse mottling of the lungs with a honeycomb appearance towards the bases, in a man aged 57 years who also has an eosinophilic granuloma of his femur and diabetes insipidus. Radiotherapy to the lungs caused an increase in breathlessness. Radiographs of his chest have remained unchanged over 6 years, but he now suffers from progressive pituitary failure. (b) Higher magnification

(b)



FIG 89 — (a) Diffuse cystic granuloma associated with diabetes insipidus and bilateral spontaneous pneumothoraces in a man aged 38 years. The heavy coarseness would appear to represent a late and chronic phase of the disease. He died from heart failure a year later. Autopsy was not performed (b) higher magnification

(a)



211

(b)




FIG. 90 — (a) Diffuse cystic granuloma with right spontaneous pneumothorax in a man aged 20 years. He died from right heart failure 10 years later autopsy revealing gross granulomatous invasion of the lungs with many cysts (b) higher magnification



(b)

# CLINICAL FEATURES OF HONEYCOMB LUNG





FIG 91 — (a) A chronic bronchitic patient aged 55 years with heart failure (b) section of whole lung showing the universal distribution of the cysts the other lung was similar



## HONEYCOMB LUNGS




FIG 92—Diffuse cystic granuloma in a pituitary dwarf the bronchogram showing that the iodized oil does not enter the cysts

Spontaneous pneumothorax develops in about 50 per cent of all patients with honeycomb lungs

Right heart failure is the usual cause of death in this almost uniformly fatal disease but it may be delayed until late middle age, it is due to a progressive diminution in effective alveolar surface the end frequently being hastened by a secondary respiratory infection

A thorough search should always be made for extra pulmonary features which might lead to the diagnosis of a general disorder having honeycomb lungs as one of its manifestations

### Radiological appearances

These differ somewhat in each group and therefore are described separately above In general the lung fields are uniformly marked by cysts and reticulation the latter showing wide variations in density Tomograms are often helpful in demonstrating the cysts On bronchography iodized oil never enters the cysts and occasionally the smaller bronchi appear to be encircling them (see Fig 92) The diagnosis is often easy from a single radiograph especially during a spontaneous pneumothorax when the cysts stand out clearly In the presence of heart failure or respiratory infection the widespread distribution of the cysts may not be demonstrable there is thus a group of patients in whom honeycomb lungs may be suspected radiologically but to whom the clinical label cannot be applied without pathological proof

## TREATMENT

### Differential diagnosis

The differential diagnosis of diffuse pulmonary reticulation is formidable, it includes pneumoconiosis, sarcoidosis, chronic miliary tuberculosis, lymphangitis carcinomatosa, and other rare diseases, in none of which have honeycomb lungs ever been found

## TREATMENT

Aside from the treatment of such general disorders as may be present, the successful management of spontaneous pneumothoraces is often a life-saving measure.

## BIBLIOGRAPHY

- Cunningham, G J, and Parkinson, T (1950) "Diffuse Cystic Lungs of Granulomatous Origin" *Thorax*, 5, 43
- Oechli W R and Miles S H (1934) "Simultaneous Bilateral Spontaneous Pneumothorax. Review of literature and report of case due to congenital cysts of lungs" *Amer Rev Tuberc*, 3 239
- Oswald, N and Parkinson T (1949) "Honeycomb Lungs" *Quart J Med.*, 18, 1
- Thannhauser S J (1947) "Eosinophilic Granuloma of Bone Synonymous with Schüller-Christian Disease. Lipid Granuloma, Essential Xanthomatosis of Normocholesteremic Type and Eosinophilic Xanthomatous Granuloma" *Arch intern Med*, 80, 283

## CHAPTER 16

### CARDIO-PULMONARY SCHISTOSOMIASIS

D EVAN BEDFORD

*Synonym* Bilharzial heart disease

PULMONARY schistosomiasis with its cardiac complications may be due to either *Schistosoma haematobium* or *S. mansoni*, and is always secondary to urinary or intestinal infestation. Pulmonary lesions were found in 33 per cent of 282 cases of schistosomiasis examined at necropsy at the Kasr-el-Aini Hospital, Cairo (Shaw and Ghareeb, 1938), and they have also been reported from other countries where bilharzial infestation is prevalent, namely North and South Africa, South America and the West Indies. Our knowledge of cardio pulmonary schistosomiasis is based mainly on the disease as encountered in Egypt, where Bilharz first discovered the parasite, where its pathology has been specially investigated by Sorour (1932) and by Shaw and Ghareeb (1938), and where its clinical manifestations were first recognized by Azmy (1932), Mainzer (1935) and

in Porto Rico

#### PATHOLOGY

Pulmonary lesions may be caused either by the worms themselves or by the ova, the latter being by far the more important cause

##### **Invasion by ova**

Ova derived from the systemic or mesenteric veins pass through the right heart to reach the lungs as emboli, lodging in the arterioles which accompany the respiratory bronchioles. In the case of *S. haematobium*, the ova are derived from the vesical, ureteric, prostatic and uterine venous plexuses and pass via the internal iliac veins to the inferior vena cava, in the case of *S. mansoni*, the ova are derived from the portal venous system and reach the vena cava via dilated porto caval anastomoses due to cirrhosis of the liver (Egyptian hepato-splenomegaly).

The ovum, once impacted in a pulmonary arteriole, produces a necrotizing arteriolitis in which the continuity of the media is destroyed, allowing the ovum to penetrate the vessel wall, outside which is formed a "bilharzial tubercle", not unlike the tubercle of Koch's infection. In the early stages there is a cellular reaction around the ovum, consisting of eosinophil leucocytes, endothelioid cells, giant cells, and later lymphocytes. This cellular reaction subsides, leaving a nodular scar containing the calcified or disintegrated ovum. The damaged arteriole is occluded by intimal thickening without thrombosis, and the occluding tissue becomes recanalized. Owing to focal destruction of the media, the newly



FIG. 93.—Section showing thickened pulmonary arteriole with adjacent nodule containing a giant cell filling the shell of a bilharzia ovum (From Bedford A. Darav and G. G. S. 1946 by courtesy of Brit. Heart J.)

In the course of time repetitive embolization leads to widespread arterial obliteration sufficient to cause pulmonary hypertension with consequent hypertrophy of the right ventricle and gross dilatation of the pulmonary arterial tree. Both the main pulmonary stem and its hilar branches may reach aneurysmal size and their walls become thickened, atheromatous, calcified or even thrombosed. Bilharzial tubercles have been found in the ventricular myocardium in a case of *S. japonicum* infection (Strong 1944) and Kenawy has reported focal cellular infiltration of the myocardium but ova were not identified.

Thus the pulmonary lesion is essentially arterial and the lung parenchyma is little affected apart from the perivascular tubercles. A broncho-pulmonary form has been recognized radiologically and evidently occurs in the earlier and more active phase of the disease when cellular reactions around the ova are present. Sorour (1932) has described an endobronchitis obliterans of the larger and medium sized bronchi but gross fibrosis of the lungs does not occur.

#### Invasion by worms

Sometimes the worms themselves—either the male or coupled male and female—reach the lungs as emboli where they cause arterial thrombosis and a focal verminous pneumonia, the foci of consolidation being 1–5 millimetres in diameter. Day (1937) has reported finding pulmonary arteries packed with coupled worms.



FIG 94 — Drawing of section showing occluded pulmonary arteriole with revascularization of lumen surrounded by angiomatoid formation characteristic of bilharzial pulmonary endarteritis. From same case as Fig 96 (From Bedford, Aldaras and Girgis, 1946 by courtesy of *Brit Heart J*)

When the inflammatory reaction subsides, the dead worm becomes calcified and enclosed in a fibrous capsule

## CLINICAL FEATURES

### Incidence

In Egypt, cardio-pulmonary bilharziasis is encountered almost entirely amongst the agricultural workers inhabiting the Nile Delta, the majority of whom are infested with bilharzia. The pulmonary lesion is commonest in males aged 12-35 years, and is always associated with either urinary or intestinal bilharziasis, in the latter case hepato-splenomegaly is usually present

### Pulmonary disease

Before the heart is seriously involved, asthmatic or bronchitic symptoms may lead to radiography of the chest, which shows a military mottling of the lung fields, not unlike that of military tuberculosis or pulmonary haemosiderosis, as described by Mainzer (1935 and 1938) and by Erfan and his colleagues (1949). The pulmonary mottling is due to dilated arterioles beaded by nodules about 1

## CLINICAL FEATURES



FIG 95 — Rad ograph from a case of pulmonary schistosomiasis in active stage showing mottled lung fields due to beaded and dilated vessels. There is some dilatation of the main pulmonary artery and its hilar branches. (From Erfan and his colleagues 1949 by courtesy of Trans R Soc trop Med Hyg.)

millimetre in diameter (Fig 95) these are due presumably, to bilharzial tubercles. The lung changes may be evident before the heart is much involved or may be associated with enlargement of the heart and main pulmonary vessels. Emphysema, with basal bronchitic or catarrhal signs, is found in some cases, but more often clinical signs in the lungs are absent. An acute focal reaction in the lungs with increased clinical or radiological signs may result from antimony treatment.

### Bilharzial heart disease

In the cardiac phase of the disease the clinical picture is that of pulmonary hypertension without gross lung disease. The apex is displaced to the left and systolic pulsation is palpable over the pulmonary artery and conus. A pulmonary systolic murmur and accentuated second sound are audible and sometimes there is a diastolic Graham Steell murmur due to relative pulmonary incompetence as in Azmy's original case (Azmy 1932). The radiograph shows enlargement of the right ventricle towards the left, and dilatation of the pulmonary trunk and its hilar branches which may pulsate excessively. Aneurysmal dilatation of the pulmonary artery is not uncommon (Fig 96). In the writer's case the aorta was hypoplastic as was also noticed by Erfan.

Cyanosis is usually absent until heart failure appears and as is usual in primary pulmonary hypertension the arterial oxygen saturation is normal or slightly reduced for example 82-95 per cent in the cases described by Soliman (1950).



## TREATMENT

to be distinguished (1) from other forms of pulmonary heart disease, (2) from mitral stenosis with haemosiderosis of the lungs and (3) from congenital heart disease, especially auricular septal defect which may give an identical radiological picture. Mitral stenosis is excluded by the absence of left auricular enlargement as shown by radioscopy in the right oblique position after a barium swallow, and by the absence of mitral diastolic or presystolic murmurs after a test exercise. An auricular septal defect can be excluded by cardiac catheterization which in bilharzial heart disease demonstrates the absence of an auricular shunt and in more advanced cases shows pulmonary hypertension (Soliman 1950 Barbato and his colleagues 1951).

## TREATMENT

The aim of treatment is preventive and urinary or intestinal infections should be treated with a full course of antimony compounds such as tartar emetic or Foudain (stibophen) with a view to preventing pulmonary embolization. In the early stages of lung involvement antimony may be tried with caution as a means of eradicating the primary visceral infection and preventing further ova from reaching the lungs but there is danger of an acute pulmonary reaction of an allergic nature. Erfan give antimony to 44 cases of schistosomiasis with pulmonary lesions and 4 of the patients with bronchitic or asthmatic symptoms improved but in other cases fever urticaria acute pulmonary congestion and eosinophilia occurred and 1 patient died suddenly (Erfan and his colleagues 1949). Caution is therefore necessary. Heart failure is an absolute contra indication to antimony treatment because besides the danger of lung reactions it has a direct toxic action on the myocardium. Congestive failure requires the usual measures such as rest in bed digitalis mercuril diuretics and a low sodium diet. Once failure has occurred the expectation of life is relatively short.

## BIBLIOGRAPHY

- Vim S (1937) Pulmonary Arteriosclerosis of Bilharzial Nature *J Egypt med A* 15 87  
 Barbato F Lma E P Merino G Cotrim E and Dantas O M (1951) Apectos  
 patológicos da Arteria Pulmonar na Esquistossomose Manson *Arq Brasl Cardl* 1  
 4 233  
 Bullard D F Adaras S M and Grigs B (1947) Bilharzial Heart Disease in Egypt  
 Cor Pulmonale due to Bilharzial Pulmonary Endarteritis *Br J Heart J* 8, 87  
 Clark F and Graef I (1935) Chronic Pulmonary Arteritis in Schistosomiasis Manson  
 associated with Right Ventricular Hypertrophy Report of a case *Amer J Path* 11 693  
 Day H B (1917) Pulmonary Bilharziasis *Trans Soc trop Med H e* 30 575  
 Erfan M Erfan H Mousa A M and Deeb A A (1949) Chronic Pulmonary Schisto-  
 soma Aetiological and radiological study *Trans Soc trop Med H e* 42 477  
 Kenawy M R (1950) The Syndrome of Cardopulmonary Schistosomiasis (Cor Pulmonale)  
*Amer Heart J* 39 64  
 Mansour I (1935) Sur la bilharziose pulmonaire maladie des poumons stimulante tuber-  
 culose *Arq Brasl Cardl* 1 85 538  
 Mansour I (1935) Clinical Aspects of Pulmonary Diseases Induced by Schistosoma Haematobium  
 and Manson *J Egypt med A* 21 762  
 Shaw S I B and Ghareeb A A (1938) "The Pathogenesis of Pulmonary Schistosomiasis in  
 Egypt with Special Reference to Azygos Disease" *J Path Bact* 46, 401  
 Soliman H S (1950) Cardiac Catheterization in Pulmonary Bilharziasis Thesis for the  
 M.D. Cairo



# CARDIO PULMONARY SCHISTOSOMIASIS

- Sorour, M F (1932) "The Pathology and Morbid Histology of Bilharzial Lesions in various Parts of the Body" *C R Congrès International de Médecine Tropicale et d'Hygiène, Cairo*, 1928 Vol 4, p 321
- Strong R P (1939) In *Stitt's Tropical Diseases*, 6th ed., Vol II, 1429 Case Report of Africa and Santa Cruz



Dje-di (1949) have recorded empyema developing as a result of pleural effusion. There is often an inflammatory reaction in the lung when the patient will have all the symptoms of bronchopneumonia. Physical examination will only reveal scattered râles throughout the lungs, until there are the signs of pleurisy and pneumonia. Diarrhoea may result from abdominal involvement, and Jacksonian epilepsy, headaches, and hemiplegia from cerebral involvement. Sometimes there is also enlargement of the lymph glands and ulceration of the skin.

### Diagnosis

Diagnosis is established by finding ova in the sputum, by the addition of 0.1 per cent sulphuric acid to the sputum the procedure is rendered more simple. Charcot-Leyden crystals are often found in the sputum. There is usually a well marked eosinophilia. The complement fixation test can be helpful and radiographic examination will show the characteristic cystic changes scattered throughout the lungs.

The disease runs a chronic course, with periodic attacks of haemoptysis, but the ultimate prognosis is always unsatisfactory owing to the difficulty of eliminating the parasites. Where the brain is involved the disease is always fatal.

### Treatment

Infection can be prevented by not eating crabs in those parts of the world where the disease is endemic. All the water in these areas should be boiled and filtered, and bathing should be avoided in contaminated rivers. Potassium iodide, 20 grains, three times a day is said to produce beneficial results and improvement has occurred after the intra-tracheal injection of Lipiodol. Emetine and tartar emetic help to relieve the cough and sputum. Protosil has proved of value in some cases. However a 15-day course of 40 millilitres (equivalent to 340 milligrams antimony) per day of Fouadin is probably the most effective treatment.

## ROUND-WORMS INFECTION

Round worms may be divided into the intestinal worms, and the tissue worms. The intestinal worms include *Strongyloides*, *Ankylostoma*, *Necator*, *Ascaris* and *Oxyuris*. In the first three of these nematodes the larvae penetrate the skin from water or damp earth and find their way into blood vessels, through which they are carried to the right side of the heart and thence to the lungs. In the case of *Ascaris* and *Oxyuris* the ova are swallowed in water or food, pass into the intestines where the larvae penetrate the bowel wall and migrate to the lungs. From the lungs both groups progress through the trachea, oesophagus and stomach to their natural habitat in the intestine.

In their passage through the lungs all these larval worms are capable of blocking the pulmonary arteries and arterioles, and thus causing small pulmonary infarcts which may give rise to symptoms. These consist of bouts of fever associated with cough, sometimes with haemoptysis, and shortness of breath. Examination of the chest may reveal areas of crepitations. Brandt (1949) had recorded a generalized pulmonary fibrosis which he attributed to *Oxyuris*. Skiagrams will show patchy transient areas of opacity. Blood count in all cases shows a marked eosinophilia (Löffler, Essellier, and Macedo, 1948).

Prophylactic treatment includes the provision of proper sanitation, the wearing of boots and the treatment of carriers. The three specific drugs are tetrachlorethylene, carbon tetrachloride, and oil of chenopodium. Tetrachlorethylene or carbon tetrachloride, 3-4 millilitres, and 1 millilitre of oil of chenopodium followed by a saline purgative is the best routine treatment. Three doses of santonin, 3 grains may be given on alternate days in the treatment of infection by *Ascaris*, and in this condition hexylresorcinol, 1 grain, may be used. Gentian violet with an adult dose of 1 grain in an enteric coated capsule thrice daily for 8 days is the best treatment for *Oxuris* and *Strongyloides* infestation.

*Filaria bancrofti* is the most important of the tissue round worms. It is common in India, the West Indies, Porto Rico, Southern China and the Pacific Islands. The intermediate host is the mosquito, which sucks the embryos out of the blood at night. Development in the mosquito then takes from 10 to 40 days and the mature embryos are then inoculated into man through the proboscis.

Most of the literature of the disease is concerned with elephantiasis, a complication of filariasis observed in natives after prolonged exposure. The acute and early stages of the disease in white man were reviewed by Coggeshall (1946) and consist mainly of lymphangitis, lymphoedema, lymphadenitis, and funiculitis, and are accompanied by aches and pains in the major lymphatic areas. Jaffe (1945) found that the lung was in rare instances, involved with the skiagrams showing a patchy type of consolidation with clearing in one part of the lung and development of patches of infiltration in other areas.

Hetrazan (1 diethyl-carbamyl-4 methylpiperazine hydrochloride) given by mouth in dosage of 1-2 milligrams per kilo of bodyweight thrice daily for a period of 2-4 weeks is the best treatment. Antimony compounds such as Neostibosan and Fouadin have also been used. Prophylaxis includes destruction of mosquitos and the use of mosquito nets.

## THORACIC AMOEBIASIS

Amoebiasis is a disease widely distributed throughout the tropics. *Endamoeba histolytica* gains entrance to the body in its cystic form through the mouth in food or water. The cysts pass through the stomach, their walls are then digested by the intestinal juices, the amoebae escape and invade the colonic mucosa giving rise to the classical intestinal amoebiasis. They may then pass through the portal vein to the liver causing either a diffuse hepatitis or a liver abscess. Mostly thoracic amoebiasis is secondary to these conditions but solitary abscesses within the lung substance and not adjacent to the right diaphragm have been described by Dormer and Friedlander (1941), Manson Bahr (1923) and Zaky (1943). The abscesses must be embolic in origin and it is most probable that the amoebae reached the pulmonary circulation by going through the haemorrhoidal veins into the inferior vena cava, or by anastomotic channels between the venules of the portal and vena caval systems within the liver. Derrick (1949) reported a fatal case of amoebiasis with lung involvement caused by *Iodamoeba butschlii* that occurred in a Japanese prisoner of war camp. There are no other pathogenic amoebae.

Pulmonary disease in relation to the right diaphragm is probably much commoner than is generally recognized. The nature and extent of the disease depending on




FIG 97 — (a)  
Amoebic hepatitis  
showing humping  
of the diaphragm and  
exaggeration of  
basal markings  
(b) showing the  
humping of the  
diaphragm in  
this patient it is  
posterior



that in the liver. These thoracic complications nearly always occur as a direct extension of the abscess or hepatitis through the diaphragm into the pleural cavity, or into the lung through adhesions that have resulted from the inflammatory process. Abscesses may form in the lung, or the disease may in fact be the drainage of an hepatic abscess through the bronchus.

In at least half the cases of thoracic amoebiasis there will be no history of previous dysentery, though the patient may have had an unexplained fever with pain over the liver from amoebic hepatitis. The most usual initial symptoms of the thoracic disease is pain in the lower right chest and in the right shoulder. Cough comes next, at first, unproductive, but later giving rise to haemoptysis and the classical "anchovy paste" sputum. There will always be fever, sometimes rigors together with anorexia, fatigue, malaise and often extreme loss of weight.

Besides this usual manifestation of thoracic amoebiasis lesser pulmonary conditions may result as sympathetic reaction to the liver disease as is stressed by Bookless (1950). These include restricted movement of the right dome diaphragm, serous pleural effusion on the right side, or basal fibrinous pleurisy with a pleural rub, bronchitis, right basal bronchopneumonia and even right lower lobar pneumonia.

Empyema is a serious and frequent complication, as may be expected in an infection which crosses the diaphragm to reach the lung. Oschner and DeBakey (1943) have recorded a mortality rate of 78 per cent of cases reported in the literature with this complication, as against 40 per cent with a lung abscess.

The x ray appearance of the chest in thoracic amoebiasis is characteristic (Fig. 97). There is a hump like localized elevation of the diaphragm, which is best seen in the lateral picture. Radiating up from this to the hilum will be exaggerated markings made by the track of the infection. If a pulmonary abscess develops there will be the usual appearance of an abscess, and similarly with pleural effusion or empyema the picture characteristic of these conditions will appear.

### Treatment

Emetine is a specific drug for the treatment of thoracic amoebiasis. As soon as the diagnosis is made a course of emetine hydrochloride, 1 grain, given intramuscularly daily for 10-12 days, should be instituted. Oschner and DeBakey (1943) showed that among 153 patients with this disease who were not treated with emetine 56.6 per cent died, whereas where emetine was used the mortality rate dropped to 1.9 per cent, while in their own personal series the mortality rate of those who did not have emetine was 60 per cent, whereas there were no deaths in those who were treated with this drug.

Elsden Dew, Armstrong and Wilmot (1952) have shown that aurcomycin and terramycin are effective in the treatment of acute amoebic dysentery, though relapse is not unusual and a state of symptomless cyst passing is apt to develop. It is probable that the main action of these drugs is on organisms other than amoebae but there is evidence that *in vitro* they do have some direct anti amoebic effect. There is therefore strong reasons for using these two drugs in conjunction with the established amoebicides.

Surgical intervention, as is pointed out by Shraw (1949) is needed only when secondary infection has occurred, when there is an empyema which cannot be cured



## CHAPTER 18

### HYDATID DISEASE OF THE LUNG

GIOFFREY FLAVELL

#### AETIOLOGY

THE DEFINITIVE host of *Taenia echinococcus* is the dog, in whose intestine the worm of 0.5 centimetre length lives, the intermediate hosts are sheep, pigs or men, in whom it undergoes its larval and encysted forms. Dogs become infested by eating sheep's offal, and man by direct contact with dogs or from eating lettuce and other produce contaminated by their faeces. In Argentina, Australia, New Zealand, the Middle East, Iceland, and some parts of Wales the disease is common, elsewhere it is rare. The liver is the favoured site of cysts, accounting for 75 per cent of the cases, but, according to experienced Australian surgeons, up to 25 per cent are found in the lung.

After the ova have been discharged from the gravid terminal segment of a worm, and have found their way into the human stomach, gastric juices digest their outer capsule freeing an embryo, or hexacanth, from its six thorns, which migrates through the stomach wall, enters the portal vein, and so reaches the liver. Should it pass this filter it finds itself in the inferior vena cava, and so passes via the right heart into the lung, where it lodges, and enters a larvated stage. The embryo vacuolates and forms a small vesicle composed of two walls—an inner nucleated germinal layer and an outer non-nucleated laminated layer. This now expands into a cyst which is small at first, but, if it is not disturbed, becomes progressively larger until it may reach a considerable size. From the germinal layer bud off groups of cells that in turn vacuolate to form proliferative "brood capsules" giving rise to scolices, the heads and source of new worms. Most of these become detached, and sink through the hydatid fluid, forming "hydatid sand" upon the floor of the cyst, while outside the laminated layer the lung reacts to its presence by condensing an adventitious layer of fibrous tissue about it, but easily separated from it, as is the skin from a banana.

#### CLINICAL PICTURE AND DIAGNOSIS

The fate of the cyst, and the symptoms of the patient, now depend upon the complications which may arise. During the period of growth the cyst gives no

fixation test is now seldom used. As the cyst increases in size pain may occur, together with pressure symptoms, and occasionally there may be anaphylaxis with



by repeated aspirations, when a persistent abnormal channel between the bile duct system and bronchi persists or when fibrosis of the pulmonary tissue continues to produce symptoms as a result of the abscess. The operations which will most frequently be required are drainage of empyema or liver abscess in the latter case the mortality will be greatly reduced by using the extra serous route, or lobectomy or thoractomy when there is damage to the lung.

## BIBLIOGRAPHY

- Bookless A S (1950) Thoracic Amoebiasis *J R Army med Cps* 94 52
- Brandt M (1949) Parasitic Pulmonary Fibrosis due to *Oxyuris Vermicularis* *Tuberculo-sis* 3, 655
- Canizares M and Celis J (1951) Observations on Paragonimiasis at the Querzon Institute *Dis eases Chicago* 19, 668
- Coggeshall L T (1946) Filariasis in the Serviceman *J Amer med Ass* 131, 8
- Dairs L J (1945) Pulmonary Ascariasis in Monkeys *Brit med J* 1, 482
- Derrick E H (1948) Fatal Case of Generalized Amoebiasis due to Protozoon closely resembling if not identical with *Iodamoeba Butschlii* *Trans Roy Soc Trop Med Hyg* 42 191
- Dormer B A and Friedlander J (1941) Amoebiasis—Pulmonary Complications *Brit med J* 2 258
- Elsden Dew R, Armstrong T G and Wilmot A J (1952) Antibiotics and Amoebic Dysentery *Lancet* 2 104
- Faust E C (1922) A New Type of Amoeba Parasitic in Man Observed in N China. *J Parasitol* 2 22
- Garduño D M and Carpio D (1950) Paragonimiasis. A report of 3 cases with pulmonary and neurological symptoms *J Philipp med Ass* 26 253
- Jaffe H L (1945) Evaluation of Roentgen Therapy in Filariasis *Amer J Roentgenol* 53 483
- Löffler W, Essellier A and Macedo M M (1948) Pathogenesis and Aetiology of Transient Pulmonary Infiltrates with Eosinophilia of the Blood *Helv med Acta* 15 371
- Manson Bahr P (1923) Pulmonary Amoebiasis *Lancet* 2 599
- Miller J J, Jun and Wilbur D L (1944) Paragonimiasis (endemic hemoptysis) report of 3 cases *US Nav med Bull* 42 108
- Ochsner A and DeBakey H (1943) Amoebic Hepatitis and Hepatic Abscess analysis of 181 cases with review of the literature *Surgery* 13 460
- Ravelli A (1949) X ray Appearances of Parasites in the Lungs of Human Beings *Med Klin* 44, 603
- Shaw R R (1949) Thoracic Complications of Amoebiasis *Surg gynaec Obstet* 88 753
- Shu Ngoh Koo and Dye di Woo (1949) Pleural Paragonimiasis complicated by Empyema Thoracis *Chin med J* 67, 211
- Soyas E and Jayawardena M D S (1945) Pulmonary Ascariasis. A possible cause of asthma *Brit med J* 1 1
- Tillman A J B and Phillips H S (1948) Pulmonary Paragonimiasis *Amer J Med* 5 167
- Zaky H A (1943) Pulmonary Amoebiasis *J R Egypt med Ass* 26, 317

## CHAPTER 18

### HYDATID DISEASE OF THE LUNG

GEOFFREY FLAVELL

#### AETIOLOGY

THE DEFINITIVE host of *Taenia echinococcus* is the dog, in whose intestine the worm of 0.5 centimetre length lives, the intermediate hosts are sheep, pigs or men, in whom it undergoes its larval and encysted forms. Dogs become infested by eating sheep's offal, and man by direct contact with dogs or from eating lettuce and other produce contaminated by their faeces. In Argentina, Australia, New Zealand, the Middle East, Iceland, and some parts of Wales the disease is common, elsewhere it is rare. The liver is the favoured site of cysts, accounting for 75 per cent of the cases but, according to experienced Australian surgeons, up to 25 per cent are found in the lung.

After the ova have been discharged from the gravid terminal segment of a worm, and have found their way into the human stomach, gastric juices digest their outer capsule freeing an embryo, or hexacanth, from its six thorns, which migrates through the stomach wall, enters the portal vein, and so reaches the liver. Should it pass this filter it finds itself in the inferior vena cava, and so passes via the right heart into the lung, where it lodges, and enters a larvated stage. The embryo vacuolates and forms a small vesicle composed of two walls—an inner nucleated germinal layer and an outer non nucleated laminated layer. This now expands into

rise to scolices, the heads and source of new worms. Most of these become detached, and sink through the hydatid fluid, forming "hydatid sand" upon the floor of the cyst, while outside the laminated layer the lung reacts to its presence by condensing an adventitious layer of fibrous tissue about it, but easily separated from it, as is the skin from a banana.

#### CLINICAL PICTURE AND DIAGNOSIS

The fate of the cyst, and the symptoms of the patient, now depend upon the complications which may arise. During the period of growth the cyst gives no

at this stage is by the intra-dermal Casoni reaction, which is not always reliable, or the existence of an eosinophilia of 4 per cent or more. The Weinberg complement fixation test is now seldom used. As the cyst increases in size pain may occur, together with pressure symptoms, and occasionally there may be anaphylaxis with



## CHAPTER 18

### HYDATID DISEASE OF THE LUNG

GEOFFREY FLAVELL

#### AETIOLOGY

THE DEFINITIVE host of *Taenia echinococcus* is the dog, in whose intestine the worm of 0.5 centimetre length lives, the intermediate hosts are sheep, pigs or men in whom it undergoes its larval and encysted forms. Dogs become infested by eating sheep's offal, and man by direct contact with dogs or from eating lettuce and other produce contaminated by their faeces. In Argentina, Australia, New Zealand, the Middle East, Iceland, and some parts of Wales the disease is common, elsewhere it is rare. The liver is the favoured site of cysts, accounting for 75 per cent of the cases, but, according to experienced Australian surgeons, up to 25 per cent are found in the lung.

After the ova have been discharged from the gravid terminal segment of a worm, and have found their way into the human stomach, gastric juices digest their outer capsule freeing an embryo, or hexacanth, from its six thorns, which migrates through the stomach wall, enters the portal vein, and so reaches the liver. Should it pass this filter it finds itself in the inferior vena cava, and so passes via the right heart into the lung, where it lodges, and enters a larvated stage. The embryo vacuolates and forms a small vesicle composed of two walls—an inner nucleated germinal layer and an outer non-nucleated laminated layer. This now expands into a cyst which is small at first, but, if it is not disturbed, becomes progressively larger until it may reach a considerable size. From the germinal layer bud off groups of cells that in turn vacuolate to form proliferative "brood capsules" giving rise to scolices, the heads and source of new worms. Most of these become detached, and sink through the hydatid fluid, forming "hydatid sand" upon the floor of the cyst, while outside the laminated layer the lung reacts to its presence by condensing an adventitious layer of fibrous tissue about it, but easily separated from it, as is the skin from a banana.

#### CLINICAL PICTURE AND DIAGNOSIS

The fate of the cyst, and the symptoms of the patient, now depend upon the complications which may arise. During the period of growth the cyst gives no

fixation test is now seldom used. As the cyst increases in size pain may occur, together with pressure symptoms, and occasionally there may be anaphylaxis with

# HYDATID DISEASE OF THE LUNG




FIG 98 — (a and b)

Large hydatid cyst situated in lower lobe of left lung in a young soldier who had served for a short time in Greece where he had kept a pet dog. A Casoni test before operation was negative, and from its position and appearance the x ray opacity was thought to be caused by a neuro-fibroma. Successful removal at operation, with no recurrence.

(a)



(b)

## CLINICAL PICTURE AND DIAGNOSIS

severe bronchospasm dyspnoea and tachycardia though this is more often seen during complications or after surgical intervention

The zone of atelectatic and fibrous tissue of the adventitious layer tends gradually to cause pressure erosion as it comes to abut upon a bronchial wall and finally a fistula is established. Air escapes into the adventitious pericyst stripping the true hydatid from it. If the cyst is small it may now be expectorated *in toto* and the patient be spontaneously cured. In many instances rupture of the cyst occurs and hooklets scolices or fragments of laminated membrane may be expectorated and confirm the diagnosis. The cyst does not necessarily become infected at this juncture though it may do so but either dies or continues to live and perhaps to produce daughter cysts.

The intrusion of air into the pericyst with separation of the hydatid sometimes gives rise to x ray appearances which Argentinian and French writers have described as perivesicular pneumonia the double dome arch or Cumbo's sign and camelote formation according to the amount of air between the pericyst and the cyst itself since this air shows as a crescentic translucent area between the two opaque convexities of the cysts. Unfortunately these so called pathognomonic signs are neither reliable nor common.

If the cyst ruptures into a bronchus or the pleura scolices are spread and multiple cysts develop simultaneously bronchial and pleural fistulae result additionally in pneumothorax (Fig 99). Metastatic hydatidosis follows erosion and rupture into a pulmonary vessel. Multiple cysts of this kind occur in about 6 per cent of infestations and are commoner in children. Rupture itself carries a heavy mortality.

Once a fistula has been established ultimate infection will probably follow bringing with it purulent sputum fever haemoptysis and general evidence of pulmonary suppuration. If a hydatid cyst has obstructed a bronchus however collapse and secondary bronchiectasis may result with similar symptoms. Infection also adds to the risk of pleural rupture since empyema will then accompany dissemination. Sometimes the cyst dies after it has become infected but daughter cysts can continue living even amongst pus. They are more commonly associated with hydatid cysts of the liver however and should they appear in the sputum hepatic involvement must be feared for liver cysts when infected can perforate the diaphragm and rupture into the lung the pleura or the mediastinum. Infestation of more than one organ occurs in less than 2 per cent of the total number of cases. Hydatid cysts seldom die except as a result of rupture or infection and even calcification of the cyst does not necessarily imply the death of the parasite. If a cyst is removed or is coughed up the pericyst is usually soon obliterated by re expansion of the lung but should it be unusually rigid may persist and in the course of time become epithelialized by extension from the bronchial mucosa and form a permanent lung cyst which in its turn may become the seat of infection.

## TREATMENT

The treatment of hydatid cysts of the lung is surgical and has for its object the complete removal of the parasite with maximum conservation of pulmonary tissue. Expectant treatment (except perhaps in the case of very small cysts detected

# HYDATID DISEASE OF THE LUNG




FIG. 99 — (a) Large hydatid cyst in left lower lobe of a school boy evacuated to Wales during World War II. While awaiting admission to hospital he played a game of football and the cyst ruptured into the pleural cavity producing a hydropneumothorax, (b) thick cyst wall can now be seen. Moderate anaphylaxis developed. At operation the cyst was found to be heavily infected and a bronchial fistula was also present, (c) left lower lobectomy with complete re-expansion of remaining lung with no recurrence.

(a)



(b)





(c)

by routine radiography), transpleural aspiration, the induction of an artificial pneumothorax, and even bronchoscopy are contra-indicated since all may favour rupture with dissemination. After careful radiographic localization the chest is opened where access to the cyst can best be obtained. Pleural adhesions cannot be relied upon securely to isolate the rest of the pleural cavity, nor does it seem profitable to try and secure adherence by packing the extrapleural space or sewing visceral to parietal pleurae, and removing the cyst at a later stage. The pleural space is carefully packed off, some surgeons employing black towels the better to detect contamination, and the cyst fluid is partially and slowly aspirated with a fine-gauge needle attached to a two-way syringe. A small incision is then made down to the pericyst and the glistening white hydatid is removed from it complete, meticulous care being taken to avoid rupture or the spilling of any of the contents. The interior of the pericyst is then inspected, bleeding points are secured, and any fistula that may exist closed. The walls of the pericyst itself may be approximated, though it usually soon obliterates. Multiple cysts are dealt with similarly.

Should serious bleeding occur, as it sometimes does, or if a cyst be very large, or accompanied by bronchiectasis, lobectomy is performed. When the cyst has already ruptured, and is infected, lobectomy may likewise be necessary, however, every effort should be made to conserve the lung, and if local removal of the cyst can be achieved without further contamination it is to be preferred, both pericyst and pleura being drained.



## HYDATID DISEASE OF THE LUNG

Barrett (1949) has pointed out that hydatid cysts of the lung show a natural tendency to herniate from the lung as they increase in size, and has recently devised a technique of removal which profits by this, and delivers the cyst intact avoiding even preliminary aspiration which with the finest needle may sometimes lead to a tear of the laminated membrane. With patient's head lowered, so that if rupture occurs into a bronchus natural drainage takes place, and with closed circuit anaesthesia, the chest is widely opened and the cyst identified. A macintosh bag, with strings round its neck and its bottom open, is then placed over the affected lobe and the strings drawn just tightly enough about the hilum not to obstruct the circulation. The open end is brought out of the wound and spread on the surface so that the part is completely isolated within its bag. The adventitia is then incised until the bluish-white laminated membrane is just exposed at one point, and freed gently for about an inch. The anaesthetist now softly increases the pressure in his circuit, and little by little over a period of 10-30 minutes, the adventitia cracks along the line of the incision and the cyst gently extrudes, to fall at last intact into the protecting bag.

Many hydatid cysts of the heart have been described, and are said to occur in some 2 per cent of infestations. Occasionally recognized radiologically, they are associated at times with paroxysmal tachycardia, and often terminate fatally by rupture which may result in embolism from the liberated daughter cysts. d'Abreu (1950) has described his successful removal of such a cyst from the wall of the left ventricle employing the usual technique, with preliminary aspiration, for hydatids of the lung, and it seems probable that his lead will be followed by increasing numbers of direct attacks upon this hitherto inviolate refuge of the parasites.

## BIBLIOGRAPHY

- d Abreu A. L. (1950) 'The Removal of a Hydatid Cyst from the Wall of the Left Ventricle' *Thorax*, 5, 362  
 Barrett N. R. (1947) 'The Treatment of Pulmonary Hydatid Disease' *Thorax*, 2, 21  
 — (1949) 'Removal of Simple Univesicular Pulmonary Hydatid Cyst' *Lancet* 2, 234

...

small Tachy-


## CHAPTER 19

### SYPHILIS OF THE LUNG

KENNETH M. A. PERRY

SYPHILIS of the lung is a very rare condition. It may be either congenital or acquired. The congenital form is the commoner and is usually found in stillborn syphilitic infants. It has been called *pneumonia alba*. The gross appearance of the lung is characteristic, the lung being of a pale greyish colour and consolidated throughout, with the essential portions sinking in water, and the surface dry and smooth. The histology is that of arrested development rather than active inflammation. Either the alveoli have failed to develop or they are smaller than normal and lined with cubical epithelium. In adults, gummas may develop in the lungs, they may be either single or multiple, when they are multiple they are usually small and do not give rise to symptoms.

FIG 100—A girl aged 15 years with congenital syphilis and a positive Wassermann reaction. The radiograph shows a gumma in the right lower lobe which was subsequently proved histologically



Large gummas, however, may be up to 1½ inches in diameter (Fig 100). They sometimes undergo softening and in such cases they may present as a lung abscess. The patient may then complain of fever, cough and persistent purulent sputum.

# SYPHILIS OF THE LUNG



FIG 101—(a) Female aged 35 years with swelling over manu-  
brum and rales at left base Wassermann  
reaction positive (b) 3 months later after  
treatment with potas-  
sium iodide 25 grains  
3 times a day and  
intramuscular injec-  
tions of bismuth



## TREATMENT

Examination of the chest may show diminished air entry and rales in the area affected. The process may progress and give rise to lesions which are similar to bronchiectasis or fibro caseous tuberculosis.

## DIAGNOSIS

The diagnosis is always difficult but the Wassermann reaction will be positive. Skiagrams of the chest show either a solid opacity or a cavity. Tubercle bacilli will be persistently absent from the sputum. The diagnosis may only be proved when an excised portion of the lung is examined histologically.

## TREATMENT

If the lesion is not soft and broken down it may respond satisfactorily to anti syphilitic treatment. Arsenic, bismuth, mercury and penicillin are the drugs used to destroy the spirochaetes but iodides are valuable in stimulating the removal of granulomatous tissue (Fig 101). The following is satisfactory treatment. On the first day the patient is given an intravenous injection of 0.45 gramme neoarsphenamine and an intramuscular injection of 0.3 gramme bismuth oxychloride followed on the next five days with intramuscular injections of 1 million units of procaine penicillin. On the seventh day 0.6 gramme of neoarsphenamine and 0.3 gramme of bismuth oxychloride are administered. This should be followed by six weekly doses of 0.6 gramme neoarsphenamine and 0.3 gramme bismuth oxychloride. A second and third course may be given if necessary. If however the lesion has broken down to form cavities and lung tissue has been destroyed the only method of treatment is by excision of the affected part of the lung. Any attempt at drainage will be followed by permanent sinus formation.

## BIBLIOGRAPHY

- Domer B A, Friedlander J and Wiles, F J (1945) Syphilis of the Lung in Bantus  
*Brit J Tuberc* 39 85  
 Hartung A and Fredman J (1932) Pulmonary Syphilis a report of 3 cases of acquired  
 lung syphilis in adults with particular reference to Roentgen aspects *J Amer med Ass*  
 98 1969  
 McIntyre M C (1931) Pulmonary Syphilis *Arch Path* 11 258  
 Morgan A D, Lloyd W E and Price Thomas C (1952) Tertiary Syphilis of the Lung  
 and its Diagnosis *Thorax* 7 125  
 O'Leary P A and Ackerly O E (1945) Syphilis of the Lung *J Lancet* 63 154

## CHAPTER 20

### PULMONARY EMBOLISM

W D W BROOKS

#### AETIOLOGY AND INCIDENCE

PULMONARY embolism is not a new disease but in recent years it has become increasingly aware of for a variety of reasons new in and treatment. It may be defined as a result of abnormal st

Many different kinds of emboli may thus reach the lung. For example, air may enter the jugular vein when the throat is cut and so enter the circulation in the heart and pulmonary in the blood and so pulmonary emboli embarrassment when rapid decompression rapidly to the surface. The larval stage of some parasites may cause pulmonary embolism, for instance, in ankylostomiasis. Bony fracture or penetrating wounds involving fatty tissues, or the use of mineral oil as a lubricant when passing a catheter may admit large amounts of fat or oil to the circulation and so cause embolism.

emboli may derive from infected thrombi in cases of septic abortion, osteomyelitis, and in the course of pyaemia or septicaemia, while perhaps the commonest bacterial embolism may occur in the early months following primary tuberculous infection. Many of these emboli may give rise to specific effects within the lungs dependent upon their nature, and military tuberculosis, multiple staphylococcal lung abscesses or metastatic carcinomas illustrate their variety. These features are dealt with elsewhere, moreover, their aggregate is small in comparison with the common form of pulmonary embolism which arises as a result of the detachment of a clot or clots from a preformed venous or cardiac thrombosis. We are concerned here with this last and most frequent form; the special types mentioned above will be considered only in so far as they contribute to the clinical features common to all forms of pulmonary embolism.

that is within the hour, for at least 60 per cent of patients with fatal emboli survive longer, some for several days. Clinically diagnosed pulmonary embolism occurs in perhaps 1 per cent of all hospital patients, while certainly many more cases are undiagnosed or are thought to be instances of pneumonia, massive collapse of the lung, or other disorder.

## ÆTIOLOGY AND INCIDENCE

The primary source within the venous system is often not obvious. It is not sufficiently recognized that in apparently healthy adults the chance that the plantar or calf veins carry thrombi is about even, that skiagrams of the pelvis commonly show phleboliths and that it is uncommon not to find thrombosed veins in the broad ligament at hysterectomy. As Virchow showed the great majority of emboli arise from thrombi in the veins in these areas—the legs and the pelvis. All other areas including the right side of the heart are relatively uncommon sources. Ochsner has distinguished between thrombophlebitis due to local trauma or bacterial invasion with a relatively firm adherent clot and well marked local symptoms and signs and phlebothrombosis due to venous stasis and alterations in the cellular and fluid constituents of the blood which increase its coagulability with a loosely attached clot and few if any local clinical manifestations. In the former unless there is suppuration embolism from the thrombus is rare in the latter it is common. Conditions which predispose to phlebothrombosis include local venous obstruction immobilization and heart failure post operative and puerperal states specific fevers and polycythaemia. The incidence of venous thrombosis increases with age and in cold seasons and climates.

## PATHOLOGY

As a rule the obstruction of one or a few minor branches of the pulmonary artery by an embolus or emboli has no demonstrable effect upon the lung. So efficient is the blood supply via the bronchial arteries that major or many minor branches of the pulmonary artery may be obstructed without infarction of the lung occurring. Such thrombi contract become organized and recanalization with restoration of the circulation takes place. However when the circulation from the bronchial arteries is defective for example in heart failure or arteriosclerosis embolism more easily causes pulmonary infarction. The infarcted area of lung is at first swollen hard dry and bloody in colour. The adjacent pleura is roughened with a layer of fibrin and later quite a considerable pleural effusion may develop. As a rule the embolus can be found to be obstructing the pulmonary artery concerned. Microscopically the area of lung involved is grossly distended with coagulated blood and in the course of time death reabsorption of the endothelial cells lining the alveoli and coagulation necrosis occurs. It is followed by much contraction of the aging infarct and replacement with fibrous tissue. A further consequence of pulmonary embolism acute cor pulmonale does not develop unless about two-thirds of the pulmonary circulation is obstructed. Thus a rider embolus at the bifurcation of the pulmonary artery or numerous simultaneous or successive emboli in each lung may provoke sufficient rise of pressure proximally to cause acute right ventricular failure which may well be fatal. Infarction of the lung occurs variably in such cases. Apart from this cause of death which may be rapid massive pulmonary embolism may kill speedily from shock asphyxia and insufficient circulation of blood to the left heart. It is thought also that widespread vagal stimulation is provoked by pulmonary embolism affecting the heart pulmonary vascular tree bronchi and gastrointestinal tract. Such autonomic effects augment the above mentioned consequences of massive embolism and may well explain the occasional fatalities.



FIG 102 —  
Multiple sar-  
comatous  
emboli—pri-  
mary in tibia  
—with small  
bilateral  
effusions

following apparently minor emboli. Air, fat, or oil may cause either simple embolism, infarction or acute cor pulmonale, and death, while recurrent embolism especially with carcinomatous emboli occasionally gives rise to relatively slowly developing right ventricular failure during the course of which the sequelae of widespread pulmonary neoplastic metastases can develop (Fig 102). Finally, it must be remembered that minor and many major consequences of pulmonary embolism may be produced by thrombosis occurring in the pulmonary arterial tree itself, and that this is especially liable to occur when the pulmonary circulation is obstructed in heart failure or mitral stenosis. There is some evidence that digitalis is a significant predisposing factor to thrombosis in pulmonary and systemic vessels.

### SYMPTOMS AND SIGNS

Pulmonary embolism commonly occurs without symptoms or signs of any kind, and it is equally true that other features of the case when heart failure is present for example, may so obscure the clinical picture as to make diagnosis almost impossible. Quite often restlessness, fear, or vague uneasiness may be present, and there may be a slight increase in the venous pressure, which is referable to the legs,

but it is uncommon to find the classical signs there of thrombophlebitis because when oedema, pain, a tender hard vein, and local discoloration are present the thrombus is firmly adherent. The really dangerous phlebothrombosis is relatively silent but may cause slight swelling of the lower leg, and since it is usually situated in the deep veins the more superficial are often distended. Local tenderness, if carefully sought, may be found in the sole of the foot near the external malleolus, or in the calf, and dorsiflexion of the foot may cause pain in the same site. The temperature of the affected leg may be raised.

Variable local features may be associated with pulmonary embolism. For example, many cases confidently diagnosed as massive collapse or bronchopneumonia following operation or childbirth are really examples of embolism. The signs and symptoms in such cases may closely resemble those of the disorders they mimic.

However, while pulmonary embolism with or without infarction often has no characteristic clinical features within the chest, it is also true that quite clearly recognizable symptoms and signs may be produced there (Fig 103). An abrupt onset with pleuritic pain, cough, haemoptysis often lasting for some days, and cyanosis and dyspnoea may occur. These may be associated with a pleural rub, signs of consolidation, persistent local adventitious sounds, fever, tachycardia, and later transient haemolytic jaundice. It is not uncommon, when infarction of the lung has resulted, for an appreciable blood-stained pleural effusion to develop and to mask other signs. Since embolism is often multiple the signs frequently involve both lungs, moreover, successive episodes may take place

and air entry, and an impaired percussion note. Compensatory emphysema is associated, but it is unusual, save with infected emboli, for bronchiectasis to develop. Only large infarcts are likely to leave behind recognizable permanent signs such as these, and the patient thus damaged is likely to be easily dyspnoeic thereafter.

The characteristic and dominant feature of pulmonary embolism of a considerable order of magnitude is its striking catastrophic onset. Scarcely less significant is the association of symptoms and signs of shock, namely faintness, pallor, sweating, a low blood pressure, vomiting, and tachycardia with those mentioned above marking the arrival of the embolus within the pulmonary arterial tree.

When massive embolism occurs death may come swiftly by shock, virtual cessation of the circulation and asphyxia. With rather more protracted survival, and if more than about 60 per cent of the circulation is obstructed the syndrome known today as acute cor pulmonale may develop. As the embolism occurs the patient is seized with a sense of suffocation, a backward gasping, a faintness, and

As a rule there are no abnormal signs to be heard over the lungs (Fig 104). The cardiac signs are those of rapidly developing dilatation, right ventricular failure, and may include marked pulsation in the second and third interspaces.



# PULMONARY EMBOLISM



FIG 103 — Recent large infarct right lower lobe with an associated pleural effusion and slight mediastinal shift to the right. This occurred on the tenth post-operative day after partial gastrectomy.




FIG 104 — Multiple pulmonary emboli producing small infarcts with acute pulmonary hypertension and right ventricular strain.

## SYMPTOMS AND SIGNS

to the left of the sternum, protodiastolic gallop rhythm best heard in the same site, accentuation of the pulmonic second sound, a pericardial rub in this same area, and venous distension in the neck, and of the liver. The blood pressure falls markedly. The electrocardiogram is helpful in the difficult distinction from coronary infarction in a minority of cases. When this is so the main characteristics in acute cor pulmonale are —

- (1) A prominent S in lead I;
- (2) T is low or inverted in lead II,
- (3) In lead III a Q wave develops or is increased, and the S T segment may be elevated, T is deeply inverted
- (4) In lead IV (R or F) T is flat or inverted, while T inversion is common in C R I

### Radiological appearances

Correlation between radiological findings and the clinical severity of the case is not marked. Serious and repeated embolism may for instance occur and the chest be normal radiologically. Almost any type of parenchymal shadow may be observed if infarction occurs, the commonest being a peripherally situated irregular area of consolidation with an associated pleural reaction. The classical wedge-shaped area extending out from the hilum is rarely seen. Elevation of the diaphragm on the affected side, pleural effusion, and atelectasis often complicate embolism, while linear and more extensive scarring may remain permanently when the patient recovers.

## DIAGNOSIS AND DIFFERENTIAL DIAGNOSIS

Pulmonary embolism should be suspected when the appropriate clinical antecedent conditions exist if otherwise unexplained fever, dyspnoea, and tachycardia develop. Evidence of phlebothrombosis will increase the probability of the diagnosis. When an abrupt worsening of the patient's condition occurs accompanied by signs and symptoms referable to the chest of the type mentioned diagnosis is as a rule easy, particularly if shock is associated. In these cases radiology is often helpful. Massive collapse of the lung also occurs abruptly but usually has characteristic physical signs, in uncomplicated cases does not quickly give rise to haemoptysis, tends to follow sooner after operation, and the radiological findings are frequently diagnostic. Broncho-pneumonia is of much slower development, the sputum is purulent, the signs even more often bilateral. Coronary thrombosis may closely resemble acute cor pulmonale, but the pain is commonly more severe, pulmonary congestion and orthopnoea are often present, restlessness greater, and the electrocardiogram as a rule abnormal and different from that set forth above.

## TREATMENT

The prevention of thrombosis in the veins of the legs and pelvis is important. Many surgical and perhaps most medical patients could with advantage have the circulation in these areas improved by repeated breathing exercises in conjunction with massage and leg exercises, and so decrease the risk of thrombosis. In

hospital many patients are unnecessarily immobilized in bed. The provision of a dining hall and recreation room would do much to reduce this and incidentally the prevalence of pulmonary embolism. Finally, the proper treatment or prevention of anaemia, dehydration, and haemoconcentration are also significant prophylactic measures.

If phlebothrombosis is strongly suspected or known to be present anticoagulant therapy is desirable. Heparin is given at 4 hourly intervals in 50-milligram doses undiluted intravenously. Simultaneously, oral administration of dicoumarol is started using 300 milligrams on the first, and 200 milligrams on the second day.

The blood prothrombin levels must be determined on alternative days, and the dosage of dicoumarol is adjusted on subsequent days to keep this level between 30 and 60 per cent of normal. Heparin is discontinued after 2 days, but, if it is essential, the dicoumarol may be continued for 1 or more months. The drug should not be discontinued abruptly, but should be reduced slowly over the period of a week. Breathing exercises at this stage are contra indicated. Tromexan may be used as an alternative to dicoumarol. Its action is quicker, and it is less dangerous. However, great variability is shown by different patients, and by the same patient from time to time, to its action. Marked drug resistance is quite commonly a most irritating disadvantage.

A loading dose of 1,200 milligrams, in an urgent case accompanied by 100 milligrams of heparin given intravenously, is followed next day by 900 milligrams, and thereafter 300 milligrams daily are given by mouth, provided estimates show that this dosage is adequate to give rise to and maintain a prothrombin time between 10-30 per cent of the normal.

Anticoagulant therapy may also of course be used after embolism has occurred, with the object of preventing further formation of thrombus and so diminishing the risk of recurrent embolism which may well be much more severe. Contra-indications to the use of this treatment include severe kidney and liver disease, blood dyscrasias, recent operation on the brain and spinal cord, high fever, and bacterial endocarditis.

The method of treatment is not without danger and, since fatal haemorrhage may occur, the above mentioned method of control is essential. Even so it is urged that if any spontaneous haemorrhage however slight occurs, or if the patient complains of backache dicoumarol administration should be stopped.

Surgical measures have been devised to prevent emboli reaching the lungs. They include proximal vein ligation, for example the femoral vein if the calf veins are the site of phlebothrombosis, or if both legs are involved ligation of the inferior vena cava. These methods are often combined with anticoagulant therapy. While the proper limitations of this method of treatment are not yet settled there exists a measure of agreement that it is of value (1) after unexpected pulmonary embolism, for which good evidence of the source exists, (2) when phlebothrombosis develops and the patient is likely to be immobilized for some other cause, and (3) to prevent spread of the clotting process.

## TREATMENT

When massive pulmonary embolism occurs atropine sulphate  $\frac{1}{2}$  grain, and  $\frac{1}{4}$ -1 grain of papaverine hydrochloride should be given intravenously at once. Oxygen should be administered, and anticoagulant therapy started. Because morphine is alleged to increase the coagulability of the blood pethidine or physeptone are preferable for the relief of pain, while papaverine is more effective in preventing vagal effects. Pulmonary embolectomy has a great mortality, and should only be used in those patients surviving the immediate risk for whom early death seems otherwise inevitable. Venesection, digitalis, and vasodilator drugs are valueless. Intravenous strophanthin 0.5 milligram followed by 0.25 milligram every 8 hours for 2 days may help increase the cardiac output. About two thirds of the cases of acute cor pulmonale survive, and for these after the emergency is over good nursing anticoagulants and possibly venous ligation are essential.

When pulmonary infarction occurs, depending on the severity of the case measures selected from those mentioned above will be needed. In addition pleuritic pain should be relieved by strapping and an electrically heated pad. Expectoration should be encouraged. In some cases, at a later stage iron by mouth and liver injections may be required if a significant anaemia has developed.

## PROGNOSIS

Recovery is finally complete in a majority of cases of pulmonary embolism. At the onset however, it is rarely possible to give any indication at all from the available clinical features of the fate of a particular case. The risk of repeated emboli should be remembered in even the mildest case. In general however, the more severe the early symptoms and signs, the graver is the prognosis.

## BIBLIOGRAPHY

- Castleman B (1940) Healed Pulmonary Infarcts *Arch Path* 30 130  
 Hampton, A O and Castleman B (1940) "Correlation of Post mortem Chest Teleroentgenograms with Autopsy Findings with Special Reference to Pulmonary Embolism and Infarction" *Amer J Roentgenol* 43, 305  
 Howkins J (1938) Movement of the Diaphragm after Operation *Lancet* 1, 85  
 Krause G R (1945) The Roentgen Diagnosis of Pulmonary Infarcts *45, 107*  
 — and Carter E H (1945) Infarction of the Lung *Arch intern Med* 67, 1144  
 Loeffler L (1941) Pulmonary Embolism and Infarction *Arch Path* 31, 93  
 Miller R and Berry J B (1951) Early Pleural Effusion in Pulmonary Embolism *Acta Rad* 29, 7  
 Noberg G (1948) A Radiological Study of Pulmonary Infarction *Quart J Med* 20, 233  
 Street D S (1951) Experiences with Venous Thrombosis and Pulmonary Embolism with Special Reference to Anticoagulant Therapy *Amer J med Sci* 219 117  
 Vanderveer J B Kuo P T and Marshall D G (1950) On the Roentgen Diagnosis of Lung Embolism *Acta Radiol* 19, 357  
 Westerman N (1938)

## CHAPTER 21

### BLAST INJURIES OF THE LUNG

KENNETH M A PERRY

#### AETIOLOGY

ZUCKERMAN (1940) showed that in animals exposed to blast, haemorrhages occurred in the lungs. These could consist in (1) a few small haemorrhages on the surface (2) patchy haemorrhages scattered throughout the lungs or (3) complete hepatization of the lungs by haemorrhages. He thought that the lesions were produced by the wave of positive pressure acting directly on the chest wall, and this theory is generally accepted.

#### CLINICAL FEATURES

##### Condition immediately after exposure

The patient is frequently unconscious for a short time at the time of the injury, when he comes round he is usually quiet and calm, in contrast with the anxiety of patients who have large haemorrhages. The most striking symptom is shortness of breath, which may last for several days after the injury. Breathing is sometimes painful, and the patient usually complains of a feeling of tightness in the chest. Apart from pain on breathing, the patients sometimes complain of retrosternal pain, or pain in the intercostal muscles. These may remain tender for a week or more. Most patients develop a cough and expectorate frothy (usually blood stained) sputum for several hours after the injury. After this the cough often becomes dry. The patients sometimes complain of epistaxis and deafness.

##### Condition on examination in hospital

Temperature, pulse and respiratory rates are usually normal by the time that the patient is admitted to hospital. In the chest, dullness, diminished breath sounds and decreased fremitus are usually present. Generalized rhonchi are found in most cases and persist for a few days. The radiological appearances are variable and there is nothing specific about them. They may be unilateral or bilateral.

the patients usually recover

#### TREATMENT

Treatment is essentially symptomatic, rest being the most important single requirement, but the areas of haemorrhage sometimes become infected and pneumonia

## BIBLIOGRAPHY

may develop. This should be treated by chemotherapy. Physiotherapy is indicated for the muscular pain which is so prominent a symptom. Breathing exercises are also of great value.

## BIBLIOGRAPHY

- Fearnley G R (1945) Blast Injury to the Lungs. Clinical and radiological findings and their relation to certain symptoms. *Brit med J* **1** 474
- Hadfield G (1940) Discussion of the Problem of Blast Injuries. *Proc R Soc Med* **34** 189
- (1941) Lung Injuries in Air Raids. A discussion on pathology and diagnosis. *Brit med J* **2** 239
- O'Reilly J N (1941) Discussion of Problem of Blast Injuries. *Proc R Soc Med*, **34** 189

## CHAPTER 22

### PULMONARY HAEMOSIDEROSIS

KENNETH M. A. PERRY

#### AETIOLOGY

PULMONARY haemosiderosis is described in two associations, firstly as an idiopathic disease, and secondly in conjunction with mitral stenosis. Idiopathic haemosiderosis is a rare condition which occurs mostly in children under the age of 16 years. Borsos-Nachtnebel (1942) described the only case in the literature in an adult; the patient was a man aged 38 years. Ceelen (1931) was the first to describe the condition. Wyllie and his colleagues (1948) collected 17 cases from the literature and described 7 of their own, and further cases have been reported by Waldenström (1948), Nancekrievill (1949), King (1949), and Walton and Williams (1951).

#### IDIOPATHIC PULMONARY HAEMOSIDEROSIS

##### Clinical picture and diagnosis

Symptoms consist in recurrent attacks of fatigue, pallor, cyanosis and increasing dyspnoea and palpitation. These symptoms may be of sudden onset. There is usually a cough, which is troublesome and often associated with vomiting. Haemoptysis is frequently present. There is sometimes severe abdominal pain and the patient becomes restless and apprehensive. Each attack commonly lasts

failure resulting from a severe disturbance of the pulmonary circulation. This is well shown by a diffuse, extreme mottling of the chest wall and show (see Fig. 105).

more accentuated during an acute attack, when there may be evidence of a partial lobar or lobular collapse. The mottling is usually much finer than that seen in miliary tuberculosis or sarcoidosis. The speckling is also smaller than is seen in occupational siderosis.

##### Necropsy findings


At necropsy the main changes are to be found in the lungs. Macroscopically these appear full and firm, of a dark reddish brown colour, with fine haemorrhages visible on the pleural surfaces. On section the lungs appear solid. Enlargement of the broncho-pulmonary and tracheo-bronchial lymph-nodes is conspicuous. The right side of the heart is usually increased in size.

# IDIOPATHIC PULMONARY HAEMOSIDEROSIS



FIG 105 —  
Radiograph of  
(a) brother and  
(b) sister showing  
idiopathic  
pulmonary  
haemosidero-  
sis

(a)





Microscopically, the alveoli are filled with haemosiderin-laden phagocytes. The epithelial cells of the alveolar walls also contain haemosiderin, and many are desquamated within the alveoli, which also contain red blood cells and haemosiderin granules lying free. Haemosiderin granules are also found in the interstitial tissues and the lymph nodes. Diffuse pulmonary fibrosis, capillary fibrosis and fragmentation of elastic fibres in the alveolar walls, deposits of haemosiderin from previous haemorrhages and fresh bleeding into the alveoli are also sometimes found.

### Course and prognosis

The cause of the condition is obscure. The prognosis is bad, and there is no treatment which appears to alter the course of the disease.

## HAEMOSIDEROSIS ASSOCIATED WITH MITRAL STENOSIS

### Pathogenesis

It had long been known that the lungs of patients with mitral stenosis show brown induration, but it was not until 1928 that Rosenhagen drew attention to the densely mottled skiagrams of the chest which are sometimes seen in patients with mitral stenosis. He showed that these were caused by nodulation and were microscopically multiple focal deposits of haemosiderin. Since then the condition has been fully described in America by Soma Weiss (1942) and Pendergrass, Lane and Ostrum (1949), and in Great Britain by Lendrum, Scott and Park (1950). Why the condition occurs in certain cases of mitral stenosis and not in others is not clear. Pendergrass and Leopold (1945) suggest that the deposit might be the result of perivascular interstitial haemorrhages, which they believed to occur at the acute phase of the rheumatic disease. Other writers still accept pulmonary congestion as an adequate explanation of the skiagraphic nodulation, but Lendrum, Scott and Park (1950) believe that the foci of the haemosiderin are situated in groups of adjacent alveoli and that the distribution is related to the wall of the terminal bronchiole, where the capillary anastomoses between bronchial and pulmonary arteries are believed to lie. At this site frequently repeated haemorrhages of mechanical origin occur. These are associated with haemoptysis. The ferrous impregnation of elastic tissue and reticulin around the haemosiderin causes the breakdown of these structures, with resulting fibrosis. This obstructs the lymphatic withdrawal and so perpetuates and intensifies the dangerous concentration of intra alveolar haemosiderin, and in some of these cases an osseous mass develops.

### Necropsy findings

Histologically the only difference between the two varieties of pulmonary

### Diagnosis

**Diagnosis**  
Skiagraphic appearances show the characteristic cardiac shadow of mitral stenosis. There are fine nodular opacities in the lungs, which are most numerous

## BIBLIOGRAPHY

in the mid zones and densest at the hila they produce a clouded or marbled effect. Towards the periphery and the bases they are fewer in number, and the apices may be clear.

### Prognosis and treatment

Haemosiderosis does not appear to affect the prognosis of mitral stenosis. Treatment should be directed to the treatment of the cardiac lesion, there is no specific treatment for the pulmonary disease.

## BIBLIOGRAPHY

- Borsos Nachtnebel O (1942) Fur pathologie der lungen hamosiderose *Zbl Allg Path path Aist* 79 174
- Coelen W Henke F and Lubarsch O (1931) In *Handb der spez path Anatomie* III 3 20
- Ellman P and Gee A (1951) Pulmonary Haemosiderosis *Brit med J* 2 384
- Gumpert T E (1947) Miliary Appearances in the Lungs in Mitral Stenosis Endogenous Pulmonary Haemosiderosis *Brit med J* 2 488
- King A B (1949) Pulmonary Haemosiderosis *Proc R Soc Med* 42, 87
- Lendrum A C Scott L D W and Park S D S (1950) Pulmonary Changes due to Cardiac Disease with Special Reference to Haemosiderosis *Quart J Med* 19 249
- Nancekivell L (1949) Acute Idiopathic Pulmonary Haemosiderosis *Brit med J* 1, 431
- Pendergrass E P Lane E L and Ostrum, H W (1949) Haemosiderosis of the Lung Due to Mitral Disease *Amer J Roentgenol* 61, 443
- Veränderungen  
38 353  
f W McLaren
- Walton M Williams A A Bodan M and Barlow A (1951) Idiopathic Pulmonary
- NY Acad Med* 18 93  
Idiopathic Haemosiderosis

## CHAPTER 23

### OCCUPATIONAL LUNG DISEASES

KENNETH M A PERRY

IT HAS been known since prehistoric times that dust will produce diseases of the lung, and conditions of this type were termed pneumoconiosis (Greek *pneumon* — lung, *konis* — dust). The term has since been incorrectly corrupted in England to pneumoconiosis, which translated literally means 'air dust'. The term is an all inclusive one but is usually adapted to refer to dusts which are either inert, such as iron, or give rise to a fibrosis of the lung as do dusts like silica, asbestos, certain other silicates and coal. The dust must be of a particle size of less than  $5\ \mu$  to pass through the smallest bronchioles and thus reach the alveoli and so give rise to pathological lesions. Besides these dusts certain metallic dusts such as manganese, vanadium, cadmium, osmium and beryllium give rise to either acute or chronic pulmonary oedema or pneumonia. Pollens and dusts of *bat and mouse* are also known to cause asthma which is described elsewhere.

the con

produce a characteristic asthma which will be referred to below. Organic dusts arising from hay, straw, grain, bagasse, and cotton produce a group of diseases which have been labelled 'asthma', 'bronchitis', and 'pneumonia'. These sometimes resolve leaving little disability but in other cases run a protracted course with fibrosis of the lung and bronchiectasis.

#### PLATINUM ASTHMA

As long ago as 1911 Karasek reported that workers in photographic studios in Chicago suffered from irritation of the throat and nasal passages, which caused violent sneezing and coughing. The bronchial irritation was sufficient to cause marked shortness of breath and a wheeze. It was found that the cause of this condition was the use of paper containing potassium chloro platinate, and when this paper was no longer used the symptoms subsided. Refining of platinum involves the production of the complex salts of platinum and at least half of the workers in this industry suffer from a syndrome similar to that produced by the photographic paper. The symptoms are repeated sneezing followed by profuse running of the nose with watery discharge. This is followed by tightness of the chest, shortness of breath and a wheeze. The symptoms persist as long as the worker remains in the factory and for about half an hour after he has left they then subside. However the patient may wake in the early hours of the morning.

# PLATINUM ASTHMA

DEATHS FROM FIBROSIS OF THE LUNG INCLUDING SILICOSIS BYSSINOSIS (ENGLAND AND WALES) 1940-1949  
(From the Annual Report of Chief Inspector of Factories 1949)

TABLE I

	1940	1941	1942	1943	1944	1945	1946	1947	1948	1949
<i>Silicosis</i>										
Refractories industries	16	12	7	7	6	9	8	10	13	13
Pottery manufacture of	53	45	47	41	32	41	49	54	48	63
Sandstone quarrying and dressing	42	24	14	24	26	21	20	12	28	15
Sandstone masons	67	58	40	53	31	44	41	43	40	36
Metal grinding	24	13	13	11	7	11	6	9	8	14
Sandblasting	13	4	4	6	7	6	4	5	5	5
Steel dressing and cleaning of castings	4	9	9	4	10	5	22	13	13	18
Stone pebble flint and sand crushing	5	1	2	1	4	—	3	1	3	—
Scouring powders manufacture of	—	2	—	1	1	—	—	—	—	—
Abrasive wheel manufacture	—	1	—	—	2	—	—	—	—	—
Glass cutting and bevelling	—	—	—	—	—	—	—	—	—	—
Millstone dressing	—	—	—	—	—	—	—	—	—	—
Slate quarrying and dressing	—	—	—	—	—	—	—	—	—	—
Granite quarrying and dressing	7	6	4	6	4	13	20	16	21	—
Tunnel mining (sewage works etc.)	1	1	3	2	4	5	3	3	2	—
Coalmining	2	—	—	—	—	—	—	—	—	—
Goldmining (South Africa)	232	196	230	276	323	343	347	322	336	—
Tin mining	10	7	7	9	4	7	5	4	4	—
Iron ore (haematite) mining	21	17	12	14	11	19	12	11	20	—
Lead mining	20	4	7	5	7	2	9	6	9	—
Copper mining	—	1	3	1	1	1	2	12	1	—
Barytes mining	—	—	—	—	—	—	—	—	—	—
Clay mining	3	3	2	1	1	1	2	12	9	—
Mining engineers	—	—	—	—	—	—	—	—	—	—
Miscellaneous	—	—	—	—	—	—	—	—	—	—
<i>Total—Silicosis</i>	2	1	2	4	5	6	11	7	7	—
<i>Asbestosis</i>	522	406	417	470	445	508	553	558	534	575
<i>Pneumoconiosis</i>	11	17	11	8	10	11	16	15	15	17
Coalmining	—	—	—	—	34	64	78	230	317	420
Other industries	—	—	—	—	3	11	5	12	19	14
<i>Byssinosis</i>	—	—	—	—	—	—	—	—	—	—
<i>Other causes of (non-occupational) Fibrosis</i>	—	—	6	7	1	10	3	4	8	7
<i>Grand Total</i>	650	461	443	493	531	529	568	616	651	554
	1 183	884	877	978	1 024	1 133	1 223	1 435	1 544	1 587

## OCCUPATIONAL LUNG DISEASES

dust of the complex salts of platinum, in a concentration of 5 micrograms per cubic metre or more. The disease can be prevented by adequate exhaust ventilation. The condition is not caused by spongy or metallic platinum even in concentrations as high as 1,000 milligrams per cubic metre, nor is it caused by the complex salts of the allied metals palladium, rhodium and ruthenium.

### ORGANIC VEGETABLE MATERIALS

There are about 800,000 persons employed in agriculture in Great Britain, and it is therefore one of the largest industries in the country. "Farmers' lung" was first described by Campbell (1932) and Fawcett (1938) in the north west areas, and by Fuller (1947) in Devonshire. It has principally occurred in hay makers and has been called "bronchitis feniiseciorum" but it has been described in all harvest workers. In Scandinavia, Tornell (1946) and Zettergren (1950) called it "threshers' lung". Hoffman (1946) reported the disease in the Canton of Appenzell in Switzerland during the harvest of 1945. There have been instances recorded where the infection has occurred in tea tasters. A similar disease occurs in horses and is known as "broken wind". Fawcett ascribes the disease to fungi but though he clearly establishes the disease as an entity, the mycotic aetiology is not by any means proved. There is no doubt that rotting vegetable material contains innumerable fungi, and if these organisms are in the dust which is inhaled they will also appear in the sputum. They are constantly present in all the dusts which give rise to these conditions, and though they may not be the immediate cause it is possible that by producing toxic substances they indirectly contribute to the aetiology of the condition.

Patients suffer from gradually increasing shortness of breath, which may get progressively worse over a period of months. They may develop an acute febrile respiratory illness with a cough, scanty frothy sputum, and occasional haemoptysis. Physical signs include patches of dullness and crepitations. The blood sedimentation rate is raised. Radiographic appearances in the earlier stages consist of a fine reticulation which is followed by a rather soft snowflake mottling, widely distributed throughout both lungs.

When the disease progresses to the chronic stage the patient will gradually become more short of breath, he will become cyanosed, develop clubbing of his fingers, and a cough with purulent sputum. X-ray examination at this stage will show an increase in the opacity of the mottling and an increase of the hilar and perihilar shadows, coalescent areas of fibrosis, marked restriction of the diaphragmatic movements and considerable areas of emphysema.

The patient eventually becomes emaciated and finally dies with right sided heart failure. If diagnosis is made early the condition responds to treatment with potassium iodide, but if it is undiagnosed it progresses and the patient becomes a chronic invalid with fibroid lungs. Necropsy shows voluminous lungs with emphysema and thickened pleurae with many bands of fibrous tissue traversing the lung substance.

### BAGASSE

Bagasse is sugar cane after the sugar has been extracted. It is used for making boarding for interior decoration and thermal insulation. One gramme of air borne

dust may contain up to 240 fungal spores. Amorphous silica forms roughly 1-2 per cent of the bagasse but quartz only 0.1 per cent with many of the quartz particles being 20-30 $\mu$  in size. It is therefore impossible that there should be any risk of silicosis. An account of the disease was first published by Castleden and Hamilton Paterson (1943) in England, and by Sodeman and Pullen (1943) in the United States, and it has been more fully described by Hunter and Perry (1946).

Workers exposed to bagasse dust for a period of 6 weeks or more develop an acute febrile illness with extreme shortness of breath, cough with scanty black stringy sputum, and occasional haemoptysis. Throughout the lungs there is dullness, diminished air entry, and many coarse crepitations. Skiagrams show miliary shadowing, which suggests an acute bronchiolitis (Fig 106). The acute illness carries a mortality rate of 10 per cent but the majority of patients make a complete recovery.

In some, however, it progresses to a chronic stage in which the patient becomes excessively short of breath and develops a cough with much sticky sputum. On examination these patients are orthopnoeic, cyanosed, and have clubbing of their fingers. They have all the physical signs of emphysema with many coarse crepitations. The chest X-ray shows many areas of consolidation. Necropsy, however, shows marked sub pleural thickening.

During the acute phase of the disease the patients are best treated with the usual methods used in the management of pneumonia, including sulphonamides and penicillin, and usually there is a marked response. The treatment of the chronic condition is symptomatic and follows the principles of treatment of chronic bronchitis and emphysema.

### COTTON (BYSSINOSIS)

Jackson as early as 1818 drew attention to the fact that those who worked in the cotton industry, particularly card room workers, suffered from a characteristic respiratory disease, it also occurs in those who work with flax. In 1862 Greenhow described the condition in a report to the Privy Council, and in 1932 the Home Office issued the Report of the Departmental Committee on Dust in Card Rooms in the Cotton Industry. This report collected the existing information about the conditions in the mills, types of machinery in use, and the nature of cotton dust. It included an account of the clinical examination and radiological findings in a group of workers at Ashton under-Lyne who suffered from the disease, and described the necropsy findings in a few cases. The disease was named byssinosis from the Greek word *βυσσος* meaning "fine linen". Foreign literature on the condition is scanty, but Zipperlen (1934) investigated patients suffering from asthma in the textile industry of Wurtemberg, and recorded that there were many with severe attacks in whom strong allergic reactions could be obtained.

OCCUPATIONAL LUNG DISEASES




FIG 106 — Skiagram showing acute bronchiolitis caused by bagasse



FIG 107 — Fibrosis of lung and bronchiol ectasis from bagasse

cotton industry. He paid particular attention to the pathogenesis of the disorder but this is still a matter for debate. The protein fraction of card room dust has been shown to be capable of producing inflammatory lesions in animals and normal men. The characteristic course of the disease is suggestive of an acquired allergic hypersensitivity, and there is some confirmation of this view in the finding of positive cutaneous reactions to the proteins in affected persons, but only rarely in unaffected persons, but the primarily toxic qualities of the protein prevent the condition from being regarded as a purely allergic one. Other factors that have been suggested as possibly concerned are a substance resembling histamine, and moulds. Histamine-like substances similar to those found in card room dust have however been found in other dusts not causing similar symptoms. Gill (1947) described the process of carding and the symptomatology of the disease.

Cotton is obtained chiefly from Egypt, the United States of America, and India, and it arrives at the mill in tightly compressed bales, which also contain much foreign material such as particles of leaf and seed coat, cotton hairs, fragments of mould, and fine sand. The raw cotton is therefore first cleaned, and the tightly compressed cotton is pulled off in layers and hand fed into the machine called the "hopper bale opener". It then passes into the blowing room where a series of machines, by means of beating actions and currents of compressed air, free it from most of the impurities. All these machines are enclosed in dust-proof covers, the machines, however, have to be cleaned, and this is a particularly dusty job for which the workers have to be suitably clothed and wear respirators. The clean cotton coming out of the blowing-room passes to the carding engines, and it is principally the strippers and grinders in charge of these engines who suffer from the disease. Usually each stripper and grinder is in charge of 16 carding engines, each engine is cleaned four times daily by a vacuum process, and twice weekly it is brush stripped. The latter process is necessary since the vacuum cleaning fails to clean sufficiently deeply. The evidence is strong that the dust liberated by the carding process in the immediate vicinity of the carding engine contains the injurious dust which causes byssinosis. The atmosphere is progressively safer the greater the distance from the carding machines.

The usual history is that after working for several years in the dusty atmosphere the man begins to sneeze and develops a dry and irritating cough with a tight feeling in the chest and restricted intake of breath. At first the trouble is temporary, passing off in 1-2 days, but it returns after a short absence from work such as a week-end. It has, in consequence, been called "Monday morning fever", "mill fever" and "factory fever". Disablement and incapacity for work does not result at this stage, and the breathing will become normal on removal from the dusty atmosphere. After exposure to the dust for 10 or more years, however, the symptoms become more severe and persistent, and the worker suffers from asthma and bronchitis associated with shallow breathing, cough, and a small amount of sticky, mucoid phlegm. He loses weight and develops a sallow complexion. Absence from work and incapacity increase, although removal from the dust causes improvement, with capacity to work in another atmosphere. In the last stage the emphysema is severe. Cough is present, with mucoid or muco-



purulent expectoration and great shortness of breath. In extreme cases the man may stand with his hands pressed against his thighs to aid his breathing. The disease is incurable at this stage and many of these workers die before the age of 50.

The physical signs are those of chronic bronchitis and emphysema, the respiration is laboured, with accessory muscles called more and more into use. The thorax becomes typically barrel shaped, and the patient tends to sit in a characteristic attitude with the body bent slightly forward, the legs widely spaced, and the hands placed on the knees. The vital capacity is greatly reduced and the

suggest considerable fibrosis (Fig. 108).

Necropsies have revealed chronic bronchitis and emphysema but do not show any specific feature which enables the bronchitis of card room operatives to be distinguished from that which occurs in the general population. Dunn and Sheehan (1932) have reported that dilatation of the right heart was frequent.

No special medicinal treatment is of any value, and timely removal from the work is the only successful method of preventing the disease from progressing. Respirators have been tried but are not satisfactory since the men find them difficult to work in and therefore do not wear them. Methods of suppression of dust by exhaust ventilation and enclosing the machinery have been highly developed in the industry and the disease is thus kept to a minimum. It is still however one of the most important industrial diseases at the present time.

First Collis (1914) and then Middleton (1926) recorded that cotton weavers in Lancashire suffered from an acute bronchitis from handling cotton fibre, which was termed 'weavers' cough'. This was an acute illness with fever, cough and sputum, and residual symptoms, which might last up to two months. The most probable cause of the outbreaks was thought to be the amount of moisture left in the warp, producing an abundant growth of fungi which developed during storage on the beams and produced, on weaving, a dust laden with conidia spores and fragments of mycelium which were breathed by the weavers.

In the United States of America Neal, Schneider and Caminita (1942) described an acute febrile illness with cough, dyspnoea, tightness of the chest and more general symptoms occurring among workers making mattresses from low grade

plants where workers were  
They thought the condition

*Aerobacter cloacae*, or its  
this may be so but cotton  
possibly be

## CARCINOMA OF THE LUNG

Agricola and Paracelsus in the sixteenth century described at length a disease of the lungs with a high mortality rate in the miners of the Erzgebirge in Saxony. The condition was known as "Bergkrankheit" (mountain disease). It was not until



FIG 108 — Severe  
byssinosis in a man  
aged 50 years

1879 that it was identified by Hærtling and Hesse as cancer of the lung. These authors attributed 75 per cent of all deaths of miners to the disease, which usually developed after a period of 20 years' work in the mines. Arnstein (1913) distinguished the tumours as carcinoma, and differentiated them from lymphosarcoma and endothelioma, as most neoplasms of lung at this time were thought to be. Since people in the district other than miners were only rarely affected by this disease, all the early workers attributed the condition to the occupation of mining. The main ores in Schneeberg are sulphides and arsenides of nickel and cobalt — "Speiskobalt" ( $\text{CoAs}_2$ ) "Wassnickelkies" ( $\text{Ni}_2\text{As}_2$ ) and "Rotnickelkies" ( $\text{NiAs}$ ) — while at Joachimstal on the southern side of the Erzgebirge there is a high content of pitchblende and radium. Rostoski, Saupe, and Schmorl (1926) showed that the bore dust of these mines contained up to 0.45 per cent of arsenic, while the dust in the pits contained 0.1 per cent, but it has not been found possible to prove that arsenic is either the cause or a contributory agent. In Southern Sweden, where Speiskobalt is also mined, pulmonary cancer is said to be unknown, and in Schneeberg no other signs of arsenic poisoning are reported. Hueck (1939) has pointed out that multiplicity of tumours which is so frequent in arsenical cancer does not occur in Schneeberg miners, while estimations of arsenic in the urine, hair, and nails (Rostoski, Saupe and Schmorl, 1926), and in the lungs (Pirchan and Siki, 1932, Zeil, 1935), give equivocal results. Work in the Joachimstal area by Loewy (1929) and Siki (1930) has suggested that the radioactivity of the ores is the principal aetiological agent. Peller (1939) showed, from



FIG. 109.—Man, aged 52 years, nickel worker for 23 years. Squamous cell carcinoma of right upper lobe.

necropsies that 50 per cent of the miners in this area died with lung carcinoma while Lange (1935) gave similar figures for Schneeberg. Recently Lorenz (1944) expressed the opinion that the radio activity could not be the sole cause of the carcinoma since similar doses of x rays or  $\gamma$  rays do not produce lung tumours in animals. He suggested that both arsenic and radio activity might be contributory factors.

Amor (1938) reported 10 cases of carcinoma of the lung (Fig. 109) and numerous cases of carcinoma of the nose and nasopharynx among workers exposed to arsenical dust for an average period of 6 years at a nickel works in South Wales. The ores were from Sudbury in Canada and contained 2 per cent of arsenic and for a period of 25 years had had the copper extracted by the addition of sulphuric acid. The workers had been exposed to arsenic dust and fumes and to the fumes of sulphuric acid.

pneumonectomy for carcinoma of the lung on 2 of these workers. Both were squamous cell carcinomas. The arsenic contents of the lungs were 3.82 and 3.40 parts per million (p.p.m.) the content of the hair of the men was 2.2 and 10.5 p.p.m. and they were excreting 41.5 and 39.8 micrograms per 24 hours.

Other evidence suggesting that arsenic produces carcinoma of the lung is the experimental work of Leitch (1922) who has succeeded in producing such growths in mice with potassium arsenite. But the most conclusive evidence is the statistical work of Bradford Hill and others (1948) which showed that in one town in England

where the principal industrial work is the manufacture of arsenical sheep-dip the mortality rate for carcinoma in the chemical workers handling arsenic is double that in the rest of the population of the town, a statistically significant difference. It was also shown that the difference is made up almost completely by carcinomas of the skin and bronchi.

It is possible that the high content of arsenic dust in cigarette fumes may account for the world-wide increase in the mortality rate for carcinoma of the lung in males, which has risen six-fold in Great Britain in the past 10 years.

Machle and Gregorius (1948) studied the mortality data for 7 chromate plants in the United States of America employing 1,445 workers. They found 42 deaths from cancer of the respiratory system among a total of 193 deaths over periods varying with the different plants. Of the 193 deaths, 66 were from cancer, thus the respiratory cancers were 63.6 per cent of all cancers, compared with 8.7 per cent in a control group of 733 deaths among Metropolitan Life Insurance Company industrial policy holders in the year 1946. Of these, 115 were from cancer, of which 10 were of the respiratory system. The mortality rate for lung cancers in the chromate workers over 50 years of age was up to forty times that for a comparable industrial group. In one plant exposure was restricted to bichromates and chromic acid, and here there was no cancer of the respiratory system, this evidence strongly suggested that the carcinogenic substances were the monochromates. This also means that nasal irritation and septal perforation may occur without associated high rates for cancer of the lung. The mean duration of exposure was 14.5 years. The disease is further reviewed by Baetjer (1950).

Wyers (1950) has shown that there is an excessive mortality from carcinoma of the lung in cases of asbestosis. In his series of 115 deaths from asbestosis 11

the more striking because 41.2 per cent of the deaths in his series were females, whereas the incidence of primary cancer of the lung in the general population is 4.4 males to 1 female.

Workers with pitch and coal tar derivatives may also develop squamous cell carcinoma of the bronchus as well as epitheliomas of the skin.

### SILICATES

Most silicates will produce dust, and if workers are exposed to these dusts they

#### China-clay or kaolin

This is hydrated silicate of aluminium, and at present is one of the most important exports from the United Kingdom. It is an important ingredient in china and earthenware pottery, and is used in the manufacture of a large number of other commodities, including cotton cloth, paint, soap, and fine art paper. It

## OCCUPATIONAL LUNG DISEASES

is formed by the decomposition of feldspar by the removal from it of silicate of potassium through the action of water containing carbon dioxide

The largest production of china clay in the world is in Cornwall, where it occurs in deep pits from which it is washed with water under pressure. Men are exposed to the dust in shovelling the clay after it has been dried in the kilns. No medical survey has been carried out in the Cornish china clay industry, and therefore little is known of the effect of this dust on the lungs. Middleton (1936), however, recorded the case of a worker aged 49 years whose skiagram showed the whole of both lung fields covered with extremely fine mottling, most marked in the middle zones and at the right base. He considered the film suggested pneumoconiosis, not of a silicotic type. Hale (1946) has some radiographs of workers in the industry showing well-marked reticulation. It is probable that these workers have few symptoms and that the dust does not produce great, if any, disability.

Thomas (1952) described in the findings in two necropsies: one a worker in the ball clay industry and one who had been engaged on grinding china stone, a granite containing china clay. He found in these workers evidence of silicosis.

### Talc or French chalk

This is hydrated magnesium silicate, and in mineral form is known as steatite or soapstone. It is used as a filler for paper, soaps, and paints, for leather dressing, in the manufacture of rubber motor tyres, in making electric switchboards, and for heat insulation of steam pipes.

The first account of changes in the lungs due to talc was given by Thorel (1896) followed by a report by Cesa-bianchi and Devoto (1911) of the necropsy of a woman who had worked in talc. The lung showed much talc dust but little resulting fibrosis. Zanelli (1931) recorded the case of a woman worker, aged 27 years, in a tyre factory, who had early tuberculosis but also reticulation in the lung radiograph. Dreessen (1933) investigated the workers in a tremolite talc mine and mull and found that those who had been employed for more than 5 years had radiographs which showed fine diffuse bilateral fibrosis but this appeared to cause little disability. Dreessen and Dalla Valle (1935) carried out a further survey in Georgia and found that 16 out of 33 mill workers exposed to high concentrations of talc and 6 of 13 miners had pneumoconiosis, in 4 cases there was evidence of added pulmonary infection. Merewether (1933, 1934) examined 24 workers who were exposed to talc in the manufacture of rubber tyres for periods of between 4½ months and 40 years. They showed no disablement though radiographic changes suggested diffuse interstitial fibrosis of the lungs (Fig. 110). Siegal, Smith and Grinnell (1934) examined 221 men and women talc millers and miners in St. Lawrence County, New York, known as asbestine.

The free silica content of the asbestine varies from 0.5 to 1.0 per cent. It contains fine, straight, needle-like fibres, and dust counts ranged from 6 million to 5,000 million particles per cubic foot. This material was obtained by underground mining with inclined shafts to depths as low as 1,000 feet; each mine employed 15-17 men underground. 221 miners and millers were examined and radiographic evidence of fibrosis was found in 32, giving an incidence of 14.5 per



FIG. 110 — Woman aged 48 years exposed for 20 years to talc in a rubber tyre factory. She complained of severe cough and dyspnoea

cent. All the cases occurred in men who had worked for 10 years or longer, and the incidence in this group was 25.9 per cent. The appearances were of fibrosis of a fine, diffuse type, showing granulation or nodulation on a hazy background. The resulting dyspnoea, cough and fatigue were disabling, and many of the men showed clubbing of the fingers. There was some increased susceptibility to tuberculosis, 3 men showing evidence of the disease. Besides the fibrosis, deposits of unidentified opaque material in the periphery of the lung called 'talc plaques' were observed in 6.3 per cent of the workers. These plaques varied from single linear deposits in the region of the diaphragm to massive deposits, bizarre in shape, extending over a large part of the lung fields; they were irregular in shape, density, and the sharpness of their outlines. No deaths were recorded, but the clinical and radiographic picture showed resemblance to asbestosis rather than silicosis. McLaughlin, Rogers and Dunham (1949) recorded the results obtained at necropsy of a man aged 51 who had worked for 37 years in a rubber tyre factory, where he was exposed to talc dust. A feature of the histology was the presence of bodies similar to asbestos bodies found in a pneumoconiotic lung. Although there was much fibrosis of the lung and abundant doubly refractile particles, x-ray analysis of the lung suggested that there was less than 0.06 per cent of free silica in the lung. It seems clear that talc and asbestos give rise to similar diseases.

#### Asbestos

The term 'asbestos' is of Greek origin and means "unconsumable". Asbestos

## OCCUPATIONAL LUNG DISEASES

was worked into textiles by the Egyptians, Romans, and the inhabitants of the ancient empire of Tartary, the wealthiest among them using it as cremation cloth. Pliny called it a "rare and costly linen," whilst Plutarch recalls its use as the perpetual wick in the lamps of the Vestal Virgins. The use of asbestos seems to have been lost in the Middle Ages and an attempt to re-introduce it in the 17th century met with little success, so that it again became indus-

is concerned with two main types of asbestos: (1) Hornblende or amphibole, to which group blue crocidolite, readily identified by its lavender-blue colour, and amosite, white or yellowish brown in colour, belong, is produced exclusively in South Africa; (2) Chrysotile, white in colour and of fine texture, is produced mainly in Canada, Rhodesia, Transvaal, Australia, Italy and Russia.

TABLE II  
TABLE OF COMPOSITION

	Hornblende		Chrysotile	
	Blue crocidolite, percentage	Amosite, percentage	Canadian, percentage	Rhodesian percentage
Silica	50	47	41	41
Alumina	1	6	3	1
Ferric oxide	35	37	2	4
Magnesia	2	6	40	39
Alkalines	8	1	—	—
Water	4	3	14	15

It is imported as the crushed rock, but the fibres can be separated to an extreme degree of fineness which renders them capable of being carded, spun into yarn and woven into cloth by processes similar to those used in the cotton industry. Besides being woven, asbestos is ground and mixed with cement and plastics to make insulating slabs, boarding, and many other articles used in industry. The dust given off during the process of manufacture is mostly less than  $5\mu$  in diameter and  $10\mu$  in length.

The first case of asbestosis was described by Murray (1907). The man had been employed on a carding machine for 14 years and was the last survivor of 10 men who had worked in the card room, the others all having died about the age of 30 with respiratory diseases. He died in Charing Cross Hospital in 1900 at the age of 34. Necropsy revealed extensive diffuse pulmonary fibrosis and the lungs were found to contain asbestos bodies. In 1906, Marchand and Reiss (Kruger, Rosloski and Saupe, 1931) noted the presence of unusual bodies in the lungs of an asbestos worker, while Auribault (1906) in France recorded the high mortality in an asbestos textile factory in Hamburg. Real attention, however, was not given until the death of an asbestos worker was recorded, who had pulmonary tuberculosis, while McDonald (1927) and Cooke and Hill (1927)

gave the first detailed descriptions of asbestos bodies and asserted that they originated in the inhaled dust. Seiler (1928), and with Gilman in 1931, recorded the necropsy findings in an asbestos worker who showed fibrosis but no evidence of tubercle. The condition was fully described by Merewether and Price (1930) and Merewether (1933, 1934). Clinically, the striking symptom is dyspnoea of gradually increasing intensity, often associated with a dry cough, in the early stages physical signs are slight or absent, but later râles throughout the lung develop and the fingers often become clubbed. Radiologically the most characteristic feature is the "shaggy" outline of the heart shadow, while in the later stages the lung fields are opaque, and the diaphragm is obscured, which is characteristic of silicosis.

At necropsy, pleural adhesions are extensive and dense, and there are often thick subpleural plaques of fibrous tissue. The consistency of the whole lung is increased, and there may be large areas of fibrous condensation which is tough rather than stony hard as in silicosis. Other parts of the lung, not affected by the fibrotic change, are congested and consolidated with inflammatory processes (Figs 112 a and b). Emphysema is extensive, but localized often to the lower and apical parts of the lung, pneumothorax sometimes occurs. The histology has

Wood and Gloyne (1934) collected 100 cases of the disease and found that 21 were complicated by pulmonary tuberculosis, but Middleton (1936) studied 54 deaths occurring between 1930 and 1934 and found that tuberculosis was present in 24 (44 per cent). The periods of employment in the industry ranged from 10 months in the disintegrating plant to 23 years' weaving, the average duration being just under 10 years. He suggested that tuberculosis was a slightly less frequent complication than in silicosis and that it is more frequent as a terminal complication than as a concomitant disease during the earlier stages.

Wyers (1947) showed that, as a result of measures enacted by the legislation of 1931, the morbidity rate from asbestosis fell from 39.1 per 1,000 in 1932 to zero in 1940.

the disease for several years (15.2 per cent in 1932, 11.2 per cent in 1933, 10.1 per cent in 1934, 10.1 per cent in 1935, 10.1 per cent in 1936, 10.1 per cent in 1937, 10.1 per cent in 1938, 10.1 per cent in 1939, 10.1 per cent in 1940).

and Wedekind (1947) have shown that the characteristic of this disease, has now largely given place to a coarser, more granular appearance. Pathologically, confluence of the blue polygonal areas is not so






FIG. 111—Man aged 56 years employed for 6 years carding in an asbestos factory. He had gross clubbing of fingers and dyspnoea.

advanced as that seen in the specimens of 15 years ago. Wyers thought these changes were due to longer exposures to lower concentrations of dust.

From his clinical studies of asbestosis Wyers concluded that it differed from silicosis in being a clinical rather than a radiological disease. Of the clinical signs he stressed the importance of clubbing of the fingers and symptoms of cardiovascular origin (raised diastolic blood pressure and accentuated second pulmonary sound) as being always present when once established, whereas the marginal crepitations and fibrotic crackles tend to be evanescent. Of the complications emphysema was always present, bronchopneumonia the commonest termination, tuberculosis was of the acute caseous type and occurred in one third of the cases, the right heart dilatation was common and bronchiectasis rare. Pregnancy could not be shown to affect the progress of the disease adversely and the obstetrician should be guided by cardiovascular rather than the pulmonary signs.

The difference between the reaction of the tissue to silica and to asbestos is even more marked than the difference between the clinical pictures of silicosis and

dust does not cause an acute inflammatory reaction. Gardner and Cummings (1931) found that asbestos accumulated in the bronchioles and then in the alveoli and fibrosis first appeared after about 500 days exposure. Gardner (1942) put



(a)



(b)

FIG 112—(a) Lung of an asbestos spinner (a patient of Dr H Wyers) (b) lung of asbestos worker showing gross thickening of pleura (a patient of Dr H Wyers) (c) asbestos bodies in sputum



(c)

forward the 'mechanical' theory as an explanation of the disease. It was the long fibre in the dust which caused asbestosis, and if the asbestos were ground sufficiently finely no disease resulted. He concluded that the disease results from the mechanical irritation produced by a combination of the long fibre and the rhythmic movement of the lung. He also showed that different pathogenic responses resulted in different animals. King, Clegg and Rae (1946), however, produced somewhat different results from the intratracheal injection of asbestos dust into rabbits: they reported that the injection of long fibres ( $15\mu$ ) produced a nodular reticulosis comparable with the experimental silicotic nodule, whereas short fibres ( $2.5\mu$ ) produced a diffuse interstitial reticulosis.

**Silimanite**

Silimanite is an aluminium silicate, and is a dense rock found in India. It is used as a refractory material in the manufacture of porcelain. When broken it produces masses of prismatic crystals in the form of fibres which cannot be broken down into fine fibres like asbestos but tend to break and become reduced to a powder.

When handled in industry the rock is crushed, ground and sieved, and then calcined, during which a considerable quantity of dust is given off. Middleton (1936) reported the investigation of 15 men who were employed handling this material. Their ages ranged from 24 to 50 years and the duration of their employment in the industry was from  $1\frac{1}{2}$  to 16 years. Chest radiographs showed abnormality in only 4 men, who had been working with the material 16, 11, 7 and 5 years, and in whom some reticulation was noted. These numbers are small, but

$5\mu$  in diameter. Necropsies were carried out on 22 of these animals. Five died from collapse or pneumonia after being dusted for periods of up to 100 days, and showed minimal fibrosis; 14 survived for periods up to a year, and in this group there was more fibrosis with thickening of the inter-lobular septa. Three of the animals survived 673 days; these showed well defined dust nodules in the inter-lobular septa and in the perivascular and peribronchial regions, consisting of dust particles with fibroblast and early connective tissue formation. These animal experiments suggest, therefore, that the dust may not be quite as harmless as the clinical observations of Middleton suggest.

**Mica**

Mica belongs to a group of widely distributed rock-forming minerals, of which muscovite and phlogopite are important commercially. Muscovite is a silicate of aluminium with potassium and phlogopite is magnesium-aluminium silicate. The former is used for its transparency and the resistance to fire for windows of stoves and lanterns while it is sometimes ground up and used in the manufacture of paper, wall-paper, and paint, and as a lubricant and absorbent. The latter is used exclusively for electrical purposes. Dust is produced when the materials are being ground and finished.

Middleton (1936) reports that Ferguson examined 12 workers who were exposed to mica dust, but only 5 of them had been exposed for more than 5 years and these complained of cough and shortness of breath. Radiographs showed some fibrosis of the peribronchial type with some diffuse shadows in the middle zone. Ferguson thought, therefore, that the dust of mica was probably capable of causing pulmonary fibrosis.

**Diatomaceous earth and tripoli**

Diatomaceous earth and tripoli are hydrated, amorphous minerals of organic origin, which are generally considered to be varieties of opal. Diatomaceous earth is also known as diatomite, kieselguhr and ferina fossil. It is found in the lake deposits of mineral diatomaceous shells. It forms a porous mass sometimes

## SILICATES

loose and floury, sometimes compact and clay like, and is always light-coloured, either white, grey or yellow. Tripoli, on the other hand, is a marine sediment. It generally contains well preserved fossils of fish and shell fish. After quarrying, draining and pulverizing, diatomaceous earth is marketed for a variety of industrial purposes including the manufacture of refractory bricks, and filters for drinking water, wines or syrups. It is used as an insulating material. Tripoli has fewer uses, but it is used for polishing metals and precious stones. Fuller's earth is also a mineral substance containing silica. Legge and Rosenkranz (1932) recorded the




FIG. 113.—A silicotic nodule

Investigation of 118 workers in a diatomite quarry in Santa Barbara, California, 81 of the patients showed radiological abnormality and in these the changes were advanced. The patients showed clubbing of the fingers and complained of shortness of breath. More recently Martin, Valland and Champey, in 1947, examined 32 workers exposed for many years to heavy concentrations of diatomite dust, but only 3 showed x ray appearances of mild interstitial pneumoconiosis, and of these only one had any disability. They concluded that while diatomaceous earth produces radiological and clinical disease it was of a comparatively mild variety. Vigliani and Mottura (1948) recorded 8 cases in which a severe pneumoconiosis developed after as little as 5 years exposure to the dust. A peculiarity of these cases was the tendency to spontaneous pneumothorax. Cases of pneumoconiosis resulting from long exposure to Fuller's earth have been recorded by Middleton (1938) in which Gloyne showed that in the lower part of the lung there were extensive patches of palpable dust fibrosis.

## SILICOSIS

Silicosis is a fibrosis producing lung disease caused by the inhalation of free, crystalline silica or quartz particles. It is the most disabling of all the dust diseases. It was originally thought that the disease was caused by the mechanical action

## OCCUPATIONAL LUNG DISEASES

of the sharp particles of quartz, but Gardner (1932) showed, by animal experiments, that silicon carbide (carborundum) the particles of which are as sharp as those of quartz, does not produce fibrosis to the same degree. These experiments resulted in the solubility theory of silicosis. Quartz is taken up by the phagocytes

TABLE III  
FATAL CASES INVESTIGATED UP TO THE END OF 1949  
(From the Annual Report of Chief Inspector of Factories 1949)

	Number of deaths	Average age at death	Duration of employment in years		
			Longest	Shortest	Average
SILICOSIS					
<i>Pottery</i>					
Silicosis	579	61.2	62.0	2.8	38.8
Silicosis with tuberculosis	423	55.5	67.0	5.0	34.6
<i>Sandstone</i>					
Silicosis	206	59.2	60.0	9.0	39.4
Silicosis with tuberculosis	209	56.7	55.0	5.0	37.2
<i>Grinding of metals</i>					
Silicosis	102	58.7	56.0	14.0	36.4
Silicosis with tuberculosis	196	53.7	53.0	2.8	32.9
<i>Sandblasting</i>					
Silicosis	61	48.7	28.0	1.7	12.3
Silicosis with tuberculosis	96	46.1	46.0	2.0	13.1
<i>Manufacture of scouring powders</i>					
Silicosis	13	37.2	37.0	2.3	8.2
Silicosis with tuberculosis	6	40.8	11.2	2.0	7.0
<i>Miscellaneous</i>					
Silicosis	211	54.6	57.0	1.5	23.3
Silicosis with tuberculosis	206	51.2	50.0	0.7	25.2
<i>Total</i>					
Silicosis	1 172	58.5	62.0	1.5	34.2
Silicosis with tuberculosis	1 136	53.8	67.0	0.7	31.1
ASBESTOS					
<i>Asbestosis</i>	186	48.2	48.0	0.5	15.6
<i>Asbestosis with tuberculosis</i>	78	39.2	29.0	0.8	10.5

in which it goes into solution. So long as it is contained within the phagocytes it is harmless but when it is released as a result of infection, either tuberculous or non-tuberculous, the typical whorled fibrous nodule of silicosis results (Fig. 113). This stimulated the work of Kettle (1932) who showed that by mixing iron oxide with silica the amount of the fibrosis that was produced in the lung was reduced. Later Denny, Robson and Irwin, in 1938, showed that aluminium-coated quartz particles did not go into solution. This led them to use powdered aluminium for the prevention and treatment of silicosis, fine black metallic aluminium powder with particles of less than  $2\mu$  being blown into the workers' changing rooms before they descend the mine. The solubility theory is, however, far from satisfactory since silicic acid normally appears as an inert substance. It seems probable there

## SILICOSIS

fore that the disease is brought about by some physico-chemical process at present not fully understood (Heffernan, 1948). It is clear, however, from these observations that the effect of silica on the lungs is greatly modified by mixture with other dusts. In industry silica rarely occurs in a pure form, and the type of disease that results from exposure to it is greatly modified by the substances with which it is mixed, as is well shown by Gloyne (1951) in his review of the pathology of 1,036 pneumoconiotic lungs. Pure silicosis occurs in workers engaged in metalliferous mining granite quarrying, metal grinding where sand stone is used, and among sand stone masons. The most rapidly developing variety of silicosis that has been described was that encountered in workers engaged in the manufacture of abrasive soaps and scouring powders, where it is possible that the presence of an alkali accelerates the action of the silica (Macdonald, Piggott and Gilder, 1930). Silicosis also occurs in the workers in the underground slate quarries in North Wales, but it has not been reported from the ancient quarry in Cornwall. This condition has been called schistosus.

### Pathology

The classical pathological picture of silicosis is that of a discrete nodular fibrosis (Fig 114). The fibroblasts forming the nodule have a typical laminated appearance, which causes the structure of the nodule to resemble the cut surface of an onion. The nodules vary in size from 0.5 centimetre to 1.0 centimetre in diameter, but they can be seen and palpated when they are only from 1 millimetre to 2 millimetres across. These are characteristically of smooth black or marbled grey black colour and may be divided into three groups. Minimal silicosis in which the whorled nodules can be detected only under the microscope, nodular silicosis in which the nodules stand out from the cut surface of the lung, most numerous in the upper lobe and with a tendency to occupy the peripheral part of the lung, thus showing a pleural drift. And lastly massive silicosis where, as the disease progresses, the nodules in some areas generally in the upper part of the lung, become more numerous and tend to conglomerate to form large hard masses. The centre of the nodules tends to undergo a degenerative process and therefore to lose the characteristics seen in the early stage. The surrounding lung tissue in classical uncomplicated silicosis shows little evidence of interstitial fibrosis. The



Fig 114 — Macroscopic appearances of  
Fig 121 African goldminer after 20  
years work showing typical whorled  
nodules and the conglomerate area  
at the apex (Section cut by Professor  
Jeihro Gough)

at the root of the lung undergo a process of fibrosis. They are hard and are usually deeply pigmented and histologically show areas in which the fibroblasts are arranged in a whorled manner. This undergoes a fibrotic change resulting in thickening and the formation of nodules. This may result from the deposition of silica in the subpleural space as a result of a retrograde lymphatic flow.

The patient must have been exposed to silica dust for a period of probably more than 10 years. This may have occurred in some early part of his life, so that a complete occupational history must be obtained. He complains of increasing breathlessness on exertion, and this may be exaggerated by a super added chronic bronchial infection, which may give rise to cough and sticky sputum. There is tightness of the chest and a wheeze. Not only does tuberculosis predispose to the development of silicosis as has been recorded above but silicosis predisposes to the development of tuberculosis. The symptoms of this disease may then be added to the above. On examination the chest will show evidence of emphysema, it will be barrel shaped, there will be diminished costal and diaphragmatic movements. There may be hyperinflation of the fingers, rhonchi will be heard scattered throughout the lungs. The chest expansion may be reduced to less than an inch. The blood sedimentation rate will be increased if infection is present and it would appear that infection and silicosis are closely allied. The vital capacity and maximum breathing capacity are reduced, the residual air increased, the blood oxygen tension is reduced. Skiagrams of the chest will show reticulation, nodulation and in many cases massive shadows, which are caused by the conglomeration of the whorled areas of fibrosis. Radiologically, similar shadows may arise in diseases such as primary tuberculosis, sarcoidosis or haemosiderosis. It is not possible to differentiate these various diseases by examination of the skiagram alone. A complete clinical and occupational history is essential for the diagnosis of silicosis (Figs 115-121).

### Treatment

Once the disease is established treatment can only be symptomatic. If the patient is freed from contact with silica before massive shadows have developed the disease will not progress, but if massive shadows are present it may continue to progress even though he is no longer in contact with silica dust. Many workers hold the view that massive shadows are a manifestation of tuberculous infection even though it is usually impossible to isolate the tubercle bacillus. Gloyne (1951) reported that in 1,036 pneumoconiotic lungs which he examined, he found evidence of pulmonary tuberculosis in 43 per cent. He found primary cancer of the lung in 7.7 per cent. Apart from removing the man from contact with the silica, treatment should be directed to treating infection. There is no scientific evidence that treatment by inhalation of aluminium powder has any therapeutic effect in relieving symptoms or preventing the disease, though workers and patients usually express satisfaction with the results of the treatment.

### Prevention

Prevention is the most important form of therapy. Managements should be

# SILICOSIS

FIG 115 —Silicosis in  
an iron moulder  
Reticulation



FIG 116 —Silicosis in a  
steel fettler Nodu-  
lation





lymphatic glands at the root of the lung undergo a process of fibrosis. They become small and hard and are usually deeply pigmented and histologically show a process of fibrosis in which the fibroblasts are arranged in a whorled manner. The pleura also undergoes a fibrotic change resulting in thickening and the formation of adhesions. This may result from the deposition of silica in the subpleural spaces as a result of a retrograde lymphatic flow.

### Clinical course

The patient must have been exposed to silica dust for a period of probably more than 5 years. This may have occurred in some early part of his life, so that a complete occupational history must be obtained. He complains of increasing shortness of breath on exertion, and this may be exaggerated by a super added catarrhal bronchial infection, which may give rise to cough and sticky sputum and later to tightness of the chest and a wheeze. Not only does tuberculosis predispose to silicosis as has been shown, but the development of tuberculosis is also influenced by the development of silicosis.

On examination the chest is flattened and has diminished costal and diaphragmatic movements. There may be clubbing of the fingers, rhonchi will be heard scattered throughout the lungs, the chest expansion may be reduced to less than an inch. The blood sedimentation rate will be increased if infection is present and it would appear that infection and disability are closely allied. The vital capacity and maximum breathing capacity are reduced, the residual air increased, the blood-oxygen tension is reduced. Skiagrams of the chest will show reticulation, nodulation and in many cases massive shadows, which are caused by the conglomeration of the whorled areas of fibrosis. Radiologically, similar shadows may arise in diseases such as miliary tuberculosis, sarcoidosis or haemosiderosis. It is not possible to differentiate these various diseases by examination of the skiagram alone. A complete clinical and occupational history is essential for the diagnosis of silicosis (Figs 115-121).

### Treatment

Once the disease is established treatment can only be symptomatic. If the patient is freed from contact with silica before massive shadows have developed the disease will not progress, but if massive shadows are present it may continue to progress even though he is no longer in contact with silica dust. Many workers hold the view that massive shadows are a manifestation of tuberculous infection even though it is usually impossible to isolate the tubercle bacillus. Gloyne (1951) reported that in 1,036 pneumoconiotic lungs which he examined, he found evidence of pulmonary tuberculosis in 43 per cent. He found primary cancer of the lung in 7.7 per cent. Apart from removing the man from contact with the silica, treatment should be directed to treating infection. There is no scientific evidence that treatment by inhalation of aluminium powder has any therapeutic effect in relieving symptoms or preventing the disease, though workers and patients usually express satisfaction with the results of the treatment.

### Prevention

Prevention is the most important form of therapy. Managements should be

# SILICOSIS

FIG 115 —Sil costs in  
an iron moulder  
Ret culation



FIG 116 —Sil costs in a  
steel fettler Nodu  
lation



## OCCUPATIONAL LUNG DISEASES

lymphatic glands at the root of the lung undergo a process of fibrosis. They become small and hard and are usually deeply pigmented and histologically show a process of fibrosis in which the fibroblasts are arranged in a whorled manner. The pleura also undergoes a fibrotic change resulting in thickening and the formation of adhesions. This may result from the deposition of silica in the subpleural lymphatics as a result of a retrograde lymphatic flow.

### Clinical course

The patient must have been exposed to silica dust for a period of probably more than 5 years. This may have occurred in some early part of his life, so that a complete occupational history must be obtained. He complains of increasing shortness of breath on exertion and this may be exaggerated by a super added catarrhal bronchial infection which may give rise to cough and sticky sputum, and later to tightness of the chest and a wheeze. Not only does tuberculosis predispose to tuberculosis as has been recorded above but silicosis predisposes to the development of tuberculosis. The symptoms of this disease may then be added.

On examination the chest will show evidence of emphysema, it will be barrel shaped and have diminished costal and diaphragmatic movements. There may be clubbing of the fingers. Rhonchi will be heard scattered throughout the lungs, the chest expansion may be reduced to less than an inch. The blood sedimentation rate will be increased if infection is present and it would appear that infection and disability are closely allied. The vital capacity and maximum breathing capacity are reduced. The residual air increased, the blood oxygen tension is reduced. Skilagrams of the chest will show reticulation, nodulation and in many cases massive shadows, which are caused by the conglomeration of the whorled areas of fibrosis. Radiologically, similar shadows may arise in diseases such as miliary tuberculosis, sarcoidosis or haemosiderosis. It is not possible to differentiate these various diseases by examination of the skilogram alone. A complete clinical and occupational history is essential for the diagnosis of silicosis (Ligs 115-121).

### Treatment

Once the disease is established treatment can only be symptomatic. If the patient is freed from contact with silica before massive shadows have developed the disease will not progress but if massive shadows are present it may continue to progress even though he is no longer in contact with silica dust. Many workers hold the view that massive shadows are a manifestation of tuberculous infection even though it is usually impossible to isolate the tubercle bacillus. Gloyne (1951) reported that in 1936 pneumoconiosis in 43 per cent. He found primary cancer of the lung in 7.7 per cent. Apart from removing the man from contact with the dust, treatment should be directed to treating infection. There is no scientific evidence that treatment by inhalation of aluminium powder has any therapeutic value in relieving symptoms or preventing the disease, though workers and their usually express satisfaction with the results of the treatment.

Prevention is the most important form of therapy.

FIG 115—Silicosis in  
an iron moulder  
Ret iculat ion



FIG 116—Silicosis in a  
steel fitter. Nodu  
lat ion



# OCCUPATIONAL LUNG DISEASES



FIG 117—Reticu-  
lation and mas-  
sive shadows in a  
brass fitter



FIG 118—Reticu-  
lation in a tin  
miner

FIG 119 —Reticulation  
and massive shadows  
in a granite quarry  
man aged 28 years



FIG 120 —Nodulation,  
massive shadows and  
opacity of mediastinal  
lymph glands  
resembling a bunch  
of grapes in a tin  
miner




FIG 121—Nodulation massive shadows, opacity of mediastinal lymph glands and basal emphysema in a South African gold miner (a patient of Dr L W Hale)

encouraged to substitute safe processes for dangerous ones, as for instance in the use of wet instead of dry methods of quarrying, blasting, drilling and other processes. Dangerous substances used in industry should be replaced by safer substances. Thus aluminium oxide should be used to make grinding wheels instead of sandstone. Mechanical methods should be used to suppress dust. This may be done by the use of wet processes, and exhaust ventilation should be installed so that the dust is removed from the air which the man breathes. Suitable provisions should be made so that the workers can have their meals away from the machinery and the dusty atmosphere. There should always be adequate medical supervision. If the engineer could prevent the creation of dust there would be no disease.

## GRAPHITE

No systematic survey of graphite workers has been carried out but Dunner (1945) described the clinical and radiological findings in men who had been exposed to pure graphite dust unloading ships for periods of about 20 years. Three of the men had cough, sputum and shortness of breath, but two had no complaints. X ray examinations showed reticulation and nodulation. Later Dunner and Bag nall (1946) reported a case of "necrosis cavitation" in the lung of a graphite worker. The sputum contained graphite although the patient had not worked for 5 years in contact with this substance. During recent years the graphite used in Great

Britain has come mainly from Ceylon and Madagascar but there has been a small quantity mined in Borrowdale in Cumberland. Dassanayake (1948) showed that the miners in the workings in Ceylon developed x-ray reticulation. Gloyne, Marshall and Hoyle (1949) recorded two fatalities and Harding and Oliver (1949) three more among men who had been exposed during the manufacture of crucibles to a mixed dust of graphite, silica, sillimanite and other substances. The patients had worked for 30 years and more in contact with the dust and had complained of increasing shortness of breath on exertion, with cough and viscid sputum. Chest movement was poor, there was diminished air-entry, and many crepitations throughout the lungs. The radiographs showed emphysema at the bases, nodulation, and massive shadows of both lungs. The patients all had massive progressive fibrosis, and dust reticulation at necropsy, and the lungs were blackened from the presence of graphite dust.

### PNEUMOCONIOSIS OF COALMINERS

Pneumoconiosis of coalminers (anthracosis) is the most important dust disease in the British Isles. The existence of the disease in South Wales has been known for a century (Meiklejohn, 1951). Early in the nineteenth century Laennec (1806) and Boyle (1815) described a peculiar form of animal production which they called melanosis. Thackrah in 1831 recognized that colliers were liable to develop "miner's asthma" and Cox (1857) described the disease as the "scourge of the mines", but it was not until Hart and Aslett (1942) carried out a survey of the industry, as a result of the rapidly increasing number of compensation certificates that were being issued, that a clear description of it was recorded in the literature. The dust inhaled by the coalminers is a mixture of coal, silicates, and silica and the resulting disease is therefore a form of silicosis. Besides the mixture of dusts the quantity of dust inhaled is a factor in the occurrence of disease. In the past the anthracite mines have been badly ventilated and therefore the quantity of dust inhaled has been high and the incidence of the disease has accordingly been highest in these mines. However, with the installation of better ventilation and the use of mechanical methods in bituminous and steam coal mines the quantity of dust in the atmosphere in all classes of mines has approximated more closely together, and there is probably not such a great difference in incidence as used to be thought.

#### Pathology

If the lungs are examined, black areas will be found scattered throughout the lungs, varying from microscopic size to about 5 millimetres, some of which are soft, others are hard, depending on the amount of fibrous tissue present (Fig. 122). These dust foci are formed around the bronchioles and their accompanying arterioles. A delicate fibrosis develops in the foci of dust which may stop at this stage or proceed to the development of hard collagen tissue. These foci have a crenated edge with processes extending from the edge and fading into unaffected tissue. The background is a pale, yellowish-grey, and the foci are black.





FIG. 122.—Macroscopic appearance of a section of lung of a miner with pneumoconiosis showing coal foci with areas of surrounding emphysema and massive apical area of fibrosis with cavity in the centre (Section by Professor Jethro Gough)

1947) It is the destruction of healthy lung and its replacement by fibrosis and focal emphysema which cause the shortness of breath and consequent disability in this disease

As a background to this simple pneumoconiosis there may be superimposed fibrosis due to infection which may be tuberculous or non tuberculous. Massive areas of fibrosis occur frequently in the upper halves of the lungs, especially in the posterior segment of the upper lobe and the apical segment of the lower. They are round or elongated and are composed of dense black tissue of a hard rubbery consistency. These areas usually have a small cavity in the centre, which may contain some black liquid. There is usually no definite evidence of tuberculosis either to the naked eye or to microscopic examination. Nevertheless it is thought by many workers, probably rightly, that these areas do represent superadded tuberculous infection which may have become dormant or died out. This view is supported by the finding of tubercle bacilli in about one third of the cases. Sometimes these areas extend into the bronchus, in which case the patient may complain of coughing up a quantity of inky black sputum. This has been called melanoptysis.

Right sided heart failure is the most common cause of death in these patients but they may die from tuberculo-toxaemia or thrombosis of the pulmonary artery. The degree of disability will depend on the amount of lung in which function has been lost or impaired. This may be brought about either by fibrosis or

emphysema, although the two generally occur together. The disease presents most rapidly in the workers who are employed in developing the roads to the coal face and are called "hard headers", but it can occur in all workers in the coal mines, or indeed, in coal trimmers working in the docks, or even in furnace men in factories, and coal-delivery men, if the exposure is long enough. The disease is similar to that already described in the section dealing with graphite.

### Clinical course

In the early stages the disease tends to pursue a slow course of dyspnoea on exertion, and this may be the only symptom. As the disease progresses, however, they may develop a cough, and advanced cases may have melanoptysis, and cough up a peculiar type of black sputum. This is associated with the breaking down of large areas of diseased lung tissue. In the late stages, paroxysms of coughing associated with tenacious mucoid sputum sometimes occur. On examination in the early stages there may be no abnormal physical signs, but as the disease develops the patients become obviously dyspnoeic on exertion and eventually orthopnoeic. There may be clubbing of the fingers. The chest tends to become barrel-shaped, and respiratory movements are poor. Flattening of the chest below the clavicles is sometimes marked, there is often hyper-resonance over the bases. Air entry becomes markedly diminished, and when bronchitis is superimposed there will be coarse crepitations throughout the lungs.

The progress of the disease is often accompanied by a rising blood sedimentation rate, but this is usually associated with infection. It is, however, the superimposed infection, which may be either a catarrhal bronchitis or pulmonary tuberculosis, which causes the disability in the condition. Tuberculosis is a common complication and Mann (1951) suggests that it is possible that coal dust exerts a modifying influence on its development and progression. Stewart (1948) found that positive sputum was present in 3 per cent of 400 patients with the disease, but Gloyne (1951) found pulmonary tuberculosis in 100 (34 per cent) out of 293 coal miners' lungs examined at necropsy. He found that the incidence of primary pulmonary neoplasm was 6.9 per cent. The vital capacity and maximum ventilatory capacity are usually below normal and the residual air is increased. Exercise tolerance tests are below standard. Measurements of arterial oxygen will show lowered saturation and tension in advanced cases. In the early stages the skiagrams show

near the hila. In more advanced cases nodulation is present, multiple, discrete shadows 2-3 millimetres in diameter are scattered throughout the lung fields (Figs 124-125). This nodulation in due course may coalesce or conglomerate so that massive shadows develop. These are dense homogeneous shadows usually with a smooth outline, and therefore well-defined, often with a symmetrical distribution, and sometimes referred to as "angel's wings". These shadows are usually associated with well marked basal emphysema. They can occur either with reticulation or nodulation. Following melanoptysis, a central area of translucency develops within a massive shadow, indicating the presence of a cavity. There is sometimes a fluid level (Fig 126).

If the man is removed from the mines in the early stages of the disease it is not progressive. In the later stages it frequently pursues a slow course, but when massive shadows are a prominent feature it may progress rapidly.

### **Treatment**

When the diagnosis is made the patient should be removed from further exposure to coal dust. If this is done in the early stages he will be able to do full time work in most industries. As the disease advances he will be suitable for work in light industry or in sheltered workshops. Treatment for the chronic bronchitis and emphysema which accompanies the disease follows the general line for treatment of these diseases. Inhalation of aluminium dust may cause the patient to express the faith that he is better, but there is no scientific evidence that it produces any change in the course of the disease.

## **PNEUMOCONIOSIS**

It is only since 1930 that radiography of the lungs has been satisfactory. It is not surprising, therefore, that much new knowledge about the effect of dust on the lungs has accumulated during the past 20 years. The characteristic radiographic change attributed to pneumoconiosis is pin-point shadowing scattered throughout the lung. In the description of this nearly every worker has found it

either fluffy and confluent, or else conglomerate, when a massive shadow is seen. The shadowing is of the x-ray type, a soft shadow, and is not a true shadow, as it is not seen on an x-ray film.

There may be no visible difference between the nodulation caused by deposits of iron in the electric arc welder's chest, and that appearing in the classical silicotic or even in the person suffering from chronic miliary tuberculosis, yet the pathological lesions are fundamentally different. The observer gives his own opinion of the cause of the shadowing.

Even though many of the dusts which cause this shadowing are inert, it is rational at least to suppose that the lung cannot become a physiological dust trap and yet retain its elasticity. It seems possible that any dust, no matter how small, in small concentrations would in large enough quantity eventually

### **Inert dusts**

#### *Baratosis*

When metallic dust is inhaled into the lung it may go into solution and be carried around the body and excreted by the kidneys and intestines, or it may remain in the lung tissues as an inert substance. Metals and their salts throw x-ray shadows, the densities of which vary with their atomic weight. Arrigoni (1933) and Pender-




FIG 123—Early reticulation in a South Wales collier




FIG 124 — More definite reticulation and early massive shadow formation in a South Wales steam collier




FIG 125 — Nodulation and massive shadow formation in South Wales anthracite hard header aged 48 years



FIG 126 — Skogram showing a cavity which appeared in a massive shadow in a coalminer's lung after a melanoptysis

grass (1938) have shown that the inhalation of the dust of barytes and barium (atomic weight 137.4) produces the densest of nodular shadows in the lung. Barium itself, however, is inert and does not produce symptoms or disability. If, however, the dust is a mixture containing quartz, silicosis may result.

#### *Siderosis*

Zenker (1866) first showed that iron (atomic weight 55.84) was deposited in the lymphatics of the lung tissue. Doig and McLaughlin (1936) describe the radiographic change which they observed in a group of 6 electric-arc welders (Fig. 127). These men suffered no symptoms and they had no physical signs as a result of inhaling the iron dust, yet their skiagrams showed dense reticulation. Enzer and Sander (1938) showed at necropsy that the iron oxide is deposited in the perivascular and peribronchial lymphatics and does not produce fibrosis in the lung. Later Doig and McLaughlin (1948) showed that the x-ray shadowing will become less marked in the course of years if the men are taken away from the occupation of welding. The study of dust diseases is inevitably complicated by the fact that the dust inhaled is of a mixed nature. Thus workers in foundries are exposed to a mixed dust of iron, silicates and silica (McLaughlin and others, 1950). In these cases it is manifestly impossible to express an opinion as to how much of the x-ray shadowing is due to the inert iron dust, and how much to fibrosis caused by the silica and silicates (Fig. 128). Likewise though the radiographic changes in the lungs of haematite iron miners are highly characteristic, these workers are exposed to a mixture of iron and quartz, resulting in sidero-silicosis with all the symptoms and disability caused by the more severe disease (Fig. 129).

#### *Argyrosiderosis*

Silver polishers polish with rouge, and are thus exposed to a dust consisting of a mixture of silver (atomic weight 107.88) and iron. They develop a cough with copious rouge-coloured sputum. McLaughlin and others (1945) showed radiographically that the lungs of these workers presented a typical picture of reticulation. No evidence of fibrotic change was shown though the lungs may be emphysematous. There were collections of inert iron in the lymphatics. Histologically there is intra-vitam staining of the elastic tissue in the arterial and alveolar walls from the inhalation of particles of silver, which is partly the cause of the x-ray shadowing (Fig. 130).

#### *Stannosis*

Pendergrass and Pryde (1948) reported the case of a workman, aged 45, who had been bagging tin (atomic weight 117.8) oxide dust obtained by treating tin cans with a caustic. The inhalation of the dust produced radiographic nodulation in the lung fields similar to that seen in arc welders. There was little fibrosis. The tin oxide dust contained 96.5 per cent of tin oxide but no silica. When an artery of a freshly excised dog's lung was injected with a saline suspension of tin oxide or the suspension was instilled into the bronchi, even small amounts of it produced appreciable densities in the lung.

#### *Vanadosis*

Wyers (1946) has shown that men employed in bagging vanadium pentoxide in

# OCCUPATIONAL LUNG DISEASES




FIG 128—Siltosis in  
steel moulder  
Nodulation



FIG 127—Siderosis in  
an electric arc welder  
aged 60 years after  
21 years of the work




FIG. 129—Appearances of siderosis in a haematite iron miner



the process of manufacturing vanadium (atomic weight 50.6) may develop radiographic reticulation in their lungs


#### *Aluminium*

Aluminium powder has been used extensively in making explosives incendiary mixtures and paints. Denny Robson and Irwin (1939) produced evidence from animal experiments that aluminium dust would prevent silicosis. Since then miners in various parts of the world have had aluminium dust blown into their changing

from Germany have shown that workers exposed to high concentrations of metallic aluminium dust complained of dry cough with pain on breathing shortness of breath and poor appetite. Their skiagrams show a rapidly progressive disease with extensive fibrosis (Fig. 131). These patients frequently developed spontaneous pneumothorax and several died. The development of the illness was reported within 3 months of starting work. Histological examination of the lungs of patients who died revealed coarse branching hyaline collagenous fibres which enclosed phagocytes containing fine and coarsely granular particles distinguished by jagged outline from carbon particles.

In 1947 Shaver and Riddell in Canada reported that a number of persons





(a)



(b)





FIG. 6. (a) (b) (c)

sect on before removal of the siver ( $\times 1600$   
reduced one quarter (approx) on repro-  
duction)

(c)

engaged in the manufacture of aluminium oxide for abrasive wheels developed a similar disease. In the manufacturing process bauxite is ground and mixed intimately with iron and coke: the mixture is shovelled into large iron pots into which carbon electrodes are lowered. The mixture is fused at about  $2000^{\circ}\text{C}$  as a result of which dense white fumes containing aluminium and silica are evolved. These fumes contain considerable quantities of alumina and silica together with small quantities of many other substances all in a fine state of division. The men first develop attacks of extreme shortness of breath and later complain of sub-sternal pain, cough with white frothy sputum, weakness, loss of appetite and marked cyanosis. Among 344 men 23 developed symptoms and 8 developed bilateral spontaneous pneumothorax. There were 7 deaths in these cases the men had had less than 5 years exposure. The chest signs were variable and depended on the presence or absence of pneumothorax. Diminished chest expansion and impaired percussion note were the principal ones observed. Rales and rhonchi were variable. Tachycardia and cyanosis were marked features. In well established cases the radiographs showed extensive fibrosis with areas of gross emphysema. The lung fields were lace like or granular in appearance. Emphysematous blebs and bulli occurred in the visceral pleura and were apt to rupture spontaneously giving rise to pneumothorax. Microscopical examination of the lungs revealed profound emphysema accompanied by overwhelming fibrous tissue formation. The ash of the lung contained 3 per cent silica and 45 per cent alumina.

FIG 131—Skriagram of man aged 38 years an alumininum stamper for 6 months showing extensive fibrosis of lung and spontaneous pneumothorax



22 2 41

## BIBLIOGRAPHY

### Platinum asthma

- Hunter D, Milton R and Perry K. M. A (1945) Asthma Caused by the Complex Salts of Platinum *Brit J Industr Med* 2 92  
 Karasek S. R., and Karasek M (1911) Report Illinois State Commission on Occupational Diseases p 97

### Organic vegetable materials

- Campbell J. M. (1932) Acute Symptoms Following Work with Hay " *Brit med J* 2, 1143  
 Fawcett R. (1935) Fungoid Conditions of the Lung *Brit J Radol.*, 9 172 354  
 — (1938) Occupational Diseases of the Lungs in Agricultural Workers *Brit J Radol* 11 378  
 — (1938) Roentgenological Recognition of Certain Bronchomycoses Involving Occupational Risks," *Amer J Roentgenol* 39 19  
 Fuller C. J. (1947) Personal Communication  
 Hoffman W. (1946) Die Dreschkrankheit *Schweiz med Wschr* 76, 988  
 Torneil, E. (1946) Threshers Lung A fungoid disease resembling tuberculosis in morbus Schauman *Acta med Scand* 125 191  
 Zettergren L. (1950) Threshers Lung (pulmonary moniliasis) An experimental investigation *Uppsala Läk Fren Förh* 55, 257

# OCCUPATIONAL LUNG DISEASES

## Bagasse

- Castleden L I M and Hamilton Paterson I L (1942) Bagassosis an Industrial Lung Disease *Brit med J* 2, 478
- Hunter D and Perry K M A (1946) Bronchiolitis Resulting from the Handling of Bagasse *Brit J Indust Med* 3 64
- Schneider R Reinhart W H and Caminita B H (1948) Aerobacter cloacae Endotoxin as a Possible Factor in the Etiology of Bagassosis *J Indust Hyg* 30 238
- Sodeman W A and Pullen R L (1943) Bagasse Disease of the Lung *Med surg J* 95 558
- (1944) Bagasse Disease of the Lungs *Arch intern Med* 73, 365
- Cotton (b) sinusosis**
- Bolen H L (1943) Byssinosis—Report of Two Cases and Review of Literature *J indust Hyg Toxicol* 25 215
- Collis E L (1915) The Occurrence of an Unusual Cough Among Weavers of Cotton Cloth *Proc R Soc Med* 8 108
- Dunn S and Sheehan H L (1932) Report on 10 Autopsies on Cotton mill Workers Appendix III *Report of the Departmental Committee on Dust in the Card rooms in the Cotton Industry* London H M Stationery Office
- Gill C I C (1947) Byssinosis in Cotton Trade *Brit J Indust Med* 4 48
- Greenhow E H (1862) Medical officer of the Privy Council Fourth Report H M Stationery Office p 173
- Jackson J (1818) On the Influence of the Cotton Manufactories on the Health *Lond med phys J* 39 464
- Middleton E L (1926) Weavers Cough *J indust hyg* 8, 428
- Neal P A (1936) Report on Acute Illness Among Rural Mattress Makers Using Low Grade Stained Cotton *J Amer med Ass* 119 1074
- Prausnitz C (1936) Report of the Departmental Committee on Dust in Card rooms in the Cotton Industry (1932) *Spec Rep Ser M R C No 212* London H M Stationery Office
- (1936) Study of sickness among operatives in Lancashire Cotton Spinning Mills (1930) *London Indust Hlth Research Bd Rep No 590*
- Zipperlin V R (1934) Über das örtliche Vorkommen von Asthma bronchiale in Württemberg und Hohenzollern *Arch Hyg* 113 1

## Carcinoma of the lung

- Amor A J (1938) Report on 8th International Congress in Industrial Accidents and Occupational Diseases *Leipzig 1939* 2, 248
- Arnstein A (1913) Über den sogenannten Schneeberger Lungenkrebs *Verh dtsch path Ges* 16 332
- Boetjers A M (1950) Pulmonary Carcinoma in Chromate Workers I A review of the literature and report of cases 2 Incidence on basis of hospital records *Arch Ind str Hyg occup Med* 2, 457
- Harting F H and Hesse W (1879) Der Lungenkrebs die Bergkrankheit in den Schneeberger Gruben *Vjschr ger chit Med* 30 296 31 102
- Hill A B and others (1948) Studies of the Incidence of Cancer in Factory Handling Inorganic Compounds of Arsenic Mortality Experience in Factory *Clinical and Environmental Investigations Brit J Indust Med* 5, 1
- Hueck W (1939) Kurzer Bericht über Ergebnisse anatomischer Untersuchungen in Schneeberg *Z Krebsst* 49 312
- Lange K (1935) Krebskrankungen und geologische Verhältnisse im Erzgebirge *Z Krebsforsch* 42 306
- Leitch A and Kennaway E L (1922) Experimental Production of Cancer by Arsenic *Brit med J* 2 1107
- Lowy J (1929) Über die Joachimstaler Bergkrankheit vorläufige Mitteilung *Med klin Berl* 25 141
- Lorenz E (1944) Radioactivity and Lung cancer critical review of lung cancer in miners of Schneeberg and Joachimsthal *J Nat Cancer Instit* 5 1

## BIBLIOGRAPHY

- Machle W and Gregonous F (1948) "Cancer of Respiratory System in United States Chromate Producing Industry" *Publ Hlth Rep Wash* 63 1114
- Feiler S (1939) Lung Cancer Among Mine-workers in Joachimsthal *Human Biol* 11 130
- Purchan A and Siki H (1932) Cancer of Lung in Miners of Jáchymov (Joachimsthal) report of cases observed in 1929-1930 *Amer J Cancer* 16 681
- Rostloski O, Saupé E and Schmorl G (1926) Die Bergkrankheit der Erzbergleute in Schneeberg in Sachsen (Schneeberger Lungenkrebs) *Z Krebsforsch* 23 360
- — — (1926) Die Bergkrankheit der Erzbergleute in Schneeberg in Sachsen (Schneeberger Lungenkrebs) *Ibid* 25 249
- Siki H (1930) Über den Lungenkrebs der Bergleute in Joachimsthal (Tschechoslowakei) *Z Krebsforsch* 32, 609
- Thompson V C (1946) Personal communication
- Wedler H W (1943) Asbestose und Lungenkrebs *Dtsch med Wschr* 69 575
- Wyers, H (1950) Asbestosis *Postgrad med J* 25 631
- Zel R (1935) Zur Frage des Lungenkrebses bei den Bergleuten Joachimsthals *Med Klin* 31, 1535
- China clay**
- Hale L W (1946) Personal communication
- Middleton E L (1936) Industrial Pulmonary Disease due to Inhalation of Dust with Special Reference to Silicosis *Lancet* 2 59
- Thomas R W (1952) Silicosis in the Ball Clay and China Clay Industries *Lancet* 1 133
- Talc**
- Cesa Bianchi D and Devoto L (1911) Patologia polmonare da inalazione di polveri *Rivista di Patologia* 481 555
- Dreessen W C (1933) Effects of Certain Silicate Dusts on the Lungs *J Industr Hyg* 15 66
- and Dalla Valle J M (1935). Effects of Exposure to Dust in 2 Georgia Talc Mills and Mines *US Publ Hlth Rep* 50 131
- McLaughlin A I G, Rogers E and Dunham K C (1949) Talc Pneumoconiosis *Brit J Ind Med* 6 184
- Merewether E R A (1933) *Annual Report of the Chief Inspector of Factories* p 63 London H M Stationery Office
- (1934) *Annual Report of the Chief Inspector of Factories* p 65 London H M Stationery Office
- Millman N (1947) Pneumoconiosis Due to Talc in the Cosmetic Industry *Occup Med* 4 391
- Schäfer W (1952) Talc Hazard in Tremolite Talc Mining *Amer J Roentgenol* 49 11
- T — — — zur pathologischen Anatomie der
- Z — — — *Il Lavoro* 1, 1
- Asbestos**
- Auribault (1906) *Bull Insp Tra Paris* p 126
- Beger P J (1933) Über die Asbestoskörperchen *Virchows Arch* 290, 280
- Cooke W E (1974) Fibrosis of the Lungs due to the Inhalation of Asbestos Dust *Brit med J* 2 147
- (1927) Pulmonary Asbestosis *Brit med J* 2, 1204
- and Hill C F (1927) Pneumoconiosis due to Asbestos Dust *J R Microsc Soc* 47 222
- Fahr T (1914) Asbestosis pneumoconiosis *Munch Med Woch* 61 625
- Gardner L U and Cummings D E (1931) Studies in Experimental Pneumoconiosis Inhalation of Asbestos Dust: Its Effect upon Primary Tuberculous Infection *J Pathol Hyg* 13 112.

# OCCUPATIONAL LUNG DISEASES

- Gardner L U (1933) *Pathological Reaction in Various Pneumoconioses* *J Amer med Ass* 101 594
- (1942) *Chrysotile Asbestos as Indicator of Subtle Differences in Animal Tissues* *Amer Rev Tuberc* 45 762
- Gloyne S R (1930) *Reaction of Tissues to Asbestos Fibre with Reference to Pulmonary Asbestosis* *Tubercle* 11, 151
- (1931) *Presence of Asbestosis Bodies in Faeces in Case of Pulmonary Asbestosis* *Tubercle* 12, 158
- (1932) *Asbestos Body* *Lancet* 1, 1351
- King E J Clegg J W and Rae V M (1946) *Effect of Asbestos and of Asbestos and Aluminum on Lungs of Rabbits* *Thorax* 1 188
- Kruger E Rostowski O and Saupe E (1931) *Über Lungenasbestose* *Arch Gewerbepath*
- Linzbach A J and Wedler H W (1941) *Beitrag zum Berufskrebs der Asbestarbeiter* *Arch Gewerbepath*
- McDonald S (1927) *Histology of Pulmonary Asbestosis* *Brit med J* 2 1025
- Merewether E R A and Price C W (1930) *Report on the Effects of Asbestos Dust on the Lungs and Dust Suppression in the Asbestos Industry* London H M Stationery Office
- (1933) *Memorandum on Asbestosis* *Tubercle* 15 69 109 152
- (1934) *Annual Report of the Chief Inspector of Factories* p 62 London H M Stationery Office
- Middleton E L (1936) *Industrial Pulmonary Disease due to Dust with Special Reference to Silicosis (Milroy Lecture)* *Lancet* 2 59
- Murray M (1907) *Dept Comm on Compensation for Industr Diseases* Cd 3495 p 14 Cd
- Page R C (1935) *Study of Sputum in Pulmonary Asbestosis* *Amer J med Sc* 189 44
- Policard A (1930) *Methode de detection et d'etude des particules minerales dans les tissus* *Bull Histol Tech met* 7 129
- Seiler H E (1928) *Case of Pneumoconiosis result of Inhalation of Asbestos Dust* *Brit med J* 2 982
- Seiler H E and Gilmour M D (1931) *Case of Pulmonary Asbestosis* *Brit med J* 1, 1112
- Simson F W and Strachan A S (1931) *Asbestosis Bodies in Sputum study of Specimens from 50 Workers in Asbestos Mill* *J Path Bact* 34 1
- Stewart M J and Haddow A C (1929) *Demonstration of the Peculiar Bodies of Pulmonary Asbestosis (Asbestos Bodies) in Material Obtained by Lung Puncture and in the Sputum.* *J Path Bact* 32 172
- Wood W B and Gloyne S R (1934) *Pulmonary Asbestosis review of 100 cases* *Lancet* 2 1383
- Wyers H (1950) *Asbestosis* *Post Grad med J* 25, 631
- Silliman  
Jotten K W and Eickhoff F W (1944) *Lung Changes from Sillimanite Dust* *Arch Gewerbepath Gewerbepath* 12, 223
- Middleton E L (1936) *Industrial Pulmonary Disease due to Inhalation of Dust with Special Reference to Silicosis* *Lancet* 2, 59
- Mica  
Ferguson T Quoted by Middleton E L (1936) *Lancet* 2, 59
- Diatomaceous earth and tripoli  
Legge R T and Rosencrantz, E (1932) *Observations and Studies on Silicosis by Diatomaceous Silica* *Amer J Publ Hlth* 22 1055
- Luton P Champere J and Faure P (1949) *Pneumoconiosis from Diatomaceous Earth in French Kieselguhr Workings* *Arch Mal prof* 10 217
- Martin Vallaud and Champere (1947) *The X ray Appearances of the Lungs of Diatom Workers* *Le Med cin d Usine* 9 295
- Middleton E L (1940) *Proceedings of the International Congress of Geneva 1938* p 134

# BIBLIOGRAPHY

Vigliani E C and Mottura G (1948) Diatomaceous Earth Silicosis *Brit J Indust Med* 5, 148

## Silicosis

- Denny J J Robson W D and Irwin D A (1939) Prevention of Silicosis by Metallic Aluminium" *Canad med ass J* 40 213
- Gardner L U (1932) Studies on Experimental Pneumoconiosis Inhalation of Quartz Dust *J Indust Hyg* 14 18
- Gloyne S R (1951) Pneumoconiosis A Histological Survey of Necropsy Material in 1205 Cases *Lancet* 1 810
- Heffernan P (1948) What is Silicosis? *Tubercle* 29 170
- Kettle E H (1932) "Interstitial Reactions Caused by Various Dusts and their Influence on Tuberculous Infections" *J Path Bact* 35 395
- Macdonald G Piggott A P and Gilder F W (1930) Two Cases of Acute Silicosis with a Suggested Theory of Causation *Lancet* 2, 846
- Meiklejohn A (1949) Silicosis in Potteries and Sandstone Workers *Brit J Indust Med* 6, 230

## Graphite

- Dassanayake W L P (1948) Health of Plumbago Workers in Ceylon *Brit J Indust Med* 5 141
- Dunner L (1945) Observations on Pulmonary Disease in Graphite Workers *Brit J Radiol* 18 33
- Dunner L and Bagnall D J T (1946) Graphite Pneumoconiosis Complicated by Cavitation of the Lungs *Br J Radiol* 19 166
- Pneumoconiosis Produced by Graphite  
in Lungs Produced by Natural Graphite

## Pneumoconiosis of coalminers

- Bayle G L (1815) *Researches on Pulmonary Phthisis* Trans by W Barrow Liverpool
- Cox W I (1857) Diseases of Special Occupations No III Diseases of Collers in South Lancashire *Brit med J* 5, 491
- Fletcher C M (1948) Pneumoconiosis of Coalminers *Brit med J* 1 1015
- Gloyne S R (1951) Pneumoconiosis A Histological Survey of Necropsy Material in 1205 Cases *Lancet* 1 810
- Gough J (1947) Pneumoconiosis in Coal Workers in Wales *Occup Med* 4 86
- Hart P D and Aslett E A (1942) Chronic Pulmonary Disease in South Wales Coalminers *Spec Rep Sen Med Res Coun London* No 243
- Lazner M (1954) *Le Bulletin de*

Man

Tuberculosis and

- Meiklejohn A (1951) History of Lung Diseases of Coal Miners in Great Britain 1800-75" *Brit med J* 8 127
- Stewart A (1948) Pneumoconiosis of Coal Miners Study of Disease after Exposure to Dust has Ceased" *Brit J Indust Med*, 5 120
- Thackeray C T (1831) *The Effect of Arts Trades and Professions on Health and Longevity* p 27 London

## Baratosis

- Arrigoni A (1933) "La pneumoconiosi da bario (Osservazioni clinico-radiologiche e sperimentali)" *Med d Lavoro* 24 461
- Pendergrass, E P (1938) *Lansilicosis and Asbestosis* p 137 London Oxford University Press
- Preti L and Talmi P C (1938) Report on Eighth International Congress on Industrial Accidents and Occupational Diseases. *Le pz g*, 2, 963

# OCCUPATIONAL LUNG DISEASES

## Siderosis

- Britten J A and Walsh E L. (1940) Health Hazards of Electric and Gas Welding. Review of Literature. Results of Examination of 286 Welders. Preventive Measures. *J Indust Hyg Toxicol* 22, 125
- Buckell M Garrad J Jupe M H McLaughlin A I G and Perry K M A (1946) Incidence of Siderosis in Iron Turners and Grinders. *Brit J Indust Med* 3, 78
- Doig A T and McLaughlin A I G (1936) X ray Appearances of Lungs of Electric Arc Welders. *Lancet* 1, 771
- Enzer N and Sander O A (1938) Clearing of X ray Shadows in Welders' Siderosis. *Lancet* 1, 789
- Simonson E and Evans A M (1945) Chronic Lung Changes in Electric Arc Welders. *J Indust Hyg Toxicol* 20, 333
- Gordon D (1950) Clinical and Physiological Observations on Welders with Pulmonary Siderosis and Foundry Men with Nodular Uncomplicated Silicosis. *J Industr Hyg Toxicol* 27, 147
- Groh J A (1944) Pneumokomosis in Queensland Foundries. *Med J Austr* 2, 217
- Hamlin L E and Weber H J (1950) Siderosis. A Benign Pneumoconiosis due to the Inhalation of Iron Dust. A Clinical Radiological and Industrial Hygiene Study of Foundry Cleaning Room Employees. *Industr Med* 19, 151
- Harding H E (1945) Radiographic and Histological Appearance of Rat Lung after Intratracheal Injection of Rouge. *Brit J Indust Med* 2, 32
- Harrold G C Meek S F and McCord C. P. (1940) Chemical and Physiological Investigation of Electric Arc Welding. Bare Washed Welding Rods. *J Industr Hyg Toxicol* 22, 347
- McLaughlin A I G Cheseaman E A Garrod J Gloyne S R Goodall K L Harding H E Jupe M H Laurie W B Perry K M A Sutherland C. L Woods H (1950) *Industrial Lung Disease of Iron and Steel Workers* London H M Stationery Office
- Pendergrass E P and Leopold S S (1945) Benign Pneumoconiosis. *J Amer med Ass* 127, 701
- Sander O A (1944) Further Observations on Lung Changes in Electric Arc Welders. *J Indust Hyg* 26, 79
- Zanetti E and Dompe M (1950) Silicosis and Tuberculosis among Steel Foundry Workers. *Med d Lavoro* 41, 321
- Zenker F A (1866) Ueber Staubinhalationskrankheiten. *Dtsch Arch klin Med* 2, 116

## Stannosis

- Pendergrass E P and Pryde A W (1948) Benign Pneumoconiosis due to Tin Oxide. A Case Report with Experimental Investigation of the Radiographic Density of Tin Oxide Dust. *J Indust Hyg Toxicol* 30, 119

## Vanadiosis

- Wyers H (1946) Some Toxic Effects of Vanadium Pentoxide. *Brit J Indust Med* 3, 177

## Aluminium

- Beit, T F and King E J (1943) Failure of Aluminium to Prevent Silicosis. *J Path Bact* 55, 69
- Crombie D W Blaisdell J L and MacPherson G (1944) Treatment of Silicosis by Aluminium Powder. *Canad med Ass J* 50, 318
- Denny J J Robson W D and Irwin D A (1939) Prevention of Silicosis by Metallic Aluminium. *Canad med Ass J* 40, 213
- — — (1939) Prevention of Silicosis by Metallic Aluminium. *Industr Med* 8, 133

# BIBLIOGRAPHY

Gorale  
St

- (1943) Weitere Erfahrungen zum Krankheitsbild der Aluminiumlung. *Deutsches Tuberk.—Bl.* 17: 3
- (1947) The Aluminum Lung, a New Industrial Disease. *Z. ges. Med.* 2, 665
- and Jäger R. (1941) Zur Klinik Pathologie und Pathogenese der Aluminiumlung. *Arch. Gene. hep. u. Gewerbeh.* 11: 102
- Hunter D. Milton R. Perry K. M. A. and Thompson D. R. (1944) Effect of Aluminum and Alumina on Lung Grinders of Duraluminum Aeroplane Propellers. *Brit. J. Indust. Med.* 1: 159
- Nguyen R. and Nguyen F. (1941) Colloid and Histochemistry. A case of Lung Infection.



## CHAPTER 24

### CYSTS OF THE LUNG

GEOFFREY FLAVELL

CYSTIC DISEASE of the lung may be congenital in origin, or it may be acquired. The distinction is often hard to make since pathological changes in the lung soon after birth may be mistaken for, or indistinguishable from, truly congenital lesions; and defects first seen in later life may have their root in developmental faults. Degenerative changes, certain specific pneumonias, the tension cavities of pulmonary tuberculosis, bronchial obstruction by neoplasm, secondary epithelialization of inflammatory spaces and parasitic infestation can on occasion all produce similar appearances. Some are truly cysts, some abscesses, and some mere tears in the parenchyma modified by the elastic tensions of the web in which they occur. The differential diagnosis between these lesions, often identical both in their clinical manifestations and their x-ray appearances, presents a task of exceptional difficulty which may be soluble only by histological means. On the other hand the history, clinical course, and allied features of many pseudo-cysts will satisfactorily identify them.

Tuberculous cavitation, for which cysts have commonly been mistaken, should be revealed for what it is by infiltration elsewhere and the finding of bacilli; pyogenic acute abscesses are generally associated with some obvious cause, and when they appear to lack this, or their clinical behaviour is atypical, they demand close investigation, those secondary to neoplasm may either fall into this latter group, or are clearly the product of an obstructing or necrotic growth. But multiple cysts may so closely resemble cystic bronchiectasis that opinion is still divided as to whether bronchiectasis is not sometimes congenital, though most cases seen in infancy are now believed to follow early atelectasis. When cysts are full of fluid there is practically no way of distinguishing them from any of the solid pulmonary tumours, or indeed sometimes from localized effusions or empyemas. Finally a division has to be made between the cysts of congenital, and mainly bronchial, origin and those which are acquired, and arise from the alveoli.

The symptoms accompanying cysts are largely the symptoms of their complications but successful treatment depends wholly upon the diagnosis of a cyst as prime cause, and failure to make this diagnosis means that treatment, too, must fail.

### CONGENITAL CYSTS

When the developing lung buds grow out from the primitive fore-gut of the embryo they may be disordered at any point along their course. If this occurs at an early stage a fragment may be pinched off near the dividing trachea and a mediastinal bronchial cyst be left, while if a more substantial part of the anlage goes astray a whole sequestered pulmonary lobe results. Further peripherally, cysts will be found in the lung, solitary if one bronchial bud is involved, multiple if many are

involved, if a bronchial communication is present the cysts will also contain air and be subjected to the changes in its pressure, but, if isolated, they may be full only of fluid or of mucus which may gradually distend them or reach a state of quiescent equilibrium.

In all cases these cysts are characterized by a lining of respiratory epithelium, cuboid or columnar, ciliated or not, as well as by a wall in which the elements of a bronchial wall, fibrous and elastic tissue, mucus glands and fragments of cartilage, are present, usually in disorderly array. It is the possession of these features which principally distinguishes them from other cysts of the lung. Kindred anomalies such as conjoined and supernumerary ribs, but particularly aberrant pulmonary vessels and unusual segmental fissures, are frequent and are strong evidence of congenital disorder.

Large cysts are detected in early infancy because they make their presence felt. They are generally air-containing, and when the first breaths enter the newly expanded lung it balloons in such a fashion that the small and pliant bronchus to which they are connected is compressed against the bulging cyst wall and its entry made more and more oblique, until at last air is admitted only during the bronchial expansion of inspiration and can no longer escape with contraction and expiry. A vicious circle is thus established and the cyst grows with every breath and cry. At first only the related lung is compressed, but soon the cyst fills the hemithorax causing respiratory distress and cyanosis, each growing more severe with every gasp until the mediastinum is displaced, and carries with it the trachea and the heart. At this stage all the signs and symptoms of a tension pneumothorax will be present, x-rays show a deviated mediastinum, the functioning lung of one side compressed by it, while upon the other is no sign of pulmonary markings, the cyst itself conforming completely to the chest wall. This predicament in infancy is practically always due to a ballooning cyst, and the insertion of a tension pneumothorax needle high in the chest, connected by tubing to a water seal, is lifesaving. Puncture of the cyst may cause pneumothorax, and then the delicate arc of the cyst wall is sometimes visible on x-ray examination, making the diagnosis plain, since air entry into the cyst is valvular, however, such pneumothoraces afford no relief but are in turn subjected to an increasing positive pressure.

Solitary fluid-containing cysts, or air cysts in which a check valve mechanism has not developed will, if large enough, cause pressure symptoms by their bulk alone, but should this not be the case they lie fallow in the lung probably about the size of oranges, until overtaken by infection (Fig 132). Multiple cysts, similar in their behaviour and in histology are in general smaller, and either cluster in one lobe like a group of two or three soap bubbles, or involve an entire lobe or even lung in diffuse honeycombing amongst which almost no parenchymal tissue survives. While this may resemble acquired "cystic" bronchiectasis, peribronchial inflammatory damage is wanting, the cysts are bigger than usually occurs in bronchiectatic dilatation, and other stigmata of congenital disturbance are often present (Fig 133). Such silent cysts will continue undetected into childhood, adolescence or early adult life, although attention may be drawn to them by an unexplained, persistent cough, dyspnoea or haemoptysis. The latter is more often found when cystic disease is diffuse, and is occasionally severe. Asthma or early



FIG 132 (a) and (b)—A typical congenital bronchial cyst communicating with a bronchus. The patient had no symptoms and the cyst has not yet become infected so that the fluid in its lower third is probably mucus

(a)



(b)




FIG 133—A case of cystic honeycombing involving the whole of the right upper lobe and associated with dextrocardia. This combination is sometimes known as Kartagena's syndrome. The congenital malposition of the heart is convincing evidence of the congenital origin of the cystic disease in the upper lobe.

clubbing in children especially if they appear somewhat undersized, and if the chest seems curiously deep or asymmetrical should indicate the possibility that congenital cysts are present. The right lung is more commonly affected than the left.

Sooner or later since the great majority communicate with the bronchial tree lung cysts are likely to become infected. If they are single chronic ill health with cough recurrent fevers and purulent sputum may follow and should x ray examination reveal a fluid level in the chest lung abscess or empyema is perhaps diagnosed. It has often been the case that large parietal fluid-containing cysts even when not infected have been mistaken for empyemas and drained by rib resection indeed ill advised transpleural aspiration of an infected cyst is calculated to result in true empyema. Drainage however does not result in the progressive diminution of the cavity and a long period of illness with a persistently draining sinus will continue until it is realized that a cyst is present and resection carried out. Infected cysts may also rupture into the pleura or suddenly establish a fistulous communication with the bronchial tree flooding it with pus. Where

cerebral abscess is ever present.

Islands of pulmonary tissue having no connexion with the adult bronchial tree or at most a tenuous one are sometimes found in the chest. They receive no blood

supply from the pulmonary circulation, but are nourished by an aberrant vessel rising directly from the aorta, or the coeliac plexus (in which case it pierces the diaphragm from below), and representing one of the foetal vascular plexuses. Analogous in origin both to bronchial and enterogenous cysts, since they are the result of an abnormal and isolated outgrowth from the primitive fore-gut, such pulmonary islands are commonly situated posteriorly in the left chest (Fig 134). They are sometimes at the apex, where they may have a narrow connexion with the side of the trachea, or wholly sequestered at the base; and they cast upon x-rays the opaque shadow of airless lung. It is not surprising, in view of their origin, that they are on occasion partially or wholly cystic, and that such cysts may become secondarily infected. Bronchography shows a normal bronchial tree having no connexions with the sequestered area. At thoracotomy they are adherent to surrounding tissues, and the artery supplying them may not only be unexpected in position, but also of formidable size.

### Treatment

The correct treatment of congenital cysts, whether they be the ballooning air


... the case of honeycombed lobes or lung, lobectomy  
tances the preservation of  
the surgeon's aim

### ACQUIRED CYSTS

The cysts of this category, with the exception ... (229), are pseudo cysts, inasmuch as they lac-  
cysts, and walls peculiar to themselves, if  
bronchiectasis and other pulmonary cavities of specific provenance are likewise  
set on one side, those remaining share also the characteristic that they arise from  
the alveolar part of the lung and not from the bronchi, and that they always  
contain air

### Blebs and bullae

... develop under  
ure the normal  
in the alveolar  
mesh, smooth rounded cavities appear surrounded by a narrow atelectatic zone,  
and grow in size disproportionately to the pressure exerted upon them  
(Moolten 1935). Not only is the lung itself an elastic net, but it is subjected  
... as a hole  
gap in the  
y coalesce  
m If the  
ht degree,



(a)



FIG 134 (a and b)—A sequestered lobe with marked cystic changes situated on the posterolateral aspect of the right lower lobe in a child of 6 years. Numerous fistulae existed between it and the base of the right lower lobe through which a great deal of pus drained. It obtained its blood supply through a large arterial trunk piercing the right diaphragm from below. At operation it was possible to cut it bodily from the base of the lower lobe without bleeding. The fistulae were closed and the child made an untroubled recovery.

(b)

air may be able to enter the alveoli supplied by it during the bronchial dilatation of inspiration, but not to escape again when contraction occurs with expiration. Thus a small alveolar space may rapidly balloon (the effect being enhanced by the forced inspirations that precede coughing) and the surrounding parenchyma compressed before the advancing bubble. Ultimately it may reach an equilibrium, or it may continue to expand until it bursts through the visceral pleura, producing a pneumothorax, or grow still vaster as ballooning congenital cysts do, producing mediastinal displacement and symptoms of compression.

A similar mechanism is well seen in the tension cavities that accompany tuberculosis, and it is perhaps significant to observe that remarkable thin-walled cystic cavities often appear during the course of acute staphylococcal pneumonias. These cavities are at first quite empty, but later have a level of staphylococcal pus in them (by which alone they are to be identified as abscesses) and, now aided by antibiotics, shrink and vanish as quickly as they came (Fig. 135). It seems probable that some at least of the ballooning cysts seen in children, when alveolar and not bronchial in type, may be "cysts" of this sort, secondary to an unrecognized staphylococcal pneumonitis, rather than cysts of congenital origin, more especially since the sudden onset of staphylococcal empyema in infancy was once common and could be explained by the same process.

The pneumatocoeles known as blebs and bullae appear first as a localized breakdown of alveolar walls weakened by infection or by the degenerative changes of chronic emphysema. The vesicle is then extended by the inherent stresses of the lung, and finally blown up by a valvular ingress of air. Blebs are defined as subpleural collections of air caused by rupture of alveoli immediately beneath the visceral pleura, while bullae result from the coalescence of one alveolus with

and stresses

earlier age-groups is obscure but may depend upon inflammatory scarring of terminal bronchioles by pneumonitis or, it has been suggested, by healed subclinical pulmonary tuberculosis. It is also conceivable that they might arise because of a congenital defect in the bronchial wall, but this seems less likely.

In infants they may attain great size, collapsing the lung from which they spring and displacing the mediastinum in the same manner as ballooning congenital cysts,

gressive respiratory distress with cyanosis accompany them, and rupture from them produces either no effect or results in pneumothorax. Occasionally they vanish as mysteriously as they appear, resembling in this respect the thin-walled "abscesses" of staphylococcal pneumonia. Radiologically there is complete absence of lung markings on the affected side, but fine trabeculae may be

# ACQUIRED CYSTS

FIG. 135 (a and b)—Acute staphylococcal abscesses in the lungs simulating cysts

(a)



(b)





FIG 136 (a and b)—A ballooning lung cyst in an adult patient. There is no evidence of emphysema in the rest of the lung fields.


(a)



(b)

# ACQUIRED CYSTS

FIG 137—(a) A giant cyst related to the left upper lobe in a man aged 66 years and simulating a spontaneous pneumothorax. At operation it was excised and fistulous communications with the compressed upper lobe were sutured. (b) The lung then re-expanded without further difficulty to fill the chest.



## CYSTS OF THE LUNG

distinguishable especially at the base of the cyst or the curvature of its edge seen in the costo phrenic angle

During the second and third decades of life similar blebs and bullae occur but do not often reach the large proportions of infancy. The lungs show no other sign of disease and are not generally emphysematous, the cysts being confined to one area. While they may cause dyspnoea by progressive enlargement their chief importance at this age lies in their being the source of idiopathic or asymptomatic spontaneous pneumothorax. A high proportion of such pneumothoraces are chronic the lung not re expanding within three months and frequently recurrent. They thus result in severe incapacitation often unnecessarily prolonged by inadequate or ill conceived treatment and carry as well a serious risk to life since an infection or similar accident upon the opposite side is likely to cause death. In spection of the lung with a thoracoscope may reveal the collapsed and punctured bleb but sometimes this is not apparent and must be searched for at thoracotomy like a leak in a bicycle tyre.

In the past such pneumothoraces have been treated by inducing chemical pleurisy with insufflated iodized talc or silver nitrate among many other irritants and withdrawing air in the hope that the lung will adhere to the chest wall. The process is painful not always successful and tends permanently to diminish respiratory efficiency. There is little doubt that in most cases and certainly in all those where the lung fails to re expand within a few weeks direct attack upon the cyst provides the best hope of early and lasting cure. Even though the patient may have a seriously reduced respiratory capacity either because of a large cyst or because of a pneumothorax as soon as the chest is opened and the bleb punctured or the leak secured it is possible to re expand the lung and restore normal respiration. These cysts can be separated from the neighbouring tissue by sharp dissection. The interior is often multilocular being divided by trabeculations which are characteristic when seen in x ray films and at the base are usually a number of small bronchial communications requiring closure. Such operations promptly carried out save the patient months of illness and distress as well as avoiding the mortal dangers of contra lateral disease.

The bullae seen in older men and associated with emphysema fortunately do not have this tendency to burst perhaps because their connexions with the

1. None

become marked local excision of the bulla attains little or no result but other bullae will undoubtedly be present in the lung probably bilaterally and each case must be judged upon its merits and necessity

## BIBLIOGRAPHY

- Adams W E (1949) Differential Diagnosis and Treatment of Congenital Cysts, Malformation of the Lung *Dis Chest* 15 60  
 Allbritten F F and Templeton J Y (1950) Treatment of Giant Cysts of the Lung *J thorac Surg* 20 749  
 Allison P R (1947) Giant Bullous Cysts of the Lung *Thorax* 2, 169  
 Anspach W E and Wolman I J (1933) Large Pulmonary Air Cysts of Infancy *Surg Gynec Obstet* 56 635

## BIBLIOGRAPHY

- Brewer L A, Dolley F S and Evans B H (1940) The Surgical Management of Chronic Spontaneous Pneumothorax *J thorac Surg* 19 167
- Hock R C (1948) Recurrent and Chronic Spontaneous Pneumothorax *Thorax* 3 88
- Cassels D, Fritz J and Adams W E (1949) Congenital Cystic Disease of the Lung *J Pediatr* 35 585
- Clagett O T (1949) Surgical Treatment of Emphysematous Blebs and Bullae "Dissecting" *Ann Surg* 123 229
- Dugan D J and Samson P C (1950) The Surgical Treatment of Giant Emphysematous Blebs and Pulmonary Tension Cysts *J thorac Surg* 20 729
- Gross R E (1946) Congenital Cystic Lung "Large Infected Solitary Pulmonary Cysts Simulating Empyema" *J thorac Surg* 9 471
- Maier H and Haight C (1940) "Large Infected Solitary Pulmonary Cysts Simulating Empyema" *J thorac Surg* 9 471
- Moersch H J and Clagett O T (1947) Pulmonary Cysts *J thorac Surg* 16, 179
- Moolten S (1935) Mechanical Production of Cavities in Isolated Lungs *Arch Pathol* 19 825
- Naciri E and Langer L (1947) Pulmonary Cysts *Surveys* 22, 516
- Nicholson H (1950) Suppurative Pneumonia *Lancet* 2, 549
- Perry K M A (1939) On Spontaneous Pneumothorax *Quart J Med* 8 1
- Ravitch M and Hardy J B (1949) Congenital Cystic Disease of the Lung in Infants and Children *Arch Surg* 49 1
- Sellers T S (1938) Congenital Cystic Disease of the Lung *Tubercle* 20 49
- Strode J E (1949) Cystic Disease of the Lung *J Thorac Surg* 18 404
- Susman M P (1949) The Treatment of Pulmonary Hydatid Disease *Thorax* 3 71
- Tyson M D (1943) The Treatment of Pulmonary Hydatid Disease *J thorac Surg* 19 422
- Pulmonary Obliteration *Ann Surg* 118 50

## CHAPTER 25

### TUMOURS OF THE LUNG

VERNON C THOMPSON

*THE ORIGIN* of a pulmonary tumour is almost always from the cellular structures of a bronchus. Far the most common tumours are those arising from the bronchial epithelium, and in this category carcinoma comprises about 98 per cent.

Although benign tumours of the lung are comparative rarities their clinical recognition is of the utmost importance. If they are diagnosed and treated the prognosis for the patient is excellent; if undiagnosed they may cause death by their complications or they may produce years of invalidism.

The following classification is offered as a basis for further study. It cannot claim to be complete as it omits such rarities as myoma, endothelioma and other pathological curiosities that have been reported from time to time, but it includes the lesions of main clinical importance, in a natural order of precedence.

<i>Primary malignant</i>	Carcinoma
	Sarcoma
	(a) round celled sarcoma
	(b) fibro sarcoma
	(c) lympho sarcoma
<i>Secondary malignant</i>	Carcinoma, sarcoma, etcetera
<i>Benign</i>	Adenoma
	Hamartoma
	(a) chondroma
	(b) fibroma
	(c) lipoma
	(d) vascular hamartoma
	Neurogenic tumours

### PRIMARY CARCINOMA OF THE LUNG

#### *Incidence and aetiology*

... the cause of the incidence of primary ... on in males as ... from this age ... mum incidence ... cline. Mason (1949) analysing 1 000 cases finds the incidence of age and percentage as follows: 30 years, 2.2 per cent; 30-40, 9.6; 40-50, 30.9; 50-60, 37.6; 60-70, 18.6; over 70, 1.1. These figures may be taken to represent common experience in any large series of cases. There is no doubt that there has been a steady increase in the incidence of cancer of the lung in both sexes in the past 50 years. Hutchison (1928) drew attention to the increasing incidence of this condition in the first quarter of the

## PRIMARY CARCINOMA OF THE LUNG

century. In the second quarter the increased incidence has far exceeded that of the first. In 1922 the number of deaths from this cause in England and Wales was 612, a fairly insignificant number. In 1949 cancer of the lung accounted for 11,272 deaths. In males the lung has now become the commonest site of primary malignant disease.

The cause of this increase has been a matter of considerable speculation and investigation. Speculation suggests that the increase may be accounted for by improved methods of diagnosis, and the survival of more people into the cancer age. These factors must undoubtedly account for at least some part of the supposed increase in incidence, but they can hardly account for the whole. Investigation has mainly been devoted to the search for some inhaled carcinogen which has increased *pari passu* with the increasing incidence of carcinoma of the lung.

The possibility of an inhaled carcinogen has been recognized for many years, and the increased incidence of carcinoma of the lung among workers in certain occupations has been studied throughout the years. The most noted example of this disease is an industrial hazard among workers in the Schneeberg mines in Czechoslovakia. In more recent years attention has been drawn to the added risks of cancer of the lung in workers with certain mineral ores. Perry (1947) has drawn attention to the increased incidence among workers in arsenic and in certain nickel and copper ores in which arsenic occurs as an impurity. Machle and Gregorius (1948) have reported that the death rate for cancer of the lung was 25 times the normal among workers in the chromate-producing industry in the United States of America.

It is notable that there is little or no evidence that chronic irritation by inflammation such as occurs in chronic bronchitis, pulmonary tuberculosis or bronchiectasis has any specific bearing on the development of carcinoma, though most clinicians of wide experience are able to recall isolated examples of the development of neoplasms in chronically inflamed lungs. No definite association has been found between carcinoma and the inhalation of dust in general, as typified by coal dust producing anthracosis in miners.

Wynder and Graham (1950) in the United States of America and Doll and Hill (1950) in Great Britain report statistical investigations into the effect of tobacco smoking on the incidence of carcinoma of the lung. They point out the correlation between the increase in the habit of cigarette smoking and the increase of lung cancer. They each report a careful review of the smoking habits of a large series of patients with carcinoma of the lung as compared with a similar number of controls. Their conclusions that heavy cigarette smokers run a far greater risk of lung cancer than non-smokers are irrefutable. Doll and Hill go so far as to say that the risk of developing the disease increases in simple proportion with the amount smoked, and after the age of 45 the risk in heavy smokers may be as much as 50 times as great as among non-smokers.

The evidence so far suggests that though chronic inflammation and irritation by non specific dusts are not entirely innocuous, the development of carcinoma of the lung is much more probable when specific carcinogens are inhaled. These carcinogens may occur in dusts or fumes associated with certain occupations and may occur in tobacco smoke. The precise nature of the carcinogens has not been

## TUMOURS OF THE LUNG

determined with any certainty. Much work has been done on this subject, but much more remains to be done in further elucidation of the problem.

### Pathology

#### Histology

There was considerable confusion about the histology and nomenclature of carcinoma of the lung until *Barnard (1926)* following *Turnbull and Bartlett* showed that malignant epithelial tumours of the lung were often pleomorphic, but all could be derived from the basement layer of the bronchial mucous membrane. Before this date many of the primary lung tumours had been classified as "oat cell sarcoma" or "lympho-sarcoma" of the mediastinum.

Perhaps the commonest histological type of lung carcinoma is the squamous celled growth which evolves with metaplasia, for squamous epithelium is not a normal component of the bronchus. For purposes of description carcinoma of the bronchus is divided into three types.

(1) The oat-cell carcinoma, so called in Great Britain (*Fig 138*). This term is employed because the cells resemble oat seeds, they have large darkly staining nuclei and little cytoplasm. They are in general undifferentiated tumours with little fibrous stroma. In other countries they are usually referred to as small-celled or anaplastic growths.

(2) The squamous-cell carcinoma (*Fig 139*), composed of flattened squamous cells with a varying degree of fibrous stroma, often showing keratinization. In the more highly differentiated types cell nests are present.

(3) The columnar-cell carcinoma (*Fig 140*) in which the growth is composed of well-established columnar cells which have an alveolar formation with well-developed stroma. There may be intra-alveolar papilliferous projections. About one half of the columnar-cell tumours secrete mucus.

Although these three main types are described, it is not always possible in any particular case to place a tumour in any one of these categories, for different types of cells may be found in the same tumour, and transitional cells such as spheroidal cells and polygonal cells are encountered. The classification of a group of tumours will therefore always depend to some extent on the personal opinion of the pathologist.

The incidence of oat cell and squamous cell tumours is about equal. The least common is the columnar-cell tumour which comprises only about 10 per cent of the total.

Although there must be innumerable exceptions to any generalization there are

When it occurs in a major bronchus it metastasizes early to the lymph nodes which become tense with growth and the pericardium is involved early (*Fig 141*). Squamous cell carcinoma is more common in elderly people, it varies greatly in its malignancy. It may grow very slowly and metastasize late. It occurs, like all lung tumours, with greatest frequency in the major bronchi, but when it develops peripherally it shows a tendency to break down in its centre and cavitate (*Fig 142*).

FIG. 138 —Oat-cell carcinoma  
( $\times 470$  Reduced one quarter  
on reproduction)





FIG. 139 —Squamous-cell carcinoma ( $\times 330$  Reduced one  
quarter on reproduction)

FIG. 140 —Columnar-cell carcinoma ( $\times 210$  Reduced one  
quarter on reproduction)








FIG 141 —Oat-cell carcinoma of lung showing med astinal extension involving and surrounding great vessels Autopsy specimen

FIG 142 —Squamous-cell carcinoma of upper lobe showing necrotic growth with cavitation surrounding lung abscess Operation specimen



In these cases it may extend centrifugally and involve the chest wall before metastasizing. The adenocarcinoma is remarkable chiefly for its incidence. Although all other types are far commoner in men than women, adenocarcinoma is almost equally divided between sexes. It shows one other characteristic which is somewhat exceptional. This is its tendency to spread by seedlings. Once it involves the pleura the latter rapidly becomes coated with small seedlings which coalesce and produce plaques of growth. It is this tendency to pleural spread which has often masked the primary tumour and caused the erroneous diagnosis of endothelioma of the pleura. Adenocarcinoma also tends to seed itself bronchogenically so that a small primary tumour may produce miliary bronchogenic seedlings throughout the lung.

#### *Macroscopic appearance*

About 75 per cent of all lung tumours arise in the major bronchi. 25 per cent arise in the smaller bronchi and are situated peripherally in the lung. Seldom do they occur in the trachea. Those occurring in the major bronchi may present as proliferating endobronchial tumours (Fig 143) or they may produce a funnel-shaped stricture by sub mucous infiltration (Fig 144). Eventually they cause obstruction of the bronchus with collapse of the alveoli distal to the growth. This

## PRIMARY CARCINOMA OF THE LUNG

Fig 143—Adenocarcinoma of lung showing intra bronchial proliferating growth from parenchymatous mass in right upper lobe. Operation specimen



Fig 144 Squamous cell carcinoma showing peribronchial infiltration and stricture of the bronchus. Operation specimen

collapse may be segmental lobar or total according to the site of the primary tumour. Secondary changes in the lung occur distal to the growth. Even in the absence of infection mucus distension and bronchiectasis appear early. If there is infection suppurative pneumonitis, bronchiectasis and lung abscess may complicate the condition. These complications may extend to the pleura so that pleural effusion or empyema may ensue. Growths situated peripherally are more likely to present in the early stages as solid rounded tumours without the complications of bronchial obstruction but as they extend they involve structures outside the lung.

### Mode of spread

A malignant tumour of the lung may spread in many ways

*Direct extension*

In tumours near the hilum this extension tends to be towards the mediastinum so that the great vessels and the pericardium and the phrenic and vagus nerves become involved. In peripheral growths the extension may reach the chest wall without other complications causing erosion of ribs and spreading through the thoracic cage into the soft tissues beyond. At the apex of the lung the brachial plexus, the sympathetic chain, the nerve roots and the vertebral column may become involved. At the base of the lung the primary growth may infiltrate the diaphragm.

*Metastasis*

Metastasis may be by the lymphatics, by the blood stream or by the bronchi.

*Lymphatic spread*

Lymphatic spread occurs as a rule first to the neighbouring lymph nodes at the hilum of the lung, thence the growth spreads into the mediastinal lymph nodes where it may cross the mid-line and extend upwards into the neck and downwards to the retroperitoneal lymphatics below the diaphragm. Another form of lymphatic spread may occur in the lung itself by a process of linear permeation, producing the condition known as carcinomatous lymphangiosis (Fig. 145).

*Blood-borne metastases*

Dissemination into the blood stream occurs as a result of the escape of emboli into the pulmonary veins. As might be expected these emboli are often carried into the cerebral vessels and cause metastases in the brain, sometimes these metastases are solitary. Once an embolus escapes into a pulmonary vein it passes into the systemic circulation and may lodge and grow anywhere in the body—usually haematogenous metastases are multiple. In addition to the brain they show a particular predilection for the liver, the bones, the suprarenals and the skin, but no tissue is immune (Fig. 146).

*Bronchogenic metastasis*

Rarely, some carcinomas tend to grow like a fern into and within the lumina of the bronchi, this is a method of direct spread from this primary growth seedlings may become separated and become dispersed into other parts of the bronchial tree where they proceed to grow separately. Usually these bronchogenic metastases occur in the same lung, but they may arise in the other lung. This particular form of spread is most often encountered with the columnar-cell growths.

*Diagnosis*


It is often a simple matter to diagnose advanced inoperable malignant disease of the lung. This diagnosis may be of some interest to the clinician, but the accuracy of the diagnosis is of little importance to the patient for by then his fate is sealed and he might just as well be suffering from any other rapidly progressive lethal condition. To diagnose tumours of the lung at an early stage is of supreme importance to the patient, and this can only be achieved by first suspecting the presence of a tumour and then carrying out a series of investigations to prove it.

# PRIMARY CARCINOMA OF THE LUNG

FIG. 145—Carcinoma  
with lymphangiosis



FIG. 146—Carcinoma of  
the left lower lobe with  
metastases throughout  
both lungs.



# TUMOURS OF THE LUNG





Fig 147—(a) A hilar mass in a young woman due to oat cell carcinoma (b) film taken on expiration shows medial displacement away from the lesion due to obstructive emphysema by partial obstruction (c) film two weeks later shows collapse of the right lung due to total obstruction of the bronchus



# PRIMARY CARCINOMA OF THE LUNG




FIG. 148.—Squamous-cell carcinoma. A rounded hilar shadow observed in a patient with signs of a cerebral tumour. A cerebral metastasis was removed and subsequently left pneumonectomy was performed. The patient has remained well and held a responsible post 3½ years after operation.





FIG 147—(a) A hilar mass in a young woman, due to oat-cell carcinoma, (b) film taken on expiration shows mediastinal displacement away from the lesion due to obstructive emphysema by partial obstruction, (c) film two weeks later shows collapse of the right lung due to total obstruction of the bronchus



PRIMARY CARCINOMA OF THE LUNG




FIG 148.—Squamous cell carcinoma. A rounded hilar shadow observed in a patient with signs of a cerebral tumour. A cerebral metastasis was removed and subsequently left pneumonectomy was performed. The patient has remained well and held a responsible post 3½ years after operation.



### *Symptomatology*

It is unfortunately well substantiated by mass radiography, and chance findings on screening the chest at radiological examination for other lesions such as peptic ulcer, that carcinoma of the lung can be quite silent and symptomless until its intra-thoracic extensions have rendered it inoperable. It is also not uncommon for the first symptom to be caused by metastases. Before resection of the lung became a safe and successful procedure, the outlook for patients with pulmonary neoplasms was indeed hopeless and there was little to be gained by making either an early or an accurate diagnosis. In spite of the prospects of successful surgery in certain cases the prognosis is still grave, partly because of the silent onset of the disease and partly because of the tendency to metastasize. The outlook however is not so grave as to justify the pessimistic and fatalistic attitude that is still prevalent and which perhaps accounts for the long delay in diagnosis that occurs even when symptoms are present. Bjork (1947) analysing 112 inoperable cases found an average delay of 8.4 months from the first symptom to reference to hospital. Mason (1949) found the average duration of symptoms in his 1,000 cases was over 6 months. In the writer's experience at the London Hospital the average duration of symptoms is 7 months.

Cough is the commonest first symptom—this is such a prevalent malady, particularly in the winter, that it cannot be fully investigated at once, but when it occurs in a patient over 40 and persists without adequate reason for more than 2 or 3 weeks a fuller investigation is indicated. The other more common early symptoms are chest pain, dyspnoea, haemoptysis and lassitude. Haemoptysis is a fairly dramatic event and should never be treated expectantly without proceeding to a diagnosis. Unfortunately it is not the commonest first symptom. Brooks and

C. — 11.72 per cent of 402

The symptomatology of carcinoma of the lung can only be fully understood by reference to its pathology. When the growth is peripheral it is silent until it breaks down or ulcerates into a major bronchus, then it may cause cough, haemoptysis or purulent sputum from secondary infection. If it extends to the pleura it causes chest pain, or it may announce itself by metastases.

When the primary growth arises in a major bronchus it causes cough, possibly haemoptysis. As it encroaches on the lumen of the bronchus it may cause a persistent wheeze and dyspnoea by obstruction of the air-way. Later there may be obstructive emphysema with increase of breathlessness. This is followed when the obstruction of the bronchus is complete by collapse of the lung distal to the growth and complications due to infection ensue (Fig 147). The symptoms may then become those of pneumonia, bronchiectasis, lung abscess, pleurisy or empyem. Loss of weight should never be regarded as a pure symptom of a primary growth of the lung when it occurs in association it is usually due to toxæmia caused either by metastasis or infection.

The mode of onset of symptoms in carcinoma of the lung may be so diverse that it is possible to describe certain quite different clinical types of onset. In a large



FIG 149—Peripheral squamous-cell carcinoma. A small round shadow in a patient aged 60 years who presented with no chest symptoms but with marked pulmonary osteoarthropathy. Treated by left lower lobectomy with immediate relief of joint pains and no evidence of recurrence 3 years later.

hospital with special departments patients may be referred on their first attendance to many different divisions of the hospital. The Department of Chest Disease inevitably invites the majority of cases because of their frank pulmonary symptoms. The Department of Cardiology sees a number of patients who are referred for dyspnoea, and sometimes patients are referred with paroxysmal cardiac irregularities which are due to involvement of the heart itself by the growth. Cases of mediastinal obstruction and superior vena cava obstruction also find their way to this department. The Departments of Neurology and Neurosurgery see a number of patients with primary carcinoma of the lung. Taylor and Waterhouse (1950) found that the first symptom was neurological in 6.1 per cent of their cases. These patients fall into three categories, they are mostly those with signs of a cerebral tumour and a metastatic one (Fig. 148). The second category is

patients complaining of hoarseness due to paralysis of the left vocal cord by involvement of the recurrent laryngeal nerve. The Orthopaedic Department sees patients with bone and joint pains or spontaneous fracture due to metastases. The Department of Physical Medicine may receive patients complaining of generalized joint pains described as rheumatism, which are actually due to pulmonary osteoarthropathy (Fig. 149). The Department of General Medicine

which always treats a number of cases of frank chest disease will also find patients with carcinoma of the lung referred for pyrexia, toxæmia or cachexia without dominant chest symptoms. This department will also find a small number of cases among those referred for abdominal symptoms.

### *Physical examination*

There are no physical signs which are absolutely conclusive in diagnosing carcinoma of the lung except those of inoperability. In early cases no abnormality whatever may be detected on ordinary clinical examination.

The physical signs in the chest itself are those of the varied lung pathology which may be caused by the tumour and its complications. Impaired air entry and a persistent rhonchus may be caused by partial bronchial obstruction, later there may be signs of pulmonary collapse with shift of the trachea or apex beat towards the side of the lesion and impaired breath sounds and impaired resonance to percussion over the collapsed portion of the lung (Fig 150). Obstructive emphysema may cause hyper-resonance and almost absent air entry when the bronchus is obstructed only on expiration.

to percussion and displacement of the mediastinal structures away from the growth

The physical signs of the lung lesion may be marked by the signs of pleural effusion. When pleural effusion is associated with underlying pulmonary atelectasis the displacement of the mediastinum that might be expected as a result of the fluid is counteracted by the opposite pull of the collapsed lung.

The general examination of the patient must be thorough. It will not help in diagnosing the early case, but it is essential to correct treatment and may reveal metastases that contra-indicate operation. No part of the body is quite immune from metastases. Special attention should be paid to the nervous system, the supraclavicular lymph nodes, the skeletal system, the liver and the skin.

Fever and cachexia are signs of secondary infection or metastases. The time to diagnose carcinoma of the lung is when the patient is still in good health, not when "he looks like a case of cancer".

### *Special investigations*

... the history, the ... the com- ... ke or

exclude a positive diagnosis of growth

### *Radiographic examination*

Tumours arising in a major bronchus cannot be revealed by a plain x-ray until they cast a shadow either by their own bulk or they produce complications such as pneumonia or atelectasis. If obstructive emphysema is suspected clinically, it should be clearly indicated to the radiologist. This condition may not be shown in the normal film taken on inspiration, but it becomes quite obvious if films are taken on expiration as well (Fig 151).

# PRIMARY CARCINOMA OF THE LUNG

FIG 150—(a) An indefinite shadow below the left hilum due to an anaplastic carcinoma of the lower lobe (b) film taken three weeks later showing collapse of the left lower lobe with a small pleural effusion

(a)

(b)

With the proviso that early growths cannot be disclosed by routine chest films, it is true to say that sooner or later the growth can be discovered by radiology. In actual practice the great majority of growths of the lung are detectable by x ray examination by the time the patient seeks medical advice. The radiological picture is either one of pulmonary obstruction and its sequelae of collapse and infection (Figs 152 and 153), or the opacity seen in the radiograph is caused by the mass itself, in which case the tumour is usually peripheral. The peripheral growths may appear as solid rounded or lobulated shadows, often with an ill defined edge with radiating striae extending from it, or they may show a cavity with a fluid level due to necrosis in the substance of the shadow (Fig 154). Mason (1949) found that in 64 per cent of cases the picture was one of collapse, and in about 10 per cent a cavity mimicking a chronic lung abscess was seen.

### Bronchoscopy

A simple and harmless procedure, tumours involving the major bronchi are seen directly and a biopsy can be taken. In other cases though a biopsy may not be possible the diagnosis may be inferred from narrowing, rigidity and distortion of the bronchial tree, when these findings are associated with the radiological appearances. Apart from diagnosis bronchoscopy is an essential investigation in determining the operability and the correct treatment of pulmonary tumours.

Only a limited part of the bronchial tree is visible on bronchoscopy, so negative findings do not necessarily exclude a diagnosis of carcinoma.

### Sputum examination

Sputum examination is not an important matter in establishing the diagnosis when the bronchoscopic findings and particularly the biopsy are positive, but when bronchoscopy is negative, examination of the sputum for malignant cells is of considerable value.

which cannot be excluded from any well organized chest unit, but its value depends entirely on the individual skill and experience of the pathologist who examines the sputum. When bronchoscopy gives no positive evidence sputum examination may reveal malignant cells in those cases of peripheral growth in which there is central necrosis and ulceration, but sputum examination may still be negative if the growth is peripheral and symptomless.

### Bronchography

an segmental filling defect by bronchography, but this is usually confirmed by radiography which demonstrates the pulmonary segmental collapse distal to the obstruction.

# PRIMARY CARCINOMA OF THE LUNG


FIG 151—Obstructive emphysema due to carcinoma of the left main bronchus (a) on inspiration shows no gross abnormality though there is mediastinal displacement to the left and some increase in the left hilar shadow (b) taken in expiration shows failure of the air to escape from the left lung with mediastinal displacement to the right. On physical examination there was no air entry to the left lung. The patient was breathless.



(a)



(b)



*FIG. 152—Collapse of right upper lobe due to carcinoma of bronchus treated by pneumonectomy (a) before operation (b) 2 years after pneumonectomy*



(b)

## PRIMARY CARCINOMA OF THE LUNG

If serial skiagrams show that a shadow in the lung field is growing steadily in size it is almost certainly a growth

A wedge shaped opacity corresponding anatomically with a pulmonary segment is due to bronchial obstruction and in patients over the age of 40 the most probable cause of the obstruction is neoplasm

The peripheral carcinoma that breaks down and presents radiographically as an abscess with fluid level can be differentiated from a chronic lung abscess by the character of its wall. In neoplasms the wall is uneven and presents irregularities and bosses which are never present in a pure pyogenic abscess. The clinical history in these two conditions is also characteristically different. In chronic abscess there is usually a history of copious purulent sputum. In neoplasm the sputum is less in amount and generally it is persistently blood stained.

The solid tuberculous focus may present great difficulty in differential diagnosis. As a rule tuberculous lesions in the lung are not solitary so the difficulty does not often arise. The presence of tubercle bacilli in the sputum is confirmatory evidence of tuberculosis but this finding does not necessarily exclude a diagnosis of carcinoma for the two diseases may co exist. A period of observation by repeated x ray examination is justifiable only if the clinician makes the reservation that the lesion should be regarded as a growth if it becomes larger.

### Treatment

The treatment of choice for carcinoma of the lung is resection. In most cases total pneumonectomy is required. For technical reasons alone it is usually necessary to remove the whole lung in order to remove the growth. It is desirable when operating for malignant disease to perform a so called radical operation by excising the area of lymphatic drainage together with the tumour. This type of operation can only be accomplished by total pneumonectomy. Even so the operation cannot be regarded as a complete radical excision as the lymphatics transgress the midline and are not all accessible to the surgeon.

The operation of lobectomy permits only a limited excision of lymphatics but it has a definite and valuable place in the treatment of lung cancer. The loss of a whole lung may cause serious incapacity to a patient over the age of 60 the loss of a lobe produces little disability. Patients with low cardio respiratory reserve may die of heart failure after total pneumonectomy or they may survive only to be crippled by dyspnoea. The preservation of healthy lung tissue in such patients is often of more importance than the hypothetical value of a radical operation.

The malignancy of primary neoplasms of the lung varies enormously and the prognosis after operation seems to depend more on the degree of malignancy than on the radical nature of the operation.

Conclusive statistical evidence exists as to the operability of malignant



*Tomography*

In delineating the depth, the shape, the edge and the relations of certain indefinite shadows tomography has some value but it cannot give a positive diagnosis. Tomography may be misleading when it is employed to demonstrate strictures of the bronchi—none of the bronchi is quite parallel with the plane of the tomograph and even when normal they may appear to become narrowed as they transect the level at which the film is focused.

*Aspiration biopsy*

Aspiration biopsy and drill biopsy are mentioned only to be condemned as normal methods of investigation. These procedures should be employed only in cases in which it is clear from other investigations that the lesion is inoperable but almost certainly neoplastic. If applied indiscriminately they may lead to complications such as pneumothorax, empyema and dissemination of the growth. They are only of value in determining questions of prognosis and sensitivity to radiotherapy.

*Differential diagnosis*

It is clear from the preceding sections that there must be a full investigation of patients suspected of pulmonary neoplasm and the steps in this investigation have been described. There will remain a small proportion in whom the diagnosis will still remain in doubt. These are the patients in whom the lesion is beyond bronchoscopic vision and radiology presents a persistent abnormality. In this connexion it is important to consider the observations of Raeburn (1951) who regards bronchoscopic evidence as a fairly late sign of most pulmonary neoplasms. Even so there are at present certain limits to our powers of diagnosis and we cannot claim to be able to diagnose carcinoma of the lung in the presence of normal x ray examination, a normal bronchoscopy and a negative sputum examination.

The differential diagnosis therefore limits itself to the interpretation of x ray shadows when bronchoscopy and sputum examination are negative. It is of course possible that diagnosis may be delayed by failure to employ bronchoscopy and sputum examination because the radiological shadows resolve as a result of chemotherapy. The antibiotic drugs are now so successful in resolving the infective complications of new growths that there is a serious danger that they may even further delay the diagnosis of the primary bronchial tumour. Patients with atypical pneumonia of any sort that does not resolve symptomatically and radiologically within one month from the date of onset, should be suspected of

Most of the delays in diagnosis are due to cases in which the radiological picture is equivocal and bronchoscopy and sputum examination are negative. It is in this particular group that the skill, the experience and the judgment of the clinician becomes of supreme importance. The type of case that falls into this difficult category is the patient whose radiograph shows an atypical peripheral shadow. It is beyond the scope of this chapter to discuss all the possibilities of diagnosis, but certain general principles may be laid down.

## PRIMARY CARCINOMA OF THE LUNG

If serial skiagrams show that a shadow in the lung field is growing steadily in size it is almost certainly a growth  
A wedge shaped opacity corresponding anatomically with a pulmonary segment is due to bronchial obstruction and in patients over the age of 40 the most probable cause of the obstruction is neoplasm

The peripheral carcinoma that breaks down and presents radiographically as an abscess with fluid level can be differentiated from a chronic lung abscess by the character of its wall. In neoplasms the wall is uneven and presents irregularities and bosses which are never present in a pure pyogenic abscess. The clinical history in these two conditions is also characteristically different. In chronic abscess there is usually a history of copious purulent sputum in neoplasm the sputum is less in amount and generally it is persistently blood stained

The solid tuberculous focus in the lung are not solitary so the difficulty does not often arise. The presence of tubercle bacilli in the sputum is confirmatory evidence of tuberculosis but this finding does not necessarily exclude a diagnosis of carcinoma for the two diseases may co exist. A period of observation by repeated x ray examination is justifiable only if the clinician makes the reservation that the lesion should be regarded as a growth if it becomes larger

### Treatment

The treatment of choice for carcinoma of the lung is resection. In most cases total pneumonectomy is required. For technical reasons alone it is usually necessary to remove the whole lung in order to remove the growth. It is desirable when operating for malignant disease to perform a so-called radical operation by excising the area of lymphatic drainage together with the tumour. This type of operation can only be accomplished by total pneumonectomy. Even so the operation cannot be regarded as a complete radical excision as the lymphatics transgress the midline and are not all accessible to the surgeon

The operation of lobectomy permits only a limited excision of lymphatics but it has a definite and valuable place in the treatment of lung cancer. The loss of a whole lung may cause serious incapacity to a patient over the age of 60 the reserve may die of heart failure after total pneumonectomy or they may survive only to be crippled by dyspnoea. The preservation of healthy lung tissue in such patients is often of more importance than the hypothetical value of a radical operation

The malignancy of primary neoplasms of the lung varies enormously and the prognosis after operation seems to depend more on the degree of malignancy than on the radical nature of the operation

### Operability

No satisfactory statistical evidence exists as to the operability of malignant growths of the lung and it is logical to assume that there never can be any true assessment. The operability of a tumour depends entirely on the date of diagnosis. Our present methods of diagnosis are insufficiently employed. Most authors are agreed that there is an average delay of 6-8 months from the date of the first

symptom until the patient comes to operation. On the other hand mass radiography has revealed many patients without symptoms in whom the lesion is inoperable, and many more patients present with initial signs of inoperability such as metastases. Although operability rates may be increased by more diligent and accurate methods of diagnosis, it seems inevitable that a large percentage of patients with carcinoma of the lung will be unsuitable for operation.

Brock (1950) reports that of 800 cases seen in the years 1941-48 the operability rate was 13.25 per cent, of 449 cases seen 1945-48 the operability rate was 19 per cent showing an increase in operability rate in the later years. These figures are probably higher than the general figure as the patients seen by an individual surgeon or in a surgical unit are preselected. In general the operability rate is low, it reflects the failure to diagnose the condition at an early stage.

*The diagnosis of inoperability*—During the routine examination and investigation attention should be directed not only to the diagnosis but also to the prospects for successful resection. Advanced age accompanied by its normal degenerative changes such as myocardial weakness and pulmonary emphysema must be assessed in relation to the disease. Age itself is not a contra-indication to operation, but there are few patients over 70 who have sufficient cardio-respiratory reserve to survive with one lung. The main general contra-indication to operation is dyspnoea. Chronic nephritis, diabetes, hypertension and similar general conditions are not contra-indications to operations for cancer.

The extension of the growth itself either by direct spread or by metastases is the usual sign of inoperability. This extension should become evident in the investigation of the patient. Particular attention should be paid to the central nervous system, the skin, the lymph nodes, particularly those in the supraclavicular region, and to the liver. Pain which is not related to the actual site of the tumour may point to metastases in bone or even in skeletal muscles.

Nerve involvement within the chest may affect the recurrent laryngeal nerve producing hoarseness by paralysis of a vocal cord. Tumours at the apex of the lung may involve the brachial plexus and the sympathetic trunk, producing a Horner's syndrome and brachial neuralgia. The phrenic nerve may be involved

trachea, or the growth may be seen to involve these structures directly.

Pleural effusion is not always a sign of inoperability. If the effusion is deeply blood stained it may be taken to denote pleural involvement, but if it is clear or even purulent the effusion may be the result of infection and in this event the lesion is not necessarily inoperable.

Direct involvement of the chest wall as indicated by rib erosion is not a contra-indication to operation as it is often possible to remove a part of the chest wall *en masse* with the lung or lobe (Fig. 155).

In spite of the most careful investigation there will remain a group of patients in whom operability can only be decided by thoracotomy. When there is no absolute contra-indication the patient should be given the chance of survival that only surgery can offer. At operation every effort should be made to remove the

FIG 155—(a) A peripheral squamous cell carcinoma involving the chest wall and eroding the 4th rib. The only symptom was pain. (b) 3 years after left upper lobectomy at which the involved area of chest wall was excised *en masse* with the tumour.

(a)

(b)

growth if this is technically possible. Even if the operative findings point to early recurrence removal of the lung or lobe is the best palliative treatment once the chest has been opened. Removal of the growth should include the mediastinal lymph nodes, but it may also involve resection of a part of the chest wall, the diaphragm or the pericardium and even in exceptional cases a part of the wall of the left auricle. Some cases will defy the most heroic attempts at removal particularly when there is massive mediastinal involvement. In the writer's experience of 232 operations for carcinoma of the lung in the period 1941-51, 27 per cent proved inoperable at thoracotomy.

### *Results of resection*

The first successful pneumonectomy for carcinoma of the lung was carried out by Graham in 1933, this operation was the herald of the great advances that have been made in the surgery of malignant disease of the lung. The technical difficulties of the operation and of anaesthesia have been overcome, so that now it is possible to remove a lung or a lobe with few technical complications. Operative mortality depends not so much on the character of the operation as on the general state of health of the patient. The antibiotics, particularly penicillin, have contributed largely to safety. Before the penicillin era healing of the bronchial stump, which was almost invariably infected, was an uncertain feature of the operation and bronchial fistula was a common occurrence. Infection of the remaining lung was also a serious contribution to morbidity and mortality. In the writer's experience in the years 1941-45 the operative mortality was 28 per cent and the incidence of bronchial fistula was 48 per cent. In the years from 1945-51 the operative mortality was reduced to 7 per cent and the incidence of bronchial

There are occasional examples of patients with growths of low grade malignancy who have survived for several years without resection of the tumour. There are also innumerable examples of patients with tumours of high grade malignancy who have died of disseminated metastases in a few months or even weeks. If patients survive for more than

for later recurrence to occur

first year after operation, the operation for advanced malignant disease. Most of the patients for whom resection is employed as a palliative procedure die within 1 year. The following Table shows the fate of 91 patients (80 pneumonectomies and 11 lobectomies) operated upon by the writer in the years 1941-48.

Since 1948 the operative mortality has been reduced, but it is unlikely that the 3-year survival rate will be improved unless patients are available for operation at a much earlier stage in their disease. Other authors have observed similar results, which are perhaps a little less favourable only because they were published earlier in a period of improving techniques in diagnosis and surgery, Tudor Edwards (1946), Ochsner and DeBakey (1948), Brock (1948), Churchill and his colleagues (1950).

# PRIMARY CARCINOMA OF THE LUNG

TABLE  
RESULTS OF 91 OPERATIONS FOR MALIGNANT DISEASE

Result	Number of cases	Percentage
Post-operative deaths	12	13
Died of recurrence in the 1st year	29	32
Died of recurrence in the 2nd year	11	12
Died of recurrence in the 3rd year	2	2
Deaths from other causes	6	7
Survivors	31	34

When the growth has extended beyond the lung either by glandular metastases or direct spread the prognosis after surgery is vastly altered. An analysis of the writer's cases shows that when there is no extension beyond the lung the 3 year survival rate was 60 per cent but when extension has occurred the survival rate is reduced to 15.7 per cent. A 60 per cent 3 year survival rate after resection for malignant disease offers a fair prognosis to patients and should be a great incentive to early diagnosis.

## Deep x ray therapy

Radiotherapy has something to offer to patients with new growths of the lung but its possibilities are limited and it is employed mainly as a palliative. The

which occur in young people and rapidly produce symptoms of mediastinal obstruction. The prognosis in such cases is almost hopeless but the symptoms

drugs should be given to clear up the infection before the treatment is started.

The best prospect for radiotherapy is probably in those cases of peripheral growth in which surgery is contra indicated only by old age or dyspnoea.

## Palliative treatment

Apart from radiotherapy the palliative treatment of carcinoma of the lung consists of the relief of pain, the relief of cough, the treatment of infection and the relief of dyspnoea.

There is much that can be done to increase the comfort of patients with inoperable cancer of the lung. Morphine and its derivatives are invaluable and should be

neurectomy is impractical as the nerve roots themselves are often involved. In

these cases the operation of cordotomy is often justifiable and earns the deep gratitude of the patient

Severe toxæmia and copious sputum may result from infection of the lung distal to the growth. Remarkable symptomatic improvement can be obtained in such cases by the employment of suitable antibiotic drugs, of which penicillin is at present the one in most common use

Pleural effusion may cause distressing dyspnoea, and should be aspirated when indicated to relieve symptoms. If the effusion is infected it should also be sterilized by the introduction of antibiotic drugs

### ALVEOLAR-CELL CARCINOMA AND ADENOMATOSIS

This section would be incomplete without reference to a type of tumour which is receiving increasing attention in the literature, to which the reader is referred for further information

The condition is described as *alveolar-cell tumour*, *alveolar carcinoma* and *pulmonary adenomatosis*. It bears a close similarity to a condition found in

acceptance of the nature and pathology of the lesion. There is for instance no agreement that the alveoli normally have an epithelial lining. Many pathologists regard the tumours as adenocarcinomas, which are showing bronchogenic dissemination. Clinically the lesion behaves as a low grade malignant growth, causing haemoptysis, dyspnoea and cough. Ultimately unless resection is successful, the patient dies of asphyxia, pneumonia or metastases. As far as treatment is concerned surgical removal of the tumour offers the best hope of survival. Ikeda (1945) has discussed this condition fully, and Griffith, McDonald and Clagett (1950) have reported 7 cases and made a statistical study of 51 other cases reported in the literature

### SARCOMA OF THE LUNG

Primary sarcoma of the lung must be considered rare. It is usually difficult to make the diagnosis during life without operation as these growths are seldom visible on bronchoscopy

Round-cell or spindle-cell growths may occur as highly malignant tumours in young people. They may present with symptoms of pain or toxæmia and radiography will demonstrate a pulmonary mass. In the absence of metastases they are best treated by a combination of surgery and deep x-ray therapy but the prognosis is poor

Fibro-sarcoma of the bronchus is a great rarity that may be detected on bronchoscopy, and may present with the signs and symptoms of bronchial obstruction. These tumours are of low-grade malignancy and the prognosis after removal of the growth is good. The subject is fully discussed by Black (1950) and by Carswell and Kraeft (1950). The writer has seen only 1 case of this condition, a child aged 2½ years who has survived for 7 years without evidence of recurrence after local excision of the tumour

Lympho-sarcoma usually involves the lung secondarily from an origin in the

## ADENOMA OF THE BRONCHUS

mediastinum but there have been a few reported instances of lympho-sarcoma starting in the lung and presenting as an intrapulmonary mass. The diagnosis in these cases is only made after resection which appears to be a successful procedure. It should be combined with deep x ray therapy. Maier (1948) describes a case of this nature.

## SECONDARY MALIGNANT TUMOURS OF THE LUNG

The lung is a common site for metastasis from malignant tumours in other parts of the body. When metastasis occurs in the lung it is usually generalized and the evidence of the pulmonary lesion is masked by the other signs of dissemination of the growth.

The secondary deposits in the lungs are caused by vascular embolic spread and consequently the lesions are usually multiple. The radiological appearance is quite characteristic and consists of multiple rounded shadows throughout both lungs (Fig 156). As time passes the shadows steadily increase in size. It is remarkable how symptomless pulmonary metastases may be often they cause no symptoms until they produce dyspnoea by their very bulk and the effects of pulmonary destruction. Pain and haemoptysis are both unusual.

Solitary metastases in the lung are uncommon and may cause difficulty in diagnosis. They present as single rounded tumours in the radiograph and it is not always possible to differentiate them from a fresh primary peripheral lung tumour (Fig 157). Bronchoscopy sometimes reveals the true nature of the growth. When there is no evidence of metastases elsewhere it is justifiable in some cases to resect the pulmonary lesion. Seiler, Clagett and McDonald (1950) have reviewed the literature on this subject and reported the results of pulmonary resection for metastatic malignant lesions at the Mayo clinic.

## ADENOMA OF THE BRONCHUS

Except for carcinoma bronchial adenoma is far the commonest and most important of the primary pulmonary neoplasms. Although uncommon in comparison with carcinoma individual chest surgeons are accustomed to treat 2 or 3 fresh cases of adenoma every year. Its incidence can only be defined in relation to carcinoma. In the writer's experience adenoma represents 10 per cent of operable pulmonary tumours. Churchill and his colleagues (1950) record 30 pulmonary resections for adenoma in the period 1930-50 in relation to 171 resections for carcinoma. It should be remembered that only about 10 per cent of carcinomas come to resection so it may be inferred that adenomas comprise only about 1/2 per cent of primary neoplasms of the lung.

In contrast to carcinoma adenoma is commoner in women than men and occurs as a rule in a younger age group though it is occasionally observed even in men over 60 years of age.

### Pathology

Kramer (1930) first drew attention to the clinical importance of adenoma of the bronchus although pathological observations had been published earlier. He suggested that the tumour arose in the mucous glands of the bronchus that it







FIG. 156 — Multiple metastases in both lungs from carcinoma of the prostate



FIG. 157 — A 'cannon ball' secondary sarcoma. Similar appearance is often produced by secondary hypernephroma

# ADENOMA OF THE BRONCHUS

FIG 158.—Adenoma of bronchus in a young woman aged 21 years. No symptoms. Not visible on bronchoscopy. Treated by right lower lobectomy.



was of a benign nature but might become malignant occasionally. During the past 20 years there has been a vast amount of discussion on the subject on the origin of the tumour and on its potential malignancy, but little real advance has been made on the views originally expressed by Kramer. Womack and Graham (1938) and Graham and Womack (1945) have held that bronchial adenomas arise from a foetal bronchial bud that they have a dangerous tendency to malignancy and should be removed by total pneumonectomy.

Much confusion has been caused by the divergent views of pathologists who have reported adenomas as carcinomas and *vice versa*. Clinically there is no doubt that there is a group of cases of innocent epithelial tumours of the bronchus that run a benign course and these tumours can be recognized by experienced pathologists. That there should be occasional examples of a malignant change is not a matter for surprise or acceptance of universal malignancy. Foster Carter (1941) has pointed to the similarity between these lesions and the tumours of the parotid gland. It is of the greatest importance to the patient that bronchial adenoma should not be confused with bronchogenic carcinoma for the prognosis of the latter is on the whole poor whereas the prognosis in adenoma is excellent provided correct diagnosis and treatment are applied.

The tumours may occur in the trachea, but they arise as a rule in a major bronchus at or near a primary division. They are thus usually visible on bronchoscopy, occasionally they arise peripherally, and are beyond bronchoscopic access (Fig 158). They may be entirely endobronchial but more commonly there is also an



FIG 159—Adenoma of bronchus showing distension of the bronchus by the tumour distal collapse, fibrosis and bronchiectasis Operation specimen



FIG 160—Bronchial adenoma ( $\times 250$  Reduced one quarter on reproduction)

extra bronchial extension. The tumour is polypoid and is covered by intact mucous membrane. As it grows it distends the bronchus and compresses the surrounding tissues (Fig 159). Histologically the tumour is covered on its exposed surface by bronchial epithelium from which it is separated by a layer of connective tissue. The cells of which the tumour is composed are arranged in solid acini, they have a remarkably orderly and uniform appearance. Mitotic figures are rare (Fig 160).

#### Clinical course

As the tumour grows it obstructs the bronchus, and like certain forms of carcinoma it gives rise to bronchial obstruction, with complications due to bronchiectasis and infection (Fig 161). Before complete obstruction occurs it may give rise to symptoms such as cough, haemoptysis and dyspnoea. Once infection has supervened the sputum becomes purulent and signs of toxæmia such as fever, malaise and loss of weight occur.

The correct diagnosis can only be made by biopsy, otherwise the signs and symptoms closely resemble those of carcinoma. Adenoma may be suspected as opposed to carcinoma by the length of history, which may extend over many years, and by the youth and sex of the patient. Haemoptysis is a commoner symptom in adenoma than in carcinoma, at times it may be severe and it may occur as the only symptom with long intervals of freedom, or it may be associated with attacks of so called "pneumonia".



Fig 161 — Bronchial adenoma causing collapse of the right lower and middle lobes with secondary infection and negative changes in the upper lobe

## Treatment

Before pulmonary resection became a relatively safe operation bronchial adenoma was treated by bronchoscopic methods by diathermic fulguration and radon implantation. These methods although satisfactory in a few early cases failed to eradicate the extra bronchial portion of the growth and failed to deal with the suppurative complications distal to the growth. They have been superseded as a general rule by open operation. At thoracotomy it is possible to define the exact extent of the tumour. If it is entirely endobronchial the bronchus may be opened and a complete local excision of the growth may be carried out. The lung is sutured and no loss of pulmonary tissue occurs. More commonly the lung distal to the tumour is irreparably damaged by obstruction and infection and it is necessary to perform a lobectomy and even in some cases a total pneumonectomy.

## HAMARTOMA OF THE LUNG

### Chondromatous hamartoma

The term hamartoma was coined by Albrecht (1904) to apply to certain tumour like malformations in which there is an abnormal mixing or development of the normal components of an organ. Although the term was not originally used by Albrecht for lesions of the lung it seems an apt word that has become generally accepted for certain benign tumours of the lung. The incidence of this type of benign lesion is second only to bronchial adenoma. The commonest pulmonary hamartoma is a tumour like mass that contains



FIG 162—Chondromatous hamartoma. The glistening white chondroma is enveloped in a well defined capsule




FIG 163—Chondromatous hamartoma showing cartilage fibrous tissue and clefts lined by epithelium ( $\times 44$  Reduced one quarter on reproduction)

elements of normal pulmonary tissue. The mass is likely to contain cartilage, epithelium, fibrous tissue, fat and vascular elements. They are liable to be described as tumours of their most dominant tissue component. The most common and characteristic hamartoma is mainly cartilaginous and may be reported as a chondroma (Fig 162). A careful histological examination reveals that in addition to cartilage, there are clefts lined by columnar or cuboidal epithelium and separate areas of fibrous and fatty tissue (Fig 163).

The rate of growth of these lesions has not been determined, it may be that growth occurs only *pari passu* with that of the individual. There is no evidence of malignant change, but such a possibility should not be disregarded. Carlsen and Kiaer (1950) have described the pathology of these tumours in considerable detail.

Cartilage is not necessarily the main constituent, and in some instances the tumour may be described as a fibroma, a fibro adenoma or even a lipoma. Many of the reported examples of this condition have been chance findings at autopsy or in operation specimens. McDonald, Harrington and Clagett (1945) reported 23 cases, of which 20 were found at necropsy and 3 were removed surgically. Most of these "tumours" occur in peripheral parts of the lung, they cause no symptoms and require no treatment (Fig 164). When the lesion is endobronchial it causes bronchial obstruction with complications of distal collapse and infection. If the lesion is accessible to the bronchoscope the diagnosis may be proved by biopsy. In these cases, in which the lesion produces symptoms, the correct treatment is

FIG. 164 — Chondromatous hamartoma. A round shadow in a man aged 50 years found on mass radiography. Thought to be a carcinoma and treated by pneumonectomy.



excision. At operation the aim should be to conserve all healthy lung tissue, thus it may be possible in some cases to excise the tumour by bronchotomy; in others a limited lung resection is necessary.

#### Vascular hamartoma

The vascular hamartomas constitute quite a separate group of developmental errors or aberrations. In recent years there has been a great increase in the literature on this subject and it seems clear that vascular lesions of the lung are more common than was believed formerly. A number of different terms have been employed in the nomenclature. The lesions may range from small telangiectases scattered throughout both lungs to a solitary pulsating vascular mass composed of a tortuous dilated branch of the pulmonary artery which communicates directly with a pulmonary vein. It is because of the wide range in the anatomical variety of these lesions that terms such as pulmonary telangiectasis, cavernous haemangioma, arteriovenous aneurysm and arteriovenous fistula have been employed.

The multiple lesions may be associated with evidence of telangiectases elsewhere, particularly on the face, the lips and the buccal mucous membrane. The writer has observed one example in a patient with hereditary familial telangiectasis. Goldman (1947) has reported the condition in two brothers. Haemoptysis is not necessarily a common symptom, though it does occur in some cases and has been recorded as a fatal complication.



FIG 165 —(a and b) Vascular hamartoma (multiple cavernous haemangiomas). In definite rounded shadows in both lower lobes in a woman who showed the classical signs of cyanosis, polycythaemia and clubbing with telangiectasis of the face and lips.



FIG 166—Vascular hamartoma (solitary arteriovenous fistula) in a young man with dyspnoea, cyanosis and clubbing. Treated by right lower lobectomy.



The signs and symptoms depend in their severity on the degree of shunt of deoxygenated blood from the pulmonary artery to the pulmonary vein. Maier and his associates (1948) record a case in which 58 per cent of the total cardiac output at rest passed through the abnormal arteriovenous communication. According to the proportion of blood that fails to become oxygenated in the lungs there are signs of cyanosis, polycythaemia and clubbing of the fingers. As a result of anoxia the patients are breathless and many suffer from attacks of faintness. These signs may suggest a diagnosis of congenital heart disease such as the tetralogy of Fallot and they have in the past no doubt led to an erroneous assumption that the patient suffers from polycythaemia vera. The key to correct diagnosis lies in a careful examination of the lungs. Auscultation over the lesion will often disclose a continuous murmur that becomes louder on inspiration. Radiology will reveal in the lung field a shadow or shadows which increase in size on inspiration (Fig 165). Kymography and screening will show expansile pulsation and tomography frequently demonstrates the continuity between the lesion and the vascular shadows of the pulmonary artery and veins. Angiography with a contrast medium is a refinement of diagnosis which is seldom if ever required.

The only treatment possible for these lesions is excision. The need for treatment depends on the severity of the symptoms. If the arteriovenous shunt in the pulmonary circulation is small there may be no disability and there is no urgent need for operation. If the lesions are multiple and bilateral total excision is impossible, considerable symptomatic improvement can be attained by excision.



of the major lesions in which a shunt occurs. In the case of a large solitary arteriovenous fistula a cure can be expected by lobectomy, and this is the operation of choice (Fig 166). In some instances such as that reported by Jones and Thompson (1944) total pneumonectomy is required.

Local excision of the lesions, even when large, is possible if temporary clamps are applied to the pulmonary artery and veins, but there is a risk of cerebral embolism from a clot in the proximal part of the ligated vein. This accident has occurred in the experience of the writer. Jones (1944) was the first to report a case treated successfully by bilateral local excision.

## NEUROGENIC TUMOURS

Tumours may arise from the nerves that supply the bronchi. They must be considered extremely rare. The writer has observed 2 cases in which a neurofibroma arose in close connexion with a bronchus. Both were removed without the necessity of resecting lung tissue. Diveley and Daniel (1951) report a case in which a neurofibroma was situated in the middle lobe and treated by lobectomy.

## BIBLIOGRAPHY

- Albrecht, E (1904) "Ueber Hamartome" *Verh dtsch path Ges*, 7, 153  
Barnard W G (1926) "The Nature of the Oat celled Sarcoma of the Mediastinum" *J Path Bact.*, 29, 241  
Bjork V O (1947) "Bronchiogenic Carcinoma" *Acta chir scand*, 95, Suppl 123  
Black, H (1950) "Fibrosarcoma of the Bronchus" *J thorac Surg*, 19, 117  
Brock, R. C (1948) "Bronchial Carcinoma" *Brit med J*, 2, 737  
— (1950) "The Early Diagnosis of Bronchial Carcinoma" *Brit med J*, 1, 116  
Brooks, W D W Davidson, M., Thomas, C P., Robson, K., and Smithers, D W. (1951)  
"Carcinoma of the Bronchus A report on the first five years' work of the joint consultation clinic for neoplastic diseases of The Brompton Hospital and The Royal Cancer Hospital"  
*Thorax*, 6, 1  
Carlsen, C J., and Kiaer, W (1950) "Chondromatous Hamartoma" *Thorax*, 5, 283  
Graham, T. H., and Keane, N U (1950) "Fibrosarcoma of the Bronchus" *J thorac Surg*, 19, 117  
J G (1950) "The Surgical Management  
of Primary Neurogenic Tumours of the Lung"
- Occurring  
Carcinoma  
Adenoma"
- J thorac Surg*, 14, 100  
Griffith, E. R., McDonald, J R., and Clagett, O T (1950) "Alveolar-cell Tumours of the Lung"  
*J thorac Surg*, 20, 949  
Hutchinson, R (1928) *The Alleged Increased Frequency of Primary Carcinoma of the Lung*  
by Local Resection  
*Lancet*, ii, 50

# BIBLIOGRAPHY

- Jones J P and Thompson W P (1944) Arterio venous fistula of the Lung *J thorac Surg* 13, 357
- Kramer R (1930) Adenoma of the Bronchus *Ann Otol etc St Louis* 39 689
- Machle W and Gregorius F (1948) Cancer of the Respiratory System in the United States Chromate producing Industry *Publ Hlth Rep Wash* 63 1114
- Maier H C (1948) Primary Lymphosarcoma of Lung *J thorac Surg* 17, 841
- Maier H C Himmelstein A Riley R L and Bunum J J (1948) Arterio-venous fistula of the Lung *J thorac Surg* 17, 13
- Mason G A (1949) Cancer of the Lung Review of a thousand cases *Lancet* 2, 587
- McDonald J R Harrington S W and Clagett O T (1945) Hamartoma (often called
- Perry K M A (1947) Diseases of the Lung resulting from Occupational Dusts other than Silica *Thorax* 2 91
- Raeburn C (1948) Primary Carcinoma of the Lung *Lancet* 2 474
- Seile (1948) Primary Resection for Metastatic
- Taylor (1948) Resection for Bronchial Carcinoma *Thorax* 5 257
- Womack N and Graham E A (1938) Mixed Tumors of the Lung so-called bronchial or pulmonary adenomata *Arch Path* 26, 165
- Wyburn Mason R (1948) Bronchial Carcinoma presenting as Polyneuritis *Lancet* 1, 203
- Wynder E L and Graham E A (1950) Tobacco Smoking and Bronchiogenic Carcinoma *J Amer med Ass* 143, 329

## CHAPTER 26

### BORNHOLM DISEASE

KENNETH M A PERRY

This epidemic disease, variously known as epidemic pleurodynia, epidemic myalgia, or "The Devil's Grip" is characterized by the sudden occurrence of violent pain in the lower part of the thorax which tends to spread to the abdomen. The incubation period is 2-5 days and patients remain infective as long as the pain persists. The disease was first described in Norway in 1872 by Daae, while two years later Finsen reported that outbreaks had occurred in Iceland in 1856 and in the Baltic. Since this time reports of many outbreaks have been reported, both in Great Britain and in the United States of America. The disease usually occurs in warm weather, when there is a considerable period of continuously high humidity. It occurs mostly in young adults and rather more commonly in females than males. The onset is abrupt with pain in the epigastrium and hypochondriac region, which spreads into one side of the chest and is accentuated by breathing. Ronald (1942) noted that in addition to the abdominal muscles, there was particular involvement of the muscles around the shoulder girdle and neck. The affected muscles are tender and swollen, and the disease is accompanied by fever usually up to 101° F, but sometimes higher, while frontal headache occurs in more than half the cases, and occasionally vomiting, diarrhoea, and epistaxis. Paraesthesia in the arms and hands occurs especially in females. There are rarely any signs in the chest, and the spleen is not enlarged. Jameson and Prinsley (1947) and Gray and Carter (1948) record cases in which a pleural friction rub was heard, and the latter authors recorded a case in which there was palpable fremitus. In a quarter of the cases one or more relapses takes place. The tenderness and swelling of the muscles and pain on exertion may last several weeks or months after the attack has subsided. Complications such as pleurisy in rare cases with effusion, conjunctivitis, orchitis, pericarditis, pneumonia and otitis media have been reported but mostly the disease follows a benign course and results in complete recovery. Nichamin (1945) noted a marked reduction in haemoglobin level, while the red cells numbered less than four millions. The white cell counts seem to vary widely, with no particular abnormality in the differential counts. The sedimentation rate is usually raised. Screening the chest reveals a freely moving diaphragm, and skiagrams show no abnormalities in the lung fields. The disease is of considerable importance because of the differential diagnosis from pleurisy.

The aetiological organism has not yet been identified but the Annual Report of the U.S. Public Health Service for 1944, records the fact that the blood, spinal fluid and nasal washings of the human cases of the disease have yielded an agent pathogenic to monkeys. Findlay and Howard (1950) carried the matter a stage

## BIBLIOGRAPHY

further by showing that sera from patients fixed complement with Coxsackie virus 2 antigen

Patients should be treated by being kept in bed until 48 hours after the pyrexia and pain have subsided. There is no specific treatment but ordinary methods should be used for the relief of pain

## BIBLIOGRAPHY

- Ann Rep U S P bl Hlth Serv* (1944) Washington D C
- Davies J B M and Warin J F (1951) "An Epidemic of Bornholm Disease" *Brit med J* 2 948
- Findlay G M and Howard E M (1950) "Coxsackie Viruses and Bornholm Disease" *Brit med J* 1 1233
- Gray J D and Carter F S (1948) "Acute Benign Dry Pleurisy Six cases in a hospital staff" *Lancet* 2 254
- Harder F K (1936) "Epidemic Myalgia or Pleurodynia in Southwestern Ohio" *Amer J med Sci* 191 678
- Howard C R G (1938) "Epidemic Myalgia Five cases in one household" *Brit med J*, 2 1203
- Howard T Weymuller C A Edson J Ittner E Watson J and Cassidy M L (1943) "Epidemic Pleurodynia in Brooklyn in the Summer of 1942" *J Amer med Ass* 121 925
- Jamieson W M and Prinsley D M (1947) "Bornholm Disease in the Tropics" *Brit med J* 2 47
- Nicham S J (1945) "Clinical and Epidemiological Aspects of Epidemic Pleurodynia" *J Amer med Ass* 129 600
- Pickles W N (1933) "Bornholm Disease Account of a Yorkshire Outbreak" *Brit med J* 2 817
- Ronald J (1942) "Bornholm Disease" *J Roy Na med Serv* 28 144
- Sylvester E (1934) *Epidemic Myalgia* English translation London Oxford University Press
- Smith R E (1937) "Cervical Myalgia in Adolescents" *Clin J* 66, 330

## CHAPTER 27

### TUMOURS OF THE CHEST WALL

GEOFFREY FLAVELL

TUMOURS arising from the chest wall may be divided into those that are superficial and those that are deep the former, comprising mainly innocent lesions such as cysts, naevi and lipomas, originate in the skin and subcutaneous tissues and are therefore not fixed, while the latter tend to involve the ribs and intercostal muscles are commonly fixed, and must all be regarded either as malignant or potentially so. Neither group differs materially from those growths to be found elsewhere on the body surface, or in skeletal structures, and those classed as superficial do not warrant further description, but the deep and fixed variety pose, from their situation, particular diagnostic and surgical problems. In a recent review of 48 such tumours, Blades and Paul (1950) found 28 benign (of which at least 7 were potentially malignant), and 20 (47 per cent) malignant; while it has been estimated that primary malignant tumours of the ribs and sternum comprise some 10 per cent of all primary malignant bone tumours.

The symptoms caused by chest wall growths are principally pain and swelling. Pain may long precede the latter, since the tumour may grow into the pleural cavity rather than present upon the surface, and it is usually boring in nature, sometimes pleuritic, or may be referred along the segmental nerves to produce abdominal pain, leading to a mistaken diagnosis of gall bladder or other intra-abdominal disease. Blades and Paul found an average of 6 months elapsed between the onset of such pain and x-ray investigation of the affected part, 2 months of which was spent in the treatment of "neuritis". When pain is constant and severe, malignancy is to be feared, on the other hand tumour masses may be quite painless and attain great size. The frequency with which these lesions are associated by patients with previous trauma is well-known, but it is doubtful if a true relationship exists, although pain (or swelling) persisting in a rib for any considerable period after injury should be most carefully investigated. Idiopathic fractures occasionally occur in the ribs, most commonly in the first, second, seventh and eighth. These have been attributed to stresses resulting from coughing and called "cough fractures" (Annotation 1946). Their importance lies in their superficial resemblance to a spontaneous fracture through a tumour of rib with which they must not be confused.

In the past many "endotheliomas" of the pleura, usually accompanied by blood-stained pleural effusions, have been described. There is now little doubt that the great majority of these were in fact undifferentiated bronchial carcinomas which invaded the pleura at an early stage and spread rapidly upon its surface, and that true endotheliomas (or mesotheliomas) arising from the pleura itself are exceptionally rare, if indeed they occur at all. The soft tissues of the chest rise to fibromas, lipomas, or fibrosarcomas which, though they may be adjacent to the pleura, or involve it, but differ in no way

## PRIMARY TUMOURS OF THE RIBS AND STERNUM

neuromas or neurilemmomas arise from intercostal nerves expand the intercostal spaces and cause smooth pressure erosion of related ribs. They may be isolated or part of generalized von Recklinghausen's disease while those much more common situated in the paravertebral gutter, are classed among the tumours of the mediastinum. It remains therefore to consider those growths which principally arise in the bony structures of the thorax.

### Secondary growths

Metastases especially from carcinoma of the thyroid kidney and breast are the commonest of chest wall tumours comprising about 38 per cent of them. They may be more conspicuous than their primary source and difficult to distinguish radiologically from intrinsic tumours of rib. When very vascular they sometimes present as pulsating masses to be differentiated from aortic aneurysm which likewise may produce a thoracic swelling which is not always pulsatile usually on the right of the sternum. Growths of the mediastinum in particular dermoids have on occasion a pre sternal element linked by a dumb-bell extension to the mass within and carcinoma of the lung often invades the chest wall especially in the superior sulcus where it produces the painful Pancoast syndrome.

### Inflammatory lesions

Specific and inflammatory lesions to be distinguished are tuberculous or pyogenic osteomyelitis empyema necessitatis or more rarely typhoid osteochondritis or gummas. Tuberculous lymphatic glands situated at the anterior ends of intercostal spaces often break down into a tense and fixed small cold abscess which is none the less fluctuant and from which sterile pus can be aspirated. Other disorders that occasionally simulate growths are mal united fractures congenital or rachitic deformities or marked unilateral protusion of the bony cage due to scoliosis. A non specific non suppurative condition causing painful swelling usually of one costo-chondral junction is Tietze's disease. Seldom encountered in civil practice it is remarkably frequent among young military recruits and often leads to unnecessary intervention. The wearing of packs has been dubiously blamed for its appearance. It runs a long course the swelling altering in size but never breaking down while histologically some fibrotic change is found in the cartilage with periarticular swelling of the soft tissues.

### Primary tumours of the ribs and sternum

Most tumours of the ribs and sternum grow from cartilage and can be classed either as chondromas or chondrosarcomas. These growths comprised 38 per cent of the series of 81 collected by Sommer and Major in 1942 and 65 per cent of Dorner and Macey's 23 examples in 1948. The ribs are the site in about 80 per cent of neoplasms the sternum 20 per cent while of cartilaginous tumours some 68 per cent are to be found situated near the costo-chondral junction the manubrium being the most commonly affected portion of the sternum. A good deal of confusion in nomenclature exists such terms as osteochondroma osteochondro sarcoma and chondromyxosarcoma tending to disguise the essentially cartilaginous character of the growths in describing bone formation or mucoid degeneration both of which are merely secondary features. Osteogenic sarcoma predominating




FIG 167 (a and b)  
A large chondroma  
involving the pos-  
terior end of the  
left 10th rib

(a)



(b)

# PRIMARY TUMOURS OF THE RIBS AND STERNUM

over chondrosarcoma elsewhere is conspicuously rare in the chest as is also the osteoclastoma a total of only 9 instances being recorded in literature

Cartilaginous tumours are hard and nodular with occasional softer or even fluctuant areas If the history is long as it frequently is they may attain considerable size X ray examination is seldom conclusive showing merely destroyed bone or an expanded and thinned out cortex though irregular flecks of calcification are common and typical (Fig 167) A firm distinction between the innocent and the malignant can in many cases be made only histologically Indeed there is a dangerous tendency to underrate the malignant potentialities of apparently innocent chondromas for mitoses are exceptional while others are clearly atypical nuclei or giant cartilage cells must be recognized as indicating malignancy Similarly some portions of a tumour may seem innocent while others are clearly invasive Enchondral bone formation suggests low malignancy while the presence of mitoses implies high malignancy Most tumours are encapsulated to some extent and even malignant ones can often be shelled out with deceptive ease The advancing edge of the growth remains discrete first displacing and distorting adjacent soft tissues then invading them Local recurrence often repeated follows inadequate removal and may show increased malignancy Metastases are unusual appearing only late in the disease and being generally blood borne Chondrosarcoma is known to develop from innocent chondroma and since a pre operative diagnosis cannot with certainty be made all should be radically extirpated along with adjoining muscle pleura or any related structures Defects resulting in the chest wall permit paradoxical respiration and must therefore be repaired either by interposing rib grafts from above or below by rigid prostheses or, if small by transposing periosteum across them so that new bone is formed X ray therapy is useless in these as in most other chest wall tumours with the exception of Ewing's sarcoma and those arising from lymphoid tissue

The next common cause of rib tumour is the fibrous dysplasia described by Schlumberger Symptoms may be altogether lacking but a history of trauma is often given as the cause of the swelling The rib is moderately expanded x ray examination revealing a fusiform area with semi translucent centre surrounded by the narrowed cortex (Fig 168) Histologically the medulla is replaced by masses of young fibrous tissue in whorls and trabeculae with patches of bony metaplasia Malignant change does not occur and providing always that a positive diagnosis has been made local excision of the affected rib suffices

Plasmocytoma and single myelomas of rib are mentioned in most series and may be more common than recorded numbers suggest (Fig 169) The latter have the character of myelomas elsewhere in the body including a raised plasma protein though Hence Jones protein is not usually present in the urine until late in the disease A similar tumour accompanies eosinophilic granuloma and may like myeloma be solitary or multiple involving in the latter case all the bones except the hands and feet Male children and adolescents are chiefly affected the symptoms of local tenderness and sometimes pathological fracture being of short duration Leucocytosis with eosinophilia pyrexia and loss of weight lead to the diagnosis and the prognosis following excision is good The condition is believed by some to manifest the same basic disorder as Letterer-Siwe or Hand-Schüller-Christian



# TUMOURS OF THE CHEST WALL




FIG 168 —Fibrous dysplasia involving the posterior portion of the right 9th rib




FIG 169 —A solitary myeloma in the mid-portion of the right 10th rib

# BIBLIOGRAPHY

disease Osteitis fibrosa, which is sometimes caused by a parathyroid tumour when the blood calcium will be raised, will on occasion cause one or more tumours in the bony thoracic cage  
Ewing's tumour in ribs is rare, only 18 cases having been reported up to the end of 1942, but when occurring it is found usually in patients during the second decade of life, and should be treated by deep x ray therapy

## BIBLIOGRAPHY

- Annotation (1946) Idiopathic Fracture of the First Rib *Brit med J* 1, 493  
Blades B and Paul J S (1950) Chest Wall Tumours *Ann Surg* 131, 976  
Dorner R A and Marcy D S (1948) Primary Rib Tumours *J thorac Surg* 17, 690  
Gill A M Jones R A and Pollak L (1942) Tietze's Disease *Brit med J* 2, 155  
Hedblom C A (1921) Tumours of the Bony Chest Wall *Arch Surg* 3, 56  
Janes R M (1939) Primary Tumours of Ribs *J Thorac Surg* 9, 145  
Kinsella T S White S M and Koucky R W (1947) Two Unusual Tumours of the Sternum *J Thorac Surg* 16, 640  
Kipp H A and Fisher E R (1951) Eosinophilic Granuloma of Rib *J thorac Surg* 21, 24  
Maurer E and De Stefano G A (1948) Eosinophilic Granuloma of Rib *J Thorac Surg* 17, 350  
O Neal L W and Ackerman L (1951) Eosinophilic Granuloma of Rib *J Thorac Surg* 21, 71  
Sommer G N S and Major R C (1942) Cartilaginous Tumours of Ribs and Sternum *Surg* 115, 51  
Zinninger M (1930) Neoplasms of the Bony Thoracic Wall *Ann Surg* 92, 1013  
Tumours of the Wall of the Thorax

## CHAPTER 28

### DISEASES OF THE MEDIASTINUM

KENNETH M A PERRY

THE MEDIASTINUM occupies the middle of the chest and lies between the two lungs. In it lie the trachea, oesophagus and heart. Diseases of the oesophagus and heart are usually considered in treatises on the digestive tract or cardiology and will not be discussed in this book. Enlargement of mediastinal glands results from reticulosis, sarcoidosis and secondary carcinoma and is discussed in the chapter dealing with these subjects.

#### MEDIASTINAL DISPLACEMENTS

In a healthy patient the mediastinum is freely mobile and is readily displaced by disease in either lung. Even lying on one side causes the mediastinum to shift to the dependent position. The mediastinum shifts towards the side of the lesion when there is fibrosis in the lung or when collapse occurs, and it is pushed away from the side of the lesion when air or fluid is present in the pleural space. In certain cases of pneumothorax there may be ballooning of the pleural cavity between the aorta and the oesophagus giving rise to a mediastinal hernia (Fig. 170). Displacement by fibrosis occurs particularly in unilateral upper lobe fibroid phthisis, while collapse giving rise to bronchiectasis or caused by neoplasm is also a common cause of shift of the mediastinum. Perhaps the most common cause, however, is a pleural effusion and if this is present without shift it may be suspected that there is an underlying carcinoma fixing the mediastinum. Expanding tumours and aneurysms of the aorta are other conditions which may cause the mediastinum to be displaced. In kypho scoliosis there is gross distortion and twisting of the mediastinum which is nearly always associated with severe chronic bronchitis and emphysema.


Mediastinal displacement gives rise to no symptoms, unless it occurs suddenly, when the patient may complain of sudden collapse and weakness. On examination the trachea will be felt to be displaced and the apex beat and area of cardiac dullness will also be displaced. Fluoroscopy is sometimes of great value since it may show a swinging shift of the mediastinum during respiration, which is not shown by the straight skiagram. Normally, however, mediastinal shift is quite evident in a skiagram of the chest. Mediastinal displacement in itself does not require treatment. This should always be directed to the underlying condition which has given rise to the shift.

#### MEDIASTINAL EMPHYSEMA

Air may reach the mediastinum along the fascial planes of the neck, through a perforation of the trachea, oesophagus or bronchus, from the retro peritoneal space, or from the interstitial tissues of the lung, and it is theoretically possible

# MEDIASTINAL EMPHYSEMA

FIG. 170.—A mediastinal hernia of left pneumothorax to right side of chest in a patient with bilateral artificial pneumothoraces



for it to happen by dissection of air between the parietal pleura and the chest wall or between the visceral pleura, or by passage of air from the pleural cavity to the mediastinum. From the practical point of view the ways that need consideration are, spread along the fascial planes of the neck, and interstitial pulmonary emphysema. It is well known that the former may occur, and mediastinal emphysema and bilateral spontaneous pneumothorax are sometimes complications of operations on the neck, particularly thyroidectomy and tracheotomy, and more rarely tonsillectomy and other operations. Since, however, these operations are always accompanied by anaesthesia through intratracheal tubes, and tracheal and bronchial obstruction and frequently coughing are all familiar causes of interstitial emphysema of the lung, it seems probable that these complications often result from rupture of pulmonary alveoli rather than direct aspiration of air into the mediastinum. The causes of interstitial pulmonary emphysema are first, traumatic, either as a result of injury to the chest, with or without fracture of ribs, or from operations of the chest including the induction of an artificial pneumothorax. Secondly, increase of pulmonary pressure caused by straining with a closed glottis, as may occur in lifting heavy weights, straining at a stool, during child birth (Fig. 171) or during attempts to resuscitate the new born, or as a result of partial or complete occlusion of the trachea or bronchus accompanied by coughing, as may occur during anaesthesia, in asthmatic attacks, in chronic bronchitis, from inhaled foreign bodies, or during the course of whooping cough or pneumonia. Thirdly, it may result from the spontaneous rupture of the alveoli. This latter condition is of





FIG. 171 — (a) Mediastinal emphysema which developed in a woman during labour (b) lateral view showing the emphysema under the sternum

(a)



(b)

particular interest in that it may occur in an apparently healthy individual without any evidence of increased intra alveolar pressure. Since the condition has a tendency to recur it seems possible that the cause is an inherited defect in the tissue of the alveolar wall.

Mediastinal emphysema may remain confined within the mediastinum or it may escape into the subcutaneous and deep tissues of the neck through the diaphragmatic openings into the retro peritoneal tissue and into the pleural cavity. A small amount of air in the mediastinum will probably cause only trivial symptoms. Larger amounts will cause pain in the chest and difficulty in breathing and the patient may lie flat in bed trying to suppress his cough. Further increase in the mediastinal pressure gives rise to symptoms similar to those of increased pericardial pressure: the veins collapse and the flow of blood into the heart is diminished. If the pressure rises sufficiently high the circulation may come to a standstill as this happens shortness of breath, cyanosis and pulmonary oedema may develop.

On physical examination of the patient if there is subcutaneous emphysema of the neck it will be evident to the touch. The most characteristic physical sign of mediastinal emphysema however consists of crackles, bubbles or churning sounds heard with each contraction of the heart. These may be heard only during systole but sometimes they occur throughout the whole cardiac cycle with accentuation during systole. Besides these characteristic signs there may be obliteration of the cardiac dullness and perhaps evidence of increased mediastinal pressure presenting as dyspnoea, cyanosis, engorged veins and circulatory failure. If pneumothoraces are present the signs of that condition will be evident. Skia-grams of the chest may show collections of air quite plainly but sometimes only a few streaks may be seen in the mediastinal tissue between the heart and the chest wall in the lateral view. A sharply defined translucent band outlining the pericardium especially along the upper part of the left border is a characteristic finding. Pneumothoraces are evident if they are present.

### Treatment

A small amount of air in the mediastinum is of no consequence as the air is soon absorbed. If tension pneumothoraces are present it is important to insert a needle into the pleural cavity and attach it to an under water drain. If pressure on the mediastinum itself gets high a catheter inserted into it through the jugulum may give relief. If there is cardiac embarrassment this may be treated with oxygen and stimulants given to combat anoxia and shock. Morphine should be given to relieve pain.

### ACUTE MEDIASTITIS

Acute mediastinitis may occur with or without suppuration. Simple mediastinitis mostly results from lacerations of the oesophagus and trachea or from injuries of the chest wall or sternum which cause haemorrhage. It may however spread from inflammatory processes in the lungs, pleurae, pericardium, peritoneum, sternum or vertebrae. Suppurative mediastinitis is a more common condition and Neuhof and Jemern (1943) and Pearse (1938) have shown that about half the cases result from trauma particularly perforation of the oesophagus. This may occur spontaneously in cases of carcinoma of the oesophagus but more often it is

## DISEASES OF THE MEDIASTINUM

the result of the passage of instruments, oesophagoscopes, gastroscopes or stomach tubes. It may also result from impacted foreign bodies especially rabbit or chicken bones. A further 25 per cent of cases are the result of spreading infection from the neck, acute pharyngitis, retropharyngeal abscess, quinsies and suppurative cervical adenitis. Other causes include the spread from adjacent organs, thus pneumonia, bronchiectasis and lung abscess may cause acute mediastinitis, as may perostitis and osteomyelitis of the vertebrae, sternum or ribs, and suppuration in the mediastinal glands. Spread may occur from pleura, pericardium or peritoneum. Suppuration in a hydatid or dermoid cyst may break into the mediastinum. Lastly, the infection may be blood borne in cases of bacterial endocarditis, pyaemia, erysipelas or typhoid fever.

### Pathology

In simple mediastinitis enlargement of the lymph nodes with hyperaemia of mediastinal connective tissues and inflammatory oedema are the only pathological changes. When suppuration occurs it may be limited to one anatomical subdivision of the mediastinum or may spread from one part to another. It may spread up into the neck, down to the abdomen or point through the chest wall. Abscesses have also been known to rupture into the oesophagus, trachea, aorta, pleura or pericardium.

### Symptoms

These vary according to the cause and the virulence of the infection, there are those symptoms caused by infection and those caused by pressure. Following perforation of the oesophagus symptoms will appear immediately, in other causes the onset may be more insidious. There is severe pain under the sternum radiating to the throat, back and shoulders associated with dysphagia, dyspnoea, fever and sometimes rigors. There may be cough with purulent sputum, and the cough may be brassy as a result of pressure on the trachea. Other pressure symptoms include hoarseness from involvement of the left recurrent laryngeal nerve, neuralgic pains from pressure on the intercostal nerves or brachial plexus, and superior mediastinal obstruction from pressure on the great veins.

Physical examination will usually reveal little. The patient will look ill and be cyanosed and dyspnoeic. There may be stridor. Enlargement of veins from superior mediastinal obstruction is more common in chronic mediastinitis. There may be painful areas over the spine or sternum. Subcutaneous emphysema may result from perforation of the oesophagus and torticollis from suppuration in the carotid sheath. Local redness, oedema and fluctuation may be caused by the pointing of the abscess. Dullness and distant breath sounds to one side of the sternum are sometimes found. Skiagrams, particularly if they are over-penetrated, may reveal a broadening of the mediastinal shadow, the trachea may be pushed forward and a foreign body may be revealed, on screening a superior mediastinal abscess may appear as a pulsating shadow. Blood examination will always show a polymorphonuclear leucocytosis.

### Prognosis

The prognosis is always grave and the condition is always acute, rapidly progressive and frequently fatal. The abscess may rupture into the oesophagus or

## CHRONIC MEDIASTITIS

a bronchus and this is often providential drainage. The suppuration, however, may spread to cause empyema, pericarditis or peritonitis. Gangrene of the mediastinum, however, may result and is always fatal. The prognosis is best when the disease is localized to the anterior mediastinum.

### Treatment

Foreign bodies should be immediately removed from the oesophagus and great care should be exercised in passing instruments when there is reason to think the oesophagus may be friable. In cervical suppuration attempts may be made to block the retro visceral space by packing the fascial plane. If mediastinal suppuration is already present attempts should be made to drain it as soon as possible, either by a posterior mediastinotomy or by resection of a costal cartilage or part of the sternum. Penicillin, sulphonamides or the appropriate chemotherapeutic agent to which the causal organism is sensitive should always be administered. If the oesophagus is perforated artificial feeding may be given through a Ryles tube, though it is usually more satisfactory to perform a gastrostomy.

## CHRONIC MEDIASTITIS

Chronic fibrous mediastinitis may be the ultimate result of the acute suppurative condition in the mediastinum which has already been described. The condition is most often caused by trauma, which has resulted in haemorrhage and organization, severe acute rheumatism that has caused a pericarditis, polyserositis, Pick's disease, tuberculosis, syphilis, actinomycosis, and Hodgkin's disease.

Chronic mediastinitis frequently results in complete occlusion or thrombosis of the superior vena cava, an occurrence which was first described by Hallet in 1848. This condition, however, may also result from idiopathic thrombosis, thrombophlebitis, and pressure from mediastinal tumours, Hodgkin's disease, retrosternal thyroids and aneurysms. Ochsner and Dixon (1936) reviewed the world literature and collected 120 cases of which 28 were due to chronic mediastinitis, of these, the mediastinitis in 11 cases was caused by syphilis, in 10 by tuberculosis, in 1 by pyogenic infection, in 1 resulted from trauma and in the remaining 5 was of unknown aetiology. Fischer (1904) first recorded actinomycosis as a cause and since then Cope (1938) has suggested that primary thoracic actinomycosis results from the fungus penetrating the wall of the oesophagus. Mountsey (1947) has recently reviewed this subject and cases from other causes have been reported in recent years by Tubbs (1946) and Renbourn (1946).

### Symptomatology

Symptoms mainly result from pressure or obstruction. They include substernal pain, shortness of breath, hoarseness, dysphagia and Horner's syndrome, resulting from pressure on recurrent laryngeal and sympathetic nerves, trachea, bronchi, oesophagus and sternum. Pottenger's sign is a deviation of the tongue to the affected side with fibrillar twitching of the muscle fibres. It may be caused by a reflex connection between the vagus and hypoglossal nerves. The principal symptoms, however, are attributed to superior mediastinal obstruction. This causes swelling of the face, neck and arms, and on examination these will be found to be swollen and oedematous, while the subcutaneous veins on the



front of the chest and abdomen are dilated and the flow in these will be downwards. Carlson (1934) states that obstruction of the superior vena cava is more severe when it occurs below the entrance of the azygos vein and Hinshaw and Rutledge (1942) have shown by infra red photography that enlarged veins extending over the costal margin on to the abdominal wall are seen only when the azygos is included in the obstruction.

X ray examination may help in the diagnosis by showing widening of the superior mediastinal shadow, but this by itself is not sufficient to warrant a diagnosis of chronic mediastinitis in the absence of obstructive symptoms.

### Treatment

Venesection frequently gives temporary relief to patients with chronic mediastinitis. Syphilitic cases should be treated with arsenic, bismuth and mercury as described in the chapter on syphilis. Actinomycosis will respond to treatment with four million units of penicillin daily for 6-8 weeks.

The problem of relief of obstruction of the superior vena cava, but Tubbs (1946) states that it is not warranted, as it has been shown to be hazardous and often, if not always, fruitless. In addition there are many cases reported who show that the condition is compatible with a long life without discomfort.

### BIBLIOGRAPHY

- Ackerman L. V. and Bricker E. M. (1941) Mediastinal Emphysema and Bilateral Pneumothoraces following Radical Dissection of the Neck. *Arch Surg* 43, 445.
- Barrie H. J. (1940) Interstitial Emphysema and Pneumothorax after Operation on the Neck. *Lancet* 1, 996.
- Banyai A. L. and Jergens G. H. (1938) Mediastinal Emphysema as a Complication of Artificial Pneumothorax. *J thorac Surg* 8, 329.
- Buford C. G. (1918) The Entrance of Air into the Mediastinum during Operations of the Base of the Neck. *Surg gynec Obstet* 26, 540.
- Carlson H. A. (1934) Obstruction of the Superior Vena Cava: an experimental study. *Arch Surg* 29, 669.
- Cope Z. (1938) *Actinomycosis* p. 141. London: Oxford University Press.
- Ehrenburg G. E. (1932) Emphysema of the Mediastinum as a Complication of Artificial Pneumothorax. *Amer Rev Tuberc* 26, 738.
- Fischer (1904) Quoted by Cope (1938).
- Gaudrault G. L. and Chalmers D. M. (1941) Emergency Treatment of Emphysema of the Mediastinum. *New Engl J Med* 224, 940.
- Gray H. K. and Skinner I. C. (1941) Constrictive Occlusion of Superior Vena Cava: Report of 3 cases in which the patients were treated surgically. *Surg Gynec Obstet* 72, 923.
- Green J. A. (1943) Unusual Sounds Emanating from the Chest. *Arch intern Med* 71, 410.
- Gumbauer B. and Cutler M. M. (1941) Spontaneous Pneumothorax in the New born. *J Amer med Assn* 117, 2050.
- Hallett C. M. (1848) Collateral Circulation in Cases of Obliteration or Obstruction of the Venae Cavae. *Edinb med Surg J* 69, 269.
- Hamman L. (1945) Mediastinal Emphysema. *J Amer med Assn* 128, 1.
- Hinshaw H. C. and Rutledge D. I. (1942) Lesions in the Superior Mediastinum. *J lab clin Med* 27, 908.

## BIBLIOGRAPHY

- Keefler C S (1938) Acute and Chronic Mediastinitis A Study of 60 cases *Arch Intern Med.* 62, 109
- Keis J (1934) Studies of the Origin of Mediastinal Emphysema and Pneumothorax at Operation of the Thyroid Gland " *Munch med Wschr* 81, 669
- Macklin, C C and Macklin M T (1944) Malignant Interstitial Emphysema of the Lungs and Mediastinum as may occur as a Complication in many Respiratory Diseases and other Conditions *Medicine* 23 281
- Michels M W (1939) Pneumothorax and Mediastinal Emphysema as Complications of Tracheotomy *Arch Otolaryng Chicago* 28, 842
- Mountsey J P D (1947) Complete Occlusion of the Superior Vena Cava with Chronic Mediastinitis in a Case of Generalized Actinomycosis *Thorax* 2, 203
- Neffson A H (1943) Tension Pneumothorax with Mediastinal Emphysema after Tracheotomy *Arch Otolaryng Chicago* 37, 23
- Neuhof H and Jemerin E E (1943) *Acute Infections of the Mediastinum* Baltimore Williams and Wilkins
- and Rabin C B (1940) Acute Mediastinitis *Amer J Roentgenol* 44, 684
- Ochsner A and Dixon J L (1936) Superior Vena Caval Thrombosis Review of the Literature and Report of Cases of Traumatic and Infectious Origin *J Thorac Surg.* 5, 641
- Pearse H E (1938) Mediastinitis following Cervical Suppuration *Ann Surg* 108, 588
- Renbourn E T (1946) Thrombosis of the Superior Vena Cava *Thorax* 1, 257
- Tubbs O S (1946) Superior Vena Caval Obstruction due to Chronic Mediastinitis *Thorax* 1 247
- Work W P (1943) Mediastinal Emphysema and bilateral Pneumothorax Complicating Tracheotomy in an Adult *Arch Otolaryng Chicago* 37, 526
- Zeman F D (1945) Occlusion of the Superior Vena Cava due to Syphilitic Mediastinitis Collateral Circulation after 19 years *J Thorac Surg* 14, 330

## CHAPTER 29

### CYSTS AND TUMOURS OF THE MEDIASTINUM

GEOFFREY FLAVELL

UNTIL comparatively recently the mediastinum was regarded almost as an abstract anatomical convention, and it is only in recent years that advances in radiology and in the techniques of thoracic surgery have revealed its practical importance and its pathological diversity. In particular the increasing use of mass radiography has led to the early recognition of a bewildering variety of cysts and tumours, and to their early removal. Differing in their anatomical relationship within the mediastinum and in their histological character, many of these lesions none the less owe their origin to a common factor—the developmental complexity of the region. For it is upon this narrow stage that the more caudal of the branchial arches develop, that the primitive foregut buds off its pulmonary diverticulum, which in turn descends into the thorax branching and dividing into the ultimate elaboration of the bronchial tree, that the enteron itself differentiates, that the heart and great vessels undergo their foetal mutations to reach their adult state, that the pleural and pericardial cavities are fashioned from the infoldings of the simple coelom, and the whole cut off by the formation of the diaphragm from the septum transversum. In addition to those growths which may be attributed to developmental faults, there remain others arising from the great diversity of adult tissues actually within or closely related to the mediastinum, and the richness of its lymphatic connexions in turn make it a frequent site for metastatic deposits from more distant organs, in particular from the lung. Such secondary growths will not be considered here, though they inevitably feature in the differential diagnosis of primary tumours as they may occur where ever lymphoid tissue is to be found in the mediastinum.

Investigation of mediastinal growths must always include good lateral, and if necessary, oblique skiagrams, as well as antero posterior views. Radiological screening is indispensable for testing mobility and watching pulsation. A barium swallow shows displacement or obstruction of the oesophagus, while tomography is sometimes useful in fixing exact relationship and revealing cavities. Bronchoscopy and oesophagoscopy are chiefly of value in excluding other growths, but may fix a point of compression from without. The induction of an artificial pneumothorax or pneumoperitoneum may assist diagnosis. Thoracoscopy has been employed, but is seldom conclusive.

Neuromas, enterogenous cysts, and meningeal growths will be found posteriorly, dermoids, teratomas, cysts of the pericardium, thymomas, and thyroid adenomas anteriorly. But others occur in any situation and much else remains to be differentiated.

Aneurysms of the great vessels, or of the heart, may be confused with new growths, and must be distinguished from them by radiological evidence of

## CYSTS AND TUMOURS OF THE MEDIASTINUM

pulsation, by demonstrating their continuity with the vascular structures, if necessary by angiography, by serological reactions, and characteristic physical signs. The oesophagus, when dilated and full of food, as may happen in long-standing cardiospasm, adopts an S shaped position in the mediastinum, bulging to the right of its upper portion and to the left lower down. In either situation it casts a rounded shadow which may greatly resemble a mediastinal tumour, but can readily be distinguished by a history of dysphagia and a barium swallow. In similar fashion diaphragmatic herniation into, or beside the mediastinum can be recognized by barium followed through the gut. Herniation of a lung across the midline often produces an appearance simulating a cyst wall, and scoliosis or vertebral disease distorts x ray appearances. Radiography of the region in infants and young children is fruitful of equivocal "tumours", since the mediastinum is highly mobile at this age and it is difficult to obtain a strictly centred film. Thus a little displacement or rotation of the heart and other structures casts disproportionate and unexpected shadows not always easy to interpret.

All these possibilities as well as that of carcinoma of an abnormal mediastinal phagus, must be borne in mind when the diagnosis of an abnormal mediastinal shadow is to be attempted, and when each in turn has been eliminated there remains to differentiate the wide field of primary cysts and tumours. Among these, characteristics of appearance and of position will sometimes permit a positive choice, and often an intelligent guess, as to their nature, but rarely indeed can there be certainty of diagnosis, and less frequently still can there be any true assurance as to a tumour's innocence or malignancy. It is generally accepted that some 10 per cent of all innocent mediastinal cysts and tumours eventually undergo malignant degeneration, the proportion being higher among teratomas and some neurogenic growths, and lower in such groups as the simple cysts.

Whether or not eventual malignancy is their fate, all are liable to enlarge, and from their situation to cause such symptoms as dyspnoea by pressure upon the adjacent bronchi or trachea, dysphagia by compressing the oesophagus, engorgement of neck veins and tachycardia by obstruction of the great vessels and displacement of the heart, and eventually all of these by a generalized blockage of the mediastinum.

Cysts are prone to establish fistulous communications with the bronchi or with the pleural cavities, generally combined with secondary infection of their contents. Such complications may carry fatal consequences. Haemorrhage can occur into them. Intrathoracic goitres are likely eventually to undergo toxic degeneration, tumours of the thymus to be associated with myasthenia gravis, enterogenous cysts may develop peptic ulceration and perforate, hydatid cysts disseminate their infestation. It is thus evident, quite apart from diagnostic uncertainty, and from the omnipresent and substantial risk of malignant change, that no case exists for tolerating these tumours, and that all, unless the age or condition of the patient forbids, should be surgically removed.

The relative frequency with which they variously occur is not easy to assess with accuracy since few surgeons trouble to record the commoner varieties, and classification of more unusual types remains unstandardized. There is little

## CHAPTER 29

### CYSTS AND TUMOURS OF THE MEDIASTINUM

GROFFREY FLAVELL

Empirically recently the mediastinum was regarded almost as an anatomical convention, and it is only in recent years that advances in and in the techniques of thoracic surgery have revealed its practical use and its pathological diversity. In particular the increasing use of thoraphy has led to the early recognition of a bewildering variety of tumours and to their early removal. Differing in their anatomical position within the mediastinum and in their histological character, many of these lesions none the less owe their origin to a common factor—the developmental complexity of the region. For it is upon this narrow stage that the more primitive branchial arches develop, that the primitive foregut buds off its foregut diverticulum which in turn descends into the thorax branching and continuing the ultimate elaboration of the bronchial tree, that the enteron itself differentiates, that the heart and great vessels undergo their foetal mutations to reach their adult state, that the pleural and pericardial cavities are fashioned from the infoldings of the simple coelom, and the whole cut off by the formation of the diaphragm from the septum transversum. In addition to those growths which may be attributed to developmental faults, there remain others arising from the diversity of adult tissues actually within or closely related to the mediastinum, and the richness of its lymphatic connexions in turn make it a frequent site for metastatic deposits from more distant organs, in particular from the lung. Such secondary growths will not be considered here, though they inevitably feature in the differential diagnosis of primary tumours as they may occur wherever lymphoid tissue is to be found in the mediastinum.

Investigation of mediastinal growths must always include good lateral and if necessary oblique skiagrams as well as antero-posterior views. Radiological screening is indispensable for testing mobility and watching pulsation. A barium swallow shows displacement or obstruction of the oesophagus, while tomography is sometimes useful in fixing exact relationship and revealing cavities. Bronchoscopy and oesophagoscopy are chiefly of value in excluding other growths but may fix a point of compression from without. The induction of an artificial pneumothorax or pneumoperitoneum may assist diagnosis. Thoracoscopy has been employed but is seldom conclusive.

Neuromas, enterogenous cysts and meningeal growths will be found posteriorly, dermoids, teratomas, cysts of the pericardium, thymomas and thyroid adenomas anteriorly. But others occur in any situation and much else remains to be differentiated.

Aneurysms of the great vessels or of the heart, may be confused with new growths, and must be distinguished from them by radiological evidence of

pulsation, by demonstrating their continuity with the vascular structures, if necessary by angiography, by serological reactions, and characteristic physical signs. The oesophagus, when dilated and full of food, as may happen in longstanding cardiospasm, adopts an S shaped position in the mediastinum, bulging to the right of its upper portion and to the left lower down. In either situation it casts a rounded shadow which may greatly resemble a mediastinal tumour, but can readily be distinguished by a history of dysphagia and a barium swallow. In similar fashion diaphragmatic herniation into, or beside the mediastinum can be recognized by barium followed through the gut. Herniation of a lung across the midline often produces an appearance simulating a cyst wall, and scoliosis or vertebral disease distorts x ray appearances. Radiography of the region in infants and young children is fruitful of equivocal "tumours", since the mediastinum is highly mobile at this age and it is difficult to obtain a strictly centred film. Thus a little displacement or rotation of the heart and other structures casts disproportionate and unexpected shadows not always easy to interpret.

All these possibilities as well as that of carcinoma of the lung or of the oesophagus, must be borne in mind when the diagnosis of an abnormal mediastinal shadow is to be attempted, and when each in turn has been eliminated there remains to differentiate the wide field of primary cysts and tumours. Among these, characteristics of appearance and of position will sometimes permit a positive choice, and often an intelligent guess, as to their nature, but rarely indeed can there be certainty of diagnosis, and less frequently still can there be any true assurance as to a tumour's innocence or malignancy. It is generally accepted that some 10 per cent of all innocent mediastinal cysts and tumours eventually undergo malignant degeneration, the proportion being higher among teratomas and some neurogenic growths, and lower in such groups as the simple cysts.

Whether or not eventual malignancy is their fate, all are liable to enlarge, and from their situation to cause such symptoms as dyspnoea by pressure upon the adjacent bronchi or trachea, dysphagia by compressing the oesophagus, engorgement of neck veins and tachycardia by obstruction of the great vessels and displacement of the heart, and eventually all of these by a generalized blockage of the mediastinum.

Cysts are prone to establish fistulous communications with the bronchi or with the pleural cavities, generally combined with secondary infection of their contents. Such complications may carry fatal consequences. Haemorrhage can occur into them. Intrathoracic goitres are likely eventually to undergo toxic degeneration. Tumours of the thymus to be associated with myasthenia gravis, enterogenous cysts may develop peptic ulceration and perforate. Hydatid cysts disseminate their infestation. It is thus evident, quite apart from diagnostic uncertainty, and from the omnipresent and substantial risk of malignant change that no case exists for tolerating these tumours, and that all, unless the age or condition of the patient forbids, should be surgically removed.

The relative frequency with which they variously occur is not easy to assess with accuracy since few surgeons trouble to record the commoner varieties, and classification of more unusual types remains unstandardized. There is little

## CHAPTER 29

### CYSTS AND TUMOURS OF THE MEDIASTINUM

GEOFFREY FLAVELL

Only recently the mediastinum was regarded almost as an anatomical convention, and it is only in recent years that advances in the techniques of thoracic surgery have revealed its practical pathological diversity. In particular the increasing use of roentgenology has led to the early recognition of a bewildering variety of lesions and to their early removal. Differing in their anatomical position in the mediastinum and in their histological character, many of these lesions owe their origin to a common factor—the developmental origin of the region. For it is upon this narrow stage that the more complex structures develop, that the primitive foregut buds off its ventral wall, which in turn descends into the thorax branching and elaborating the bronchial tree, that the enteron itself differentiates, that the heart and great vessels undergo their foetal mutations to reach their adult state—that the pleural and pericardial cavities are fashioned from the infoldings of the simple coelom, and the whole cut off by the formation of the diaphragm from the septum transversum. In addition to those growths which are attributable to developmental faults, there remain others arising from the replacement of adult tissues actually within or closely related to the mediastinum. The richness of its lymphatic connexions in turn make it a common site for metastatic deposits from more distant organs, in particular from the bronchi. Secondary growths will not be considered here, though they inevitably feature in the differential diagnosis of primary tumours as they may occur where ever lymphoid tissue is to be found in the mediastinum.

Investigation of mediastinal growths must always include good lateral and if necessary, oblique skiagrams, as well as antero posterior views. Radiological screening is indispensable for testing mobility and watching pulsation. A barium swallow shows displacement or obstruction of the oesophagus, while tomography is sometimes useful in fixing exact relationship and revealing cavities. Bronchoscopy and oesophagoscopy are chiefly of value in excluding other growths, but may fix a point of compression from without. The induction of an artificial pneumothorax or pneumoperitoneum may assist diagnosis. Thoracoscopy has been employed, but is seldom conclusive.

Neuromas, enterogenous cysts, and meningeal growths will be found posteriorly, dermoids, teratomas, cysts of the pericardium, thymomas, and thyroid adenomas anteriorly. But others occur in any situation and much else remains to be differentiated.

Aneurysms of the great vessels, or of the heart, may be confused with new growths, and must be distinguished from them by radiological evidence of

FIG. 172—(a) A typical neurofibroma. Note the characteristic position in the paravertebral gutter.  
(b) A right artificial pneumothorax has been induced but the tumour has not descended with the lung showing that it is extrapulmonary.

(a)





however that neuromas top the list, followed by bronchogenic cysts and teratomas, various tumours of lymphoid tissue, pericardial cysts and enterogenous cysts, in that order, with rarer lesions following. If retrosternal thyroid enlargements are included they are probably of all and certainly must be considered among growths of the superior

mediastinal tumours removed in United States Army thoracic surgical period of 3 years, 90 were innocent, 15 malignant. Among the 23 neuromas 23 bronchogenic cysts 14 dermoids and teratomas 14 cysts, the malignant numbered 6 teratomas, 4 cases of Hodgkin lymphoblastomas, 2 thymomas, and a neurosarcoma. Ulrich 60 primary mediastinal tumours removed at the Mayo Clinic but roughly similar proportions.

### NEUROGENIC TUMOURS

In the posterior mediastinum are, in 92 per cent of cases, neurogenic tumours occurring elsewhere in the mediastinum are excessively rare, but common. They are divided broadly into those which originate from the so-called generally called neurofibromas or neurilemmomas, and those in which cell predominate, and which arise from the sympathetic ganglia. In 1944 Kent collected 78 cases from the literature and found 41 (52 per cent) malignant. He added 18, classifying 41 per cent as malignant. Harrison in 1949 recorded 51 neuromas of which 59 per cent were malignant. Godwin and associates in 1950 described 24 cases, 2 (8 per cent) malignant and ganglioneuromas characteristically occupy a position anteriorly, evidently closely related to the sympathetic chain. They are more common in the upper part of the chest than in the lower but are also found in the gutter. The vast majority are encapsulated and appear on x-ray film as smooth rounded or slightly lobulated tumours typically posterior (Fig 172). They are distinguished from intrapulmonary tumours by the induction of an artificial pneumothorax. Not infrequently they cause pressure erosion of adjacent ribs and widening of the intercostal spaces. Sometimes the intervertebral foramina also are widened and the tumour may extend dumb bell fashion into the neural canal causing pressure symptoms on the cord. Such instances are generally ganglioneuromas, but may be malignant. A meningioma may produce a similar lesion and several instances of meningocele presenting in the thorax have been reported. The clinical course may be benign or malignant. In some instances they encircle major vessels or extend into the neck making removal difficult and are much more prone to malignant change than the encapsulated type about 13 per cent undergoing degeneration. It has been estimated that a quarter of all neurogenic sarcomas occur in patients having the stigmata of von Recklinghausen's disease.

for diagnostic reasons. Infected bronchial cysts may give rise to long periods of chronic ill health resembling that caused by severe bronchiectasis, if the lesion is not recognized and removed. Furthermore both Moersch and Clagett (and Everts Graham) have recorded carcinoma developing from a bronchial cyst.

An unruptured cyst presents a smoothly round or oval homogeneously opaque shadow on x ray examination. At this stage it is difficult to make a positive diagnosis since only by exploratory thoracotomy can it be distinguished from any of the solid mediastinal tumours. Bronchoscopy or oesophagoscopy are of little use but aneurysms and oesophageal dilatations are eliminated by radiological means. When however the cyst has established a communication its character at least becomes evident since a fluid level will at once be apparent in it. The cyst wall is also then visible and is seen to be characteristically thin and regular. It is still necessary to differentiate it from a breaking down carcinoma, an abscess, a diaphragmatic hernia, an hydatid or any other cyst. And while the fineness and regularity of its wall may make the correct diagnosis probable it is seldom that it can be made certainly. This uncertainty, especially when infection has already occurred, indicates removal.

Histologically bronchial cysts are lined by cuboidal or columnar epithelium, which may or may not be ciliated. It is sometimes irregular or even trabeculated and the walls contain characteristically bronchial elements in disorderly arrangement such as cartilage, smooth muscle, mucus glands and elastic tissue. The contents may be clear or viscid and gelatinous fluid, or consist of blood, pus and air.

Surgical removal is planned through whichever side of the chest seems to afford best access to the cyst and removal presents no particular difficulty except that especial care must be devoted to closing any bronchial fistula.

### DERMOIDS AND TERATOMAS

No fundamental difference exists between these tumours which share a common origin and a common character. In each all three germinal layers may be represented. As the name implies, however, in the dermoid the ectodermal derivatives greatly predominate and the tumour itself is generally cystic. The teratoma on the other hand is composed of all constituents and is mainly solid. Several theories exist, none wholly satisfactory to explain their source, but that of Collenberg seems best to fit the facts of their position, their nature and their relationship. He holds all due to abnormalities in the development of the third and fourth branchial arches at whose union in the foetal midline all the germinal layers are represented in great complexity and from which groups of primitive cells are carried down into the anterior mediastinum by the descent of the heart and great vessels to which they remain intimately related.

Be this as it may the teratoid tumours are almost invariably found in the anterior mediastinum, only 3 exceptions being known among 250 recorded cases. This is so much the case that any tumour of the anterior mediastinum should be regarded as teratoid until otherwise proved. Furthermore they are intimately related to the anterior surface of the pericardium and to the great vessels which emerge from it, this constituting one of the chief surgical difficulties in removing them.

Simple neurofibromas and ganglioneuromas seldom cause symptoms until they do so by reason of their size, which may become considerable. Neurological lesions or pain are generally absent, and many are first detected by routine chest radiography. Later cough or dyspnoea result as their bulk increases, and it may not be until the third or fourth decade of life or even later. At the same time cystic degenerative changes are common, with myxomatous and xanthomatous changes and haemorrhages into the cysts. Histologically both neurofibromas and neuromas are encapsulated, the former being composed of cells from the sheath of Schwann, neurones, and much fibrous tissue, the latter of mature nerve cells set in a stroma of myelinated and non-myelinated nerve fibres. Metastatic change is quite exceptional.

Surgically they are best approached by a transpleural postero lateral thoracotomy unless they are small when a direct extrapleural removal may be possible. They are covered on the anterior surface by the parietal pleura, behind they lie in the chest wall and which must be incised. Before this is done they appear fixed to the chest wall, but as soon as the pleura is stripped from them it is very easy to mobilize and secure the short pedicle that attaches them to the anterior end of some intercostal space, and either dissect them from the chest wall or resect, if need be, the nerve from which they grow. Bleeding from the closely related intercostal vessels is practically the only surgical difficulty, and is usually easily controlled. Tumours having a dumb-bell extension through the intervertebral foramen into the spinal canal however are not so easily dealt with. These generally present with symptoms of cord compression, and a neurosurgical attack as a rule must precede the removal of the intrathoracic portion.

### BRONCHOGENIC CYSTS

These arise in any part of the mediastinum but are most often closely related to the pleura to which they may be attached by a pedicle, or to the trachea or a major bronchus. If more distally disposed on the bronchial tree they cease to be related to the mediastinum and must be distinguished from other cysts of the lung. They originate essentially in a developmental fault, probably by a "pinching off" process from one of the outgrowing lung buds and may therefore be regarded as first cousins to enterogenous cysts, sequestered pulmonary segments, or even to dermoids. They undoubtedly occur more frequently than published series suggest, although 44 are recorded in the Mayo clinic during the past decade. Most have been recognized in the earlier decades of life and very large ones are sometimes seen in infancy, often associated with other developmental abnormalities, and causing respiratory distress and cyanosis. More usually they remain symptomless until their presence is complicated by infection, coupled with a fistulous communication with the bronchial tree or with the pleura. Persistent cough or dyspnoea may precede this stage, due merely to the pressure effects of the cyst, but once a fistula is established with the bronchus expectoration of the infected contents follows, accompanied by haemoptysis, fever, and of course purulent sputum. Rupture of such a cyst into the pleural cavity results in an empyema which will not heal after simple drainage and indeed this accident has been surgically induced by attempts to aspirate cysts mistaken for encysted effusions or

for diagnostic reasons. Infected bronchial cysts may give rise to long periods of chronic ill health, resembling that caused by severe bronchiectasis, if the lesion is not recognized and removed. Furthermore both Moersch and Clagett (and Everts Graham) have recorded carcinoma developing from a bronchial cyst.

An unruptured cyst presents a smoothly round or oval homogeneously opaque shadow on x ray examination. At this stage it is difficult to make a positive diagnosis since only by exploratory thoracotomy can it be distinguished from any of the solid mediastinal tumours. Bronchoscopy or oesophagoscopy are of little use, but aneurysms and oesophageal dilatations are eliminated by radiological means. When however the cyst has established a communication its character at least becomes evident since a fluid level will at once be apparent in it. The cyst wall is also then visible, and is seen to be characteristically thin and regular. It is still necessary to differentiate it from a breaking down carcinoma, an abscess, a diaphragmatic hernia, an hydatid or any other cyst. And while the fineness and regularity of its wall may make the correct diagnosis probable it is seldom that it can be made certainly. This uncertainty, especially when infection has already occurred, indicates removal.

Histologically, bronchial cysts are lined by cuboidal or columnar epithelium, which may or may not be ciliated. It is sometimes irregular or even trabeculated, and the walls contain characteristically bronchial elements, in disorderly arrangement, such as cartilage, smooth muscle, mucus glands and elastic tissue. The contents may be clear, or viscid and gelatinous fluid, or consist of blood, pus and air.

Surgical removal is planned through whichever side of the chest seems to afford best access to the cyst and removal presents no particular difficulty, except that especial care must be devoted to closing any bronchial fistula.

### DERMOIDS AND TERATOMAS

No fundamental difference exists between these tumours which share a common origin, and a common character. In each all three germinal layers may be represented. As the name implies, however, in the dermoid the ectodermal derivatives greatly predominate, and the tumour itself is generally cystic. The teratoma on the other hand is composed of all constituents and is mainly solid. Several theories exist none wholly satisfactory to explain their source, but that of Collenberg seems best to fit the facts of their position, their nature and their relationship. He holds all due to abnormalities in the development of the third and fourth branchial arches at whose union in the foetal midline all the germinal layers are represented in great complexity and from which groups of primitive cells are carried down into the anterior mediastinum by the descent of the heart and great vessels, to which they remain intimately related.

Be this as it may the teratoid tumours are almost invariably found in the anterior mediastinum only 3 exceptions being known among 250 recorded cases. This is so much the case that any tumour of the anterior mediastinum should be regarded as teratoid until otherwise proved. Furthermore they are intimately related to the anterior surface of the pericardium and to the great vessels which emerge from it, this constituting one of the chief surgical difficulties in removing them.

Simple neurofibromas and ganglioneuromas seldom cause symptoms until they do so by reason of their size, which may become considerable. Neurofibromas or pain are generally absent, and many are first detected by routine radiography. Later cough or dyspnoea result as their bulk increases, and is not be until the third or fourth decade of life or even later. At the same time degenerative changes are common, with myxomatous and xanthomatous and haemorrhages into the cysts. Histologically both neurofibromas and ganglioneuromas are encapsulated the former being composed of cells from Schwann neurones, and much fibrous tissue, the latter of mature Schwann cells in a stroma of myelinated and non myelinated nerve fibres. The histology is quite exceptional. If they are best approached by a transpleural postero-lateral thoracotomy they are small when a direct extrapleural removal may be possible. They are covered on the anterior surface by the parietal pleura, behind the vertebral column and which must be incised. Before this is done they appear to be attached to the chest wall, but as soon as the pleura is stripped from them it is easy to mobilize and secure the short pedicle that attaches them to the chest wall, and either dissect them from the chest wall or resect, if need be, the nerve from which they grow. Bleeding from the intercostal vessels is practically the only surgical difficulty, and is easily controlled. Tumours having a dumb-bell extension through the intervertebral foramen into the spinal canal however are not so easily dealt with. They are generally present with symptoms of cord compression, and a neuro-myelogram must precede the removal of the intrathoracic portion.

### BRONCHOGENIC CYSTS

They are in any part of the mediastinum but are most often closely related to the trachea, which they may be attached by a pedicle, or to the trachea or a bronchus. If more distally disposed on the bronchial tree they cease to be related to the mediastinum and must be distinguished from other cysts of the lung. They originate essentially in a developmental fault, probably by a 'pinching off' process from one of the out growing lung buds and may therefore be regarded as first cousins to enterogenous cysts, sequestered pulmonary segments or even to dermoids. They undoubtedly occur more frequently than published series suggest although 44 are recorded in the Mayo clinic during the past decade. Most have been recognized in the earlier decades of life and very large ones are sometimes seen in infancy often associated with other developmental abnormalities and causing respiratory distress and cyanosis. More usually they remain symptomless until their presence is complicated by infection, coupled with a fistulous communication with the bronchial tree or with the pleura. Persistent cough or dyspnoea may precede this stage, due merely to the pressure effects of the cyst but once a fistula is established with the bronchus expectoration of the infected contents follows accompanied by haemoptysis, fever, and of course purulent sputum. Rupture of such a cyst into the pleural cavity results in an empyema which will not heal after simple drainage and indeed this accident has been surgically induced by attempts to aspirate cysts mistaken for encysted effusions or

for diagnostic reasons. Infected bronchial cysts may give rise to long periods of chronic ill health, resembling that caused by severe bronchiectasis, if the lesion is not recognized and removed. Furthermore both Moersch and Clagett (and Evans Graham) have recorded carcinoma developing from a bronchial cyst.

An unruptured cyst presents a smoothly round or oval homogeneously opaque shadow on x ray examination. At this stage it is difficult to make a positive diagnosis since only by exploratory thoracotomy can it be distinguished from any of the solid mediastinal tumours. Bronchoscopy or oesophagoscopy are of little use, but aneurysms and oesophageal dilatations are eliminated by radiological means. When however the cyst has established a communication its character at least becomes evident since a fluid level will at once be apparent in it. The cyst wall is also then visible, and is seen to be characteristically thin and regular. It is still necessary to differentiate it from a breaking down carcinoma, an abscess, a diaphragmatic hernia, an hydatid or any other cyst. And while the fineness and regularity of its wall may make the correct diagnosis probable it is seldom that it can be made certainly. This uncertainty especially when infection has already occurred, indicates removal.

Histologically, bronchial cysts are lined by cuboidal or columnar epithelium, which may or may not be ciliated. It is sometimes irregular or even trabeculated, and the walls contain characteristically bronchial elements, in disorderly arrangement, such as cartilage, smooth muscle, mucus glands and elastic tissue. The contents may be clear, or viscid and gelatinous fluid, or consist of blood, pus and air.

Surgical removal is planned through whichever side of the chest seems to afford best access to the cyst, and removal presents no particular difficulty, except that especial care must be devoted to closing any bronchial fistula.

### DERMOIDS AND TERATOMAS

No fundamental difference exists between these tumours which share a common origin, and a common character. In each all three germinal layers may be represented. As the name implies however in the dermoid the ectodermal derivatives greatly predominate, and the tumour itself is generally cystic. The teratoma on the other hand is composed of all constituents and is mainly solid. Several theories exist none wholly satisfactory, to explain their source, but that of Collenberg seems best to fit the facts of their position, their nature and their relationship. He holds all due to abnormalities in the development of the third and fourth branchial arches at whose union in the foetal midline all the germinal layers are represented in great complexity and from which groups of primitive cells are carried down into the anterior mediastinum by the descent of the heart and great vessels to which they remain intimately related.

Be this as it may the teratoid tumours are almost invariably found in the anterior mediastinum, *only 3 exceptions being known among 250 recorded cases.* This is so much the case that any tumour of the anterior mediastinum should be regarded as teratoid until otherwise proved. Furthermore they are intimately related to the anterior surface of the pericardium and to the great vessels which emerge from it, this constituting one of the chief surgical difficulties in removing them.

Such a position affords little room for expansion since the rigid barrier of the sternum limits it in front, and as these tumours tend to increase in size during adolescence and early adult life, it is then they are apt to cause pressure symptoms, in general throughout the mediastinum, but particularly upon the heart, so that palpitations frequently accompany them. They may, however, remain symptomless until overtaken by one or other of the complications to which they are prone. Of these, infection is common and occurs readily without any fistulous communication, being presumably blood borne. Should it be slight it results only in fever of unexplained origin, and in a formidable increase of those adhesions already binding the tumour to surrounding structures. Often, however, it results in rupture into a bronchus or into the pleura with disastrous results. Bronchial fistulae on occasion develop without evident infection, so that the contents of a dermoid may be expectorated bit by bit. Thus sebaceous material, and particularly hair, may appear in the sputum, confirming in spectacular manner the diagnosis, or it may be aspirated from the pleural cavity or by direct puncture of a cyst. Sometimes the tumours present in the neck, or beside the sternum, and either spontaneously or after puncture discharge their characteristic contents upon the surface. Pre-sternal dermoids occur, and may be linked dumb-bell fashion with a mediastinal extension, this being sometimes facilitated by sternal defects.

The most serious complication of all, however, is malignant change which Rusby (1944) records as affecting 12.9 per cent of all teratoid tumours, the proportion being lower for the more simple dermoid, higher for the solid teratoma, 30 per cent of such malignant growths arise in the former, 70 per cent in the latter. Once such a change has occurred there is little or no hope of successful removal.

Apart from such pathognomonic symptoms as trichoptysis (the spitting up of hairs), and others resulting from their complications, teratoid growths are diagnosed principally by their radiological appearance. This is of a rounded or lobulated shadow, anterior in the mediastinum, bulging to one or other side of the sternum, and in its earlier phases with a clearly defined edge. Later on successive attacks of inflammation reduce this sharpness and the edge may become almost as ill defined as that of carcinoma though no invasion has occurred. When fistulae are established a fluid level may be seen, and even before this an apparent level called Phemister's line is sometimes distinguished where the fat in such cysts separates at body temperature and floats upon the top of the aqueous content. Teeth may be visible in skiagrams of dermoids, or a regular line of punched-out translucent patches representing tooth buds in the midst of a denser germinal ridge. Calcification commonly occurs in their walls, but, of course, may also be seen in hydatids, goitres, around tuberculous lesions, or in the pericardium itself (Fig. 173). Exploratory puncture should never be resorted to since it may carry infection into the pleural cavity, or even worse, malignant cells. The next step in diagnosis must be exploratory thoracotomy.

Dermoids usually consist of one cyst, but may be polycystic. They contain thick gelatinous mucin and sebaceous matter, and from the inner cyst lining protrude finger-like processes or hard ridges (which may contain cartilage or bone) and from which sprout clumps of lank hair. Teeth are often present, as well as other ectodermal derivatives such as neural or retinal tissue. They are some-

FIG. 173. (a and b) A dermoid cyst with marked calcification in its wall. Note the characteristically anterior position in the mediastinum.





times attached to the mediastinum by a pedicle, but in any case are adherent to the pericardium, the pleura and the chest wall. Accessory dermoids may cluster about the main cyst, having no direct communication with it, or being joined by a narrow stalk.

Teratomas share the general character of dermoids but are mainly solid though often having cystic and haemorrhagic areas. Their components are more complex representing all layers and often including glandular and skeletal elements.

The surgical removal of these tumours may present great difficulty owing to their close and frequently dense and tough adhesion to the anterior surface of the pericardium and the great vessels. Such adhesions may be broken comparatively readily, but usually require tedious and dangerous sharp dissection in a region where serious haemorrhage can easily result. There is seldom any way of telling before operation whether or not malignant invasion has added to these hazards. The aim of surgery, none the less, should invariably be complete excision, and not the marsupialization sometimes practised formerly.

### TUMOURS OF LYMPHOID TISSUE

Lymphatic tissue is richly disposed in the mediastinum. The thoracic duct itself transverses it, lymph from the oesophagus and from the lungs drains into it, and chains of lymphatic glands are found clustered on either side of the trachea, at the carina, about the hila, and in both pulmonary ligaments. It is thus not surprising that these mediastinal glands are involved both in localized tumours and granulomas, and in generalized disorders of lymphatic tissue.

The commonest cause of local enlargement of the mediastinal glands is without doubt tuberculous adenitis, an invariable, but usually transitory, accompaniment of the primary tuberculous lesion. It may however be associated with atelectasis of one or other of the pulmonary lobes from external pressure on a bronchus. In later life also, tuberculous adenitis may be mistaken for a primary mediastinal tumour or cyst, and tuberculous glandular masses, sometimes breaking down and semi-cystic, have been removed. Such glands are adherent to the neighbouring vessels and bronchi, rendering dissection from the former difficult, and sometimes ulcerating into the lumina of the latter, producing strictures or discharging calcified material into them in the form of "lung stones" or broncholiths. The pre-existence of pulmonary tuberculosis, and frequent calcification in the mass, aids diagnosis, but such masses are not necessarily at the hilum, and may simulate cysts or breaking down growths.

Metastatic deposits, especially from bronchial carcinoma, are common, and occasionally are more conspicuous than the parent growth. On the left side they are often associated with recurrent laryngeal nerve palsy and hoarseness, the only innocent lesion commonly causing this being an aortic aneurysm.

Lymphoblastoma and lymphosarcoma occur, the former mainly in children, the latter in young adults, and present smoothly lobulated massive shadows generally responsive to deep x-ray therapy, though lymphoblastomas may not be so, and ultimately fatal. When the presence of either is suspected, a diagnostic dose of x-rays provides the best method of identification since they, with lymphadenoma, are practically the only mediastinal tumours which will so respond.




FIG. 174—A case of Hodgkin's disease treated by deep x-ray therapy

Reticuloses and Hodgkin's disease similarly treated may be part of a generalized disorder or, not infrequently, appear to be localized to the mediastinum spreading to lymphoid tissue elsewhere only later in the course of the disease and running a comparatively chronic course (Fig. 174). Biopsy of accessible lymphatic glands, the blood picture, or a Peltz-Ehrlich syndrome may assist diagnosis.

Lymphangioma or cystic hygroma rarely occurs in the mediastinum and some reported instances are believed to be confusions with cysts of other origins. Arising as a congenital malformation of lymphatic vessels it is unencapsulated, multilocular, intimately incorporated among the structures surrounding it (so that it cannot be shelled out from them) and lined by flattened mesothelium.

### PERICARDIAL CYSTS

These not uncommon thin-walled cysts, lined by a single layer of endothelial cells, and largely filled with clear fluid are almost always situated anteriorly and closely related to the cardio-phrenic angle. 70 per cent of them are found on the right side (Fig. 175). Some have a pedicle attaching them to the pericardium and there are cases on record where fluid in the cyst could be expressed into the pericardium by a narrow communication so that the cyst was in reality a diverticulum. On the other hand many have no such attachment and are either loosely related to pericardial fat in the angle or attached to the antero-medial half of the diaphragm. That they have a developmental origin is not in doubt but differing theories have been advanced to explain their relationship. One suggests that they represent




FIG. 175—(a and b)  
A pericardial coelomic cyst situated typically in the right cardio-phrenic angle. At operation the cyst was found to be thin walled and contained clear fluid. A tenuous pedicle attached it to the pericardium near its junction with the diaphragm.

(a)



(b)

mesenchymal lacunae arising during the formation of the pericardial coelom, another that they are pinched off from the pneumato-enteric recess of the pleuro-peritoneal cavity by the elements of the primitive diaphragm, and constitute the "bursa infracardiaca", a third that they are formed from a constant "ventral parietal recess" of the pericardium during early foetal life.

Be that as it may, the great majority are found in the right cardio-phrenic angle, although they do occur upon the opposite side, and rarely, elsewhere in the mediastinum. As a rule symptomless, they may become sufficiently large to cause dyspnoea, but are usually discovered on routine radiography. Most are flaccid so that their shadow is irregularly convex and not commonly really spheroidal. Easily aspirated anteriorly, the fluid contents is as a rule so clear that they have been called "spring-water" cysts. No example of infection or of malignancy has so far been reported. They are easily removed by thoracotomy.

### THYMOMA

Some 15 per cent of patients with myasthenia gravis have a thymoma and 75 per cent of patients with a thymoma have myasthenia gravis. No constant relation has been detected between the two, nor has a curare like factor been found in the thymomas, and in a quarter of the patients with a thymoma there is no sign of myasthenia. No constant effect has been observed in myasthenics following removal of the tumour, but Keynes believes that their presence worsens the prognosis, results in greater resistance to Prostigmin and that they should be sought for in every myasthenic patient.

Essentially tumours of the anterior mediastinum, they are usually found in its upper part related to the thymus itself, but may occur much lower down on the pericardium. On lateral x ray films they appear sharply outlined and rounded (Fig. 176), though some show as a flattened bulge on the front of the cardiac shadow. Calcification may be seen in them either in a cyst wall or diffusely scattered.

The tumour itself is slowly growing and composed of thymic constituents epithelial cells and thymocytes, palisaded around cystic spaces and blood vessels. Hassall's corpuscles are usually absent. The whole is wrapped in a thick fibrous capsule from which trabeculae traverse the gland and which adheres closely to contiguous structures. In spite of this appearance and whether or not calcification is present, about a quarter of thymomas extend beyond their capsules and become locally invasive rendering removal difficult or impossible. Distant metastases however have not been recorded. In the past they have probably been confused with lymphosarcomas or anaplastic bronchial carcinomas since the small round thymocyte may closely resemble the cells of these tumours.

In general thymomas cause no symptoms unless associated with myasthenia but malignant change may be signalled by pain in the chest, and superior vena caval and innominate obstruction have been described. The onset of substernal pain in a myasthenic is therefore of grave significance and any anterior mediastinal tumour in a myasthenic is virtually certain to be a thymoma.

Surgical access is best gained by splitting the sternum down to the third or fourth interspace. The thymus lies directly beneath between the right and left




FIG 176 —A malignant thymoma in a girl of 19 years without myasthenia. It was however still removable although this involved resection of part of the middle lobe all the anterior pericardium some of the opposite pleura and the thymus gland

pleurae, and can be removed largely by blunt dissection, and ligation of its main venous communication with the left innominate vein. Should surgical removal of an invasive thymoma prove impossible, deep x-radiation is worth trial since some tumours are responsive.

### ENTEROGENOUS CYSTS

Most of the reported cases have been in infants under 1 year of age, suggesting that their presence is not long compatible with life, but a few examples are found in older groups. They have a similar origin to bronchial cysts, arising as diverticulae from the developing foregut during foetal life, and are characterized by the possession of a mucosal lining resembling some portion of the gut, but especially and principally the stomach. This may be complete, and secrete gastric juice copiously with all its components, so that the cyst resembles an accessory but isolated stomach, or represented by such small fragments of gastric mucosa that its identity may pass unrecognized and the cyst contain little but mucus. As is the case with most congenital anomalies, other malformations may co-exist, such as cysts and diverticulae elsewhere, or spina bifida.

Generally the cyst is closely related to the oesophagus and so is found posteriorly, and in nine cases out of ten upon the right side — possibly because of the presence of the aorta on the left. It may be densely adherent to its parietes, and the wall is sometimes as thick as that of the stomach and includes submucosal and muscular

## INTRATHORACIC GOITRE

layers. Such cysts are found only in infants since the gastric juices they secrete cause them rapidly to distend and either rupture into the oesophagus, lung, or pleura, or cause death by interfering with respiratory function. Those seen in later life, while also occupying a posterior position will contain little secreting mucosa and have a thinner wall. Seven examples of gastric ulceration within enterogenous cysts are recorded, and 4 have penetrated into bronchi.

In infancy cough, dyspnoea, and cyanosis occur, due to the cyst's bulk or collapse of the adjacent lung and haemoptysis haematemesis, and melaena are all described depending upon the course taken by the ulceration. Once fistula is established a fluid level will be seen across the x-ray shadow, revealing its cystic character. Should aspiration have been carried out under the impression that an effusion was being dealt with the presence of hydrochloric acid or pepsin in the fluid withdrawn make the diagnosis clear though this measure is most unwise. Cysts containing little or no gastric mucosa remain symptomless.

Surgical removal is life saving in infancy, and advisable at any age.

## INTRATHORACIC GOITRE

Colloid goitres frequently become substernal occupying a position in the anterior part of the superior mediastinum and may here be mistaken for any of the tumours or cysts intrinsic to the region in particular those arising from the thymus. They may attain great size but retain from the nature of their growth their vascular attachments in the neck. In time most undergo toxic degeneration with resultant tachycardia or auricular fibrillation. From their situation in the rigid and constricted thoracic outlet pressure signs are often in evidence with dilatation of neck veins and compression of the trachea which is displaced to one or other side according to the thyroid lobe principally enlarged or flattened from in front. Symptoms as well as evidence of toxicity may include an irritative cough, dyspnoea or dysphagia. Should haemorrhage occur into one of the cystic spaces of the gland acute exacerbation of symptoms is apt to result.

Clinically such patients may have evidence of thyroid enlargement in the neck, or the scar of previous thyroidectomy but sometimes the enlargement is wholly substernal. It is usually possible however to palpate the upper lobe of the tumour in the suprasternal notch or supraclavicular fossa especially on deep inspiration and it characteristically rises on deglutition distinguishing itself thus from other intrathoracic growths.

In antero-posterior skiagrams of the chest an irregularly rounded shadow is seen in the superior mediastinum generally more prominent upon one side and displacing the trachea to the other. Its convexity is downwards and its upper limits cannot be defined shading off into the base of the neck. Lateral views show it to lie anteriorly and often reveal calcification which may be extensive (Fig. 173). The use of radioactive iodine isotopes is now increasingly used to identify intrathoracic thyroid masses.

The surgical approach should always be by means of an orthostotic thyroidectomy collar incision in the neck since it is here the vascular pedicle is to be secured. If necessary the infrahyoid muscles on one side are divided and dissection carried down to the capsule of the gland. The thyroid vessels are secured and the mass




FIG 177 —Petrosternal thyroid (a) showing the tumour chiefly to the right of the mediastinum, (b) lateral film shows its anterior position and reveals considerable calcification in its substance

(a)



(b)

delivered into the neck by a combination of gentle traction and leverage from below, on occasion with a spoon. Some surgeons recommend evacuating the colloid material from the interior of larger glands but all are agreed that sternum splitting is seldom if ever required. Subtotal thyroidectomy is then performed care being taken not to damage the recurrent laryngeal nerve as it courses on the postero-lateral surface of the glands. True ectopic intrathoracic thyroids (having no vascular connexions with neck) have been recorded but are exceedingly rare. They are diagnosed and can be removed, only by exploratory thoracotomy.

### RARE TUMOURS

Such then are the commoner tumours encountered in the mediastinum a variety of rarer growths remain. Among them are found lipomas 40 of which have been recorded. All are extrapleural and may attain great size being signalled by a progressive and ultimately severe dyspnoea. Most remain wholly within the thorax but 9 have presented dumb-bell fashion beside the sternum and 7 in the root of the neck, where they are felt to be soft lobulated and to transmit an impulse on coughing. X ray appearances are those of an opacity of which the edges may be perceptibly more translucent than the centre but which otherwise resembles an encysted or interlobar effusion.

Some dozen examples of meningocele protruding into the chest are described most of them at first mistaken for neurofibromas because of their smooth outline and posterior situation. They may cause pressure symptoms and bone erosion of intervertebral foramina and might be diagnosed (were their nature suspected) by myelography. The writer has also seen a meningioma presenting in the upper part of the chest.

Haemangioma of the mediastinum may be single and encapsulated in multiple foci as part of generalized congenital telangiectasia or frankly malignant. At least 17 instances occur in the literature of which 8 were malignant and 2 were part of many primary lesions. Appearing anywhere in the mediastinum they are not to be identified before operation though a vascular bruit may be audible over them. This feature however they share with any very vascular tumour with aneurysms, and with arteriovenous communications.

Other, and more exceptional tumours numbered only in ones or twos but appearing from time to time still further to complicate the already difficult differential diagnosis of mediastinal neoplasms are leiomyoma an innocent growth associated with the smooth muscle of the oesophagus parathyroid adenoma with the skeletal changes raised serum calcium and generalized osteitis fibrosa cystica single or multiple innocent or malignant fibroma chondroma myxoma, and sarthoma.

### BIBLIOGRAPHY

- Blades, B. (1946). "Mediastinal Tumours." *Ann. Surg.* 123, 42.  
 Blakes, A. (1944). "Thymectomy in the Treatment of Myasthenia Gravis." *J. thorac. Surg.* 13, 116.  
 Blum, F. N., Allig, E. E. and Samson, P. C. (1949). "Intra-Serous Meningocele." *J. thorac. Surg.* 18, 294.  
 Clagett, O. T. and Ross, G. T. (1944). "Surgical Approach for Tumours of the Thorax." *Surg. Gynec. Obstet.* 78, 12.






FIG 177—Petrosternal thyroid, (a) showing the tumour chiefly to the right of the mediastinum, (b) lateral film shows its anterior position and reveals considerable calcification in its substance

(a)



(b)

## RARER TUMOURS

delivered into the neck by a combination of gentle traction and leverage from below, on occasion with a spoon. Some surgeons recommend evacuating the colloid material from the interior of larger glands, but all are agreed that sternum splitting is seldom if ever required. Subtotal thyroidectomy is then performed, care being taken not to damage the recurrent laryngeal nerve as it courses on the postero-lateral surface of the glands. True ectopic intrathoracic thyroids (having no vascular connexions with neck) have been recorded, but are exceedingly rare. They are diagnosed, and can be removed, only by exploratory thoracotomy.

## RARER TUMOURS

Such then are the commoner tumours encountered in the mediastinum, a variety of rarer growths remain. Among them are found lipomas, 40 of which have been recorded. All are extrapleural and may attain great size, being signalled by a progressive and ultimately severe dyspnoea. Most remain wholly within the thorax, but 9 have presented dumb bell fashion beside the sternum, and 7 in the root of the neck, where they are felt to be soft, lobulated, and to transmit an impulse on coughing. X-ray appearances are those of an opacity of which the edges may be perceptibly more translucent than the centre, but which otherwise resembles an encysted or interlobar effusion.

Some dozen examples of meningocele protruding into the chest are described, most of them at first mistaken for neurofibromas because of their smooth outline and posterior situation. They may cause pressure symptoms and bone erosion of intervertebral foramina, and might be diagnosed (were their nature suspected) by myelography. The writer has also seen a meningioma presenting in the upper part of the chest.

Haemangioma of the mediastinum may be single and encapsulated, in multiple foci as part of generalized congenital telangiectasia or frankly malignant. At least 17 instances occur in the literature, of which 8 were malignant, and 2 were part of many primary lesions. Appearing anywhere in the mediastinum they are not to be identified before operation, though a vascular bruit may be audible over them. This feature however they share with any very vascular tumour, with aneurysms, and with arteriovenous communications.

Other, and more exceptional tumours, numbered only in ones or twos, but appearing from time to time still further to complicate the already difficult differential diagnosis of mediastinal neoplasms are leiomyoma, an innocent growth associated with the smooth muscle of the oesophagus, parathyroid adenoma, plasmocytomas, single or multiple, innocent or malignant, fibroma, chondroma, myxoma, and xanthoma.

## BIBLIOGRAPHY

- Blades, B. (1946) Mediastinal Tumours. *Ann. Surg.* **123**, 741.  
 Bialock, A. (1944) Thymectomy in the Treatment of Myasthenia Gravis. *J. thorac. Surg.* **13**, 316.  
 Byron F. A. Alling, I. F. and Simon P. C. (1949) Intrathoracic Meningocele. *J. thorac. Surg.* **18**, 294.  
 Clagett O. T. and Root G. T. (1944) Surgical Approach for Tumours of the Thymus - *Surg. Gynec. Obstet.* **78**, 397.

## DIASTINUM

- is " *J thorac Surg*, 20, 195  
a " *Thorax*, 2, 163  
oc, 5, 1  
rnat Abstr Surg, 84, 195  
University of Minnesota  
inn Surg, 123, 377  
*J thorac Surg*, 13, 16  
of Intestinal and Bronchial

## CHAPTER 30

### DISEASES OF THE DIAPHRAGM

KENNETH M A PERRY

UPWARD displacement of the diaphragm may result from the pressure from below of ascites, pregnancy and abdominal tumour or from pull from above by pulmonary fibrosis or collapse. In all these cases, however, it moves normally. When paralysis is present, there will always be paradoxical movement visible on screening.

#### DIAPHRAGMATIC PARALYSIS

The phrenic nerve is sometimes divided as a therapeutic measure. If it is crushed paralysis will last for 6 months, if it is avulsed the paralysis will be permanent. Trophic changes in the diaphragm follow as early as 3 days after paralysis and atrophy is well established after a month. The muscle is reduced in thickness, fat is deposited and fibrous tissue develops. The phrenic nerve may be divided by trauma or disease processes involving the peripheral course of the nerve, spinal cord or medulla. Thus diaphragmatic paresis may occur in acute anterior poliomyelitis, tumour, progressive muscular atrophy, myelitis, haemorrhage and injuries and carries of the cervical vertebrae. Peripheral lesions of the phrenic nerve include neck wounds, destruction by neoplastic masses, diphtheritic and other types of peripheral neuritis and avulsion from birth injury as an extension of Erb's paralysis. Blattner (1942) and Light (1944) have recorded cases in newborn infants resulting from intra-abdominal malposition. Mediastinal tumours, and chronic inflammatory processes in the mediastinal and paratracheal lymph nodes not infrequently involve the phrenic nerve (Fig 178). Friedman (1950) recorded 6 cases of phrenic paralysis in association with pneumonia. Phrenic paralysis is occasionally found in patients when no explanation can be given for its development but it is extremely doubtful whether stretching of the nerve by aortic aneurysm ever causes paralysis though Sangunette and Galzerano (1943) have suggested that this may happen.

#### EVENTRATION OF THE DIAPHRAGM

Eventration of diaphragm is the name applied to a condition in which the diaphragm is abnormally high, as contrasted with an upward displacement from pathological causes (Fig 179). It is a rare condition and is commoner on the left side than the right (Korns, 1921, Reed and Borden, 1935, Evans and Simpson, 1950). The condition is probably of congenital origin but may result from birth injury to the phrenic nerve. The diaphragmatic muscle is usually atrophied and replaced by a related translucent membrane of white fibrous tissue with few muscle fibres. It is frequently paralysed and shows paradoxical movement, but this is not always so. The condition does not as a rule cause symptoms and in the rare cases where they do occur they are usually of a dyspeptic nature. Shortness of breath on

exertion and cough and pain in the left side may occur. Physical examination will reveal poor movement of the left side of the chest and its paradoxical nature may be evident. Fluoroscopy will confirm these findings while the use of a barium meal will differentiate the condition from a short oesophagus with herniation. The stomach often appears as a characteristic inverted fish hook, being rotated so that the greater curvature is uppermost under the dome of the diaphragm, the pylorus and pyloric antrum lie anteriorly in the left upper quadrant of the abdomen (Rosenfeld, 1944). A pneumoperitoneum will assist in outlining the diaphragm.

Treatment is not satisfactory. Dyspepsia can be treated with a light diet, phenobarbitone, and alkaline and antispasmodic mixtures. Surgery has been used in the treatment of the condition, the diaphragm being plicated or incised, overlapped and sutured. At the operation the appearance and calibre of the phrenic nerve is always normal.

### DIAPHRAGMATIC TRICHINIASIS

Trichiniasis is caused by eating pork or bear meat which contains living larvae of the nematode *Trichinella spiralis*. These larvae are liberated in the gastrointestinal tract during the digestion of the meat. They quickly mature and then mate, when the female produces new larvae which penetrate the wall of the gut to enter the blood stream. Some are deposited in the muscles where they remain alive for years and become encysted; others lodge in other organs where they die. The human illness occurs only during the first few weeks of infestation.

Symptoms rarely arise during the stage of intestinal invasion and there are no symptoms after the larvae become encysted. These usually arise 7–12 days after ingestion and consist of fever, facial oedema, sometimes oedema elsewhere, skin rashes and muscular pains. Chest pain is common and is attributed to infestation of the intercostal muscles and the diaphragm. There is sometimes hiccough; cough is also common and is unproductive, probably resulting from the presence of parasites in the lungs. haemoptysis occurs occasionally.

Eosinophilia is the best guide to early diagnosis. The parasites can sometimes


*Trichinella* antigen are useful diagnostic procedures.

The condition quickly resolves and little is required in the way of treatment. The disease can readily be prevented by ensuring that pork is well cooked. An outbreak occurred in Wolverhampton in 1941 as a result of people eating uncooked sausage meat (Sheldon, 1941).

### DIAPHRAGMATIC HERNIA

There are two types of diaphragmatic hernia—those due to trauma and those resulting from congenital weakness. Traumatic herniae may either be due to penetrating or non penetrating injuries or be a post operative occurrence (Fig 180). Where they result from non penetrating injuries, the injury is usually of a crushing nature but it may be by less violent trauma such as falling down stairs. In the case of penetrating or post operative injuries the cause is always

FIG. 178. — Diaphragmatic paralysis resulting from involvement of phrenic nerve in a tuberculous adenitis (a patient of Dr. Sita Lun-pien)



evident. When the hernia is on the right side the liver and colon are the herniated viscera while on the left side the stomach and colon are herniated and perhaps if the rupture is large the ileum and spleen as well (Figs 181-182). Non traumatic diaphragmatic herniae may either occur through the oesophageal hiatus or through defects in the muscular and tendinous portions of the diaphragm. The latter are more common on the left side than on the right. They can occur through the foramen of Bochdalek the foramen of Morgagni through a congenital absence of the posterior fourth or through defects in various parts of the dome. On the left side they are most commonly posterior on the right side where the herniae are often parasternal they are usually anterior and contain colon or liver. The oesophageal hiatus herniae may be associated with a short oesophagus oesophagitis and peptic ulceration or with a normal oesophagus when they may be of the para-oesophageal or sliding gastric oesophageal type. They develop in adult life as a result of obesity and laxity of the tissues at the hiatus from advancing age or increased intra abdominal pressure. Oesophagoscopy will show if oesophagitis or peptic ulceration is present.

In infancy shortness of breath cyanosis colicky pain vomiting and abdominal distension are the principal symptoms. These are often improved by holding the child in the upright position thus enabling the hernia to reduce itself. In later life the symptoms are usually dyspnoea or retro-sternal pain resembling that caused by coronary ischaemia. Although is an occasional symptom. Screening the patient in the recumbent position after a barium meal is the most important aid to diagnosis. In right sided cases a barium enema is most helpful.





FIG 179 — Typical example of eventration of the left side of the diaphragm (a) Postero anterior view, (b) lateral view. An example of the rare right sided eventration is seen in (c) and (d) (c) Postero-anterior and (d) lateral

(a)



(b)

# DIAPHRAGMATIC HERNIA



(c)



(d)



Skiagrams usually show the abnormality, which has to be differentiated from cysts and tumours of surrounding structures or from pleural effusion.

If a hernia is liable to strangulate, surgical repair is indicated. Treatment of hiatal herniae should be medical and the peptic ulcer régime used if oesophagitis is present. The patient should sleep propped up. Left phrenic paralysis gives good results in older patients but repair of the defect by the transthoracic approach is usually the best treatment. If stricture or severe ulceration is present, dilatation of the oesophagus with application of silver nitrate or resection of the affected portion may be necessary.

### SUB-DIAPHRAGMATIC ABSCESS

This is not truly a disease of the chest but it so often causes sympathetic lesions above the diaphragm that a book of diseases of the chest is not complete without some consideration of the condition. There are five spaces in which these abscesses may form, right and left supra-hepatic are the commonest sites, the others being right infra-hepatic and left anterior and posterior infra-hepatic, the frequency being in that order. The supra-hepatic lies between the diaphragm and the dome of the liver while the infra-hepatic space is the under surface of the liver and the transverse colon. The supra-hepatic space is divided into right and left portions by the falciform ligament and the infra-hepatic is similarly divided by the ligamentum teres and ligamentum venosum.

Most cases of subdiaphragmatic abscess arise from pocketing of pus following a general peritonitis resulting from appendicitis, perforation of peptic ulcers, or malignant neoplasms of stomach or colon, cholecystitis and liver abscess. In the last mentioned case there is a high mortality rate. The majority of these occur post-operatively. Some may result from direct spread of the pyogenic infection, but there is little evidence to confirm that they can result from lymphatic spread as was once thought. Occasionally they occur as part of a pyaemic process and rarely from a spread from inside the chest. They may occur at any age but are

effusions are usually sterile, but empyema is a frequent complication. In a quarter of the cases (Harley, 1949) and suppurative pneumonitis also results in a few. Perforation of the diaphragm either from spread of the infection or as a result of transpleural aspiration or damage is the most common cause of the suppurating conditions.

### Clinical features

The clinical manifestations of this disease tend to appear 3-4 weeks after the causal laparotomy. Faxon (1940) in reviewing 111 patients from the Massachusetts General Hospital found an average interval of 26 days. The usual presenting symptoms are fever and pain along one of the costal margins. The onset is sometimes abrupt with rigors, sweating, vomiting, abdominal pain and collapse. More often it is subacute and in a few it is slow and insidious and in this last group it

FIG 180—Traumatic hernia of liver and colon through right side of diaphragm. (b) L. podol. n. bronchus and barium in colon (a patient of Dr. S. A. L. pl. n.)



(a)



(b)




FIG 181 —Diaphragmatic hernia of colon through an anterior parasternal defect in the right side of diaphragm (b) lateral view showing barium in colon

(a)



(b)




FIG 182 —Diaphragmatic hernia of stomach and intestines through the left diaphragm

may go unrecognized for many months. Other symptoms which may occur include cough, shortness of breath, pain in the lower part of the chest or generalized toxæmia.

The physical signs are scanty unless a pleural effusion is present, when there will be dullness and absent breath sounds over the lower part of the chest. Otherwise localized tenderness over the site of the abscess will be the only sign that can be elicited. A blood count will always reveal a polymorphonuclear leucocytosis. Diagnostic aspiration is inadvisable owing to its unreliability and because of the danger of infecting the pleura. The most significant radiological finding is fixation and elevation of the diaphragm, and on the right side if the diaphragm is normal in position and moves freely on respiration a diagnosis of subdiaphragmatic abscess can be excluded. This rule however is not absolute on the left side. When air is present under the diaphragm it is a valuable aid to diagnosis but this occurs in only about a quarter of the cases. On the left side it must be distinguished from gas in the stomach. In about 25 per cent of cases a pleural effusion is present and this will be evident in the skiagrams as well as pulmonary consolidation if it is present. It is important to take lateral pictures as well as antero-posterior ones while postero-anterior ones in lateral recumbency are sometimes helpful.

#### Treatment

Chemotherapy with adequate dosage of sulphonamides, penicillin and streptomycin will prevent the development of a large number of sub-diaphragmatic

abscesses, and make the condition a rare one. Chloromycetin and terramycin will also play a part in achieving this.

Once there is an established abscess surgical drainage becomes essential and the mortality rate is greatly reduced if an extra serous method of drainage is adopted. An incision can be made over the twelfth rib which is resected subperiosteally. The abscess can then be drained by a transverse incision through the bed of the rib at the level of the spinal process of the first lumbar vertebra. Ochsner and DeBakey (1938) give a mortality rate of 20.8 per cent for retroperitoneal drainage compared with 36.2 per cent for transpleural and 35.1 per cent for transperitoneal and Harley (1949) gives 11 per cent for extra serous drainage and 33 per cent for trans serous.

Serous pleural effusion will usually absorb spontaneously when the abscess has been drained but intrathoracic suppuration usually makes the condition more serious. When the abscess is drained, however, by the establishment of a bronchial fistula the mortality is significantly diminished.

### DIAPHRAGMATIC TUMOURS

Primary cysts and tumours are rare. They were first described by Grancher (1868),

the condition was carefully reviewed by Cruickshank and Cruickshank (1951). These congenital cysts appear to be of mesothelial origin (Fig 183). They arise from that part of the coelom which becomes the pleuro pericardial canal and are closely related to pleuro pericardial cysts. Many are situated within the substance of the diaphragmatic muscle. Histologically they have a fibrous wall and are lined by a single layer of flattened cells.

Tumours of various kinds occur in the diaphragm, sarcomas, which have been described as rhabdomyosarcoma, fibromyosarcoma, mixed cell sarcoma, myoblastic sarcoma, leiomyosarcoma, myosarcoma and endothelial sarcoma, are the commonest. Lipomas are the next most frequent tumours, but fibromas, mesotheliomas, neurofibromas, fibromyomas, chondromas, angiofibromas and haemangio endotheliomas have been described. Endometriomas may also occur in the diaphragm.

Severe pain in the lower part of the chest aggravated by deep breathing is the commonest symptom. Cough with a little mucoid sputum occasionally streaked with blood has been reported. Arkless (1942) described the coincidence of a rhabdomyosarcoma of the diaphragm, idiopathic hypoglycaemia and retroperitoneal sarcoma. On examination a tumour is sometimes palpable between the lower ribs or in the upper abdomen. It moves with respiration. It has on occasion been mistaken for the liver or spleen. Skiagrams will show a tumour in the lower part of the chest which is difficult to separate from surrounding structures. It moves synchronously with the diaphragm. Diagnostic pneumothorax or pneumoperitoneum may help to define its situation. Thoracoscopy may also help in the diagnosis.

Prognosis depends on the nature of the tumour, in the malignant conditions it is always bad.

# DIAPHRAGMATIC TUMOURS

FIG 183 — Mesothelial cyst of the diaphragm lying in the left costo diaphragmatic angle (b) lateral view of same (a patient of Mr G Flavell)



(a)



The cysts and tumours should always be removed surgically, since more than half are malignant and it is only after removal that the diagnosis can be finally established

## BIBLIOGRAPHY

- Ackerman, A J (1942) "Primary Tumours of the Diaphragm Roentgenologically Considered" *Amer J Roentgenol*, 47, 711
- Allison, P R (1946) "Peptic Ulcer of the Oesophagus" *J Thorac Surg*, 15, 308
- Arkless, H A (1942) "Coincidence of Rhabdomyosarcoma of the Diaphragm, Idiopathic Hypoglycaemia and Retroperitoneal Sarcoma" *Med Bull Veterans' Adm, Wash*, 19, 225
- Barnard, H. L. (1908) "Surgical Aspects of Subphrenic Abscess" *Brit Med J*, 1, 371
- Barrett, N R., and Wheaton G E W (1934) "The Pathology, Diagnosis and Treatment of Congenital Diaphragmatic Hernia in Infants" *Brit J Surg*, 21, 420
- Beeson, P B (1941) "Trichiniasis, Clinical Manifestations and Diagnosis" *Lancet*, 2, 67
- Binney, H (1931) "Tumours of the Diaphragm" *Ann Surg*, 94, 524
- Blattner, R J (1942) "Unilateral paralysis of the Diaphragm without Involvement of the Brachial Plexus" *J Pediat*, 20, 225
- Clagett, O T., and Johnson, M A (1949) "Tumours of the Diaphragm" *Amer J Surg*, 78, 526
- Cruikshank, G., and Cruikshank, D B (1951) "Intradiaphragmatic Mesothelial Cysts" *Thorax*, 6, 145
- Delle Vide, B L., and Dyke, S C (1941) "Blood Picture in Trichiniasis" *Lancet*, 2, 69
- Donovan, E. J (1945) "Congenital Diaphragmatic Hernia" *Arch Surg*, 122, 569
- Dunhill, T (1935) "Diaphragmatic Hernia" *Brit J Surg*, 22, 475
- Evans, C J., and Simpson, J A (1950) "Fifty-seven Cases of Diaphragmatic Hernia and Eventration" *Thorax*, 5, 343
- Faxon, H H (1941) "Logical Approach to Subphrenic Abscess" *Amer J Surg*, 54, 114
- Freedman, B (1950) "Unilateral Paralysis of Diaphragm and Larynx Associated with Inflammatory Lung Disease" *Thorax*, 5, 169
- Gale, J W., and Edwards, S R (1939) "Malignant Tumours of the Diaphragm" *J thorac Surg*, 9, 185
- Graucher (1868) "Tumeur végétante du centre phrénique du diaphragme" *Bull Soc Anat, Paris*, 43, 1874
- Gross, R. E (1946) "Congenital Hernia of the Diaphragm" *Amer J Dis Child*, 71, 579
- (1940) "Subphrenic Abscess A report of 111 consecutive operative cases" *New Engl J Med*, 222, 289
- Harley, H R S (1949) "Subphrenic Abscess" *Thorax*, 4, 1
- Hochberg, L. A (1942) "Diagnostic Criteria for Subphrenic Abscess Based on a study of 139 cases" *Ann intern Med*, 17, 183
- Harrington, S W (1942) "Diaphragmatic Hernia of Children" *Arch Surg*, 115, 705
- (1940) "Diagnosis and Treatment of Various Types of Diaphragmatic Hernia" *Amer J Surg*, 50, 377
- Husteldt, E., Thomsen, G., and Wamberg, E (1951) "Hiatal Hernia and Short Oesophagus in Children" *Thorax*, 6, 56
- Korns, A M (1921) "The Diagnosis of Eventration of the Diaphragm" *Arch intern Med*, 28, 192
- Light, J S (1944) "Respiratory Shaft in Epigastric abdominal wall, a Physical Sign seen with complete Unilateral Paralysis of the Diaphragm in Infants and Children" *J Pediat*, 24, 627
- Neuhof, H., and Schlossmann, N C. (1942) "Left Subphrenic Abscess" *Surg Gynec and Obstet*, 75, 751
- Ochsner, A., and DeBakey, M (1938) "Subphrenic Abscess Collective review and an analysis of 3,608 collected and personal cases" *Internat Abstr Surg*, 66, 426
- Pancoff, H K (1926) "The Roentgenological Diagnosis of Liver Abscess with or without Subdiaphragmatic Abscess" *Amer J Roentgenol*, 16, 303
- Reed, J A., and Burden, D L (1935) "Eventration of Diaphragm and Report of Two Cases" *Arch Surg*, 31, 30

# BIBLIOGRAPHY

- Robson K and Collis J L (1944) Tumours of the Diaphragm with Report of a Diaphragmatic Cyst *Brit J Tuberc* 38, 3
- Rosenfeld D H (1944) Unusual Type of Inversion of Stomach Associated with Diaphragmatic Eventration and other Anomalies *Amer J Roentgenol* 52, 607
- Sanguinetti A A Galzerano D A (1943) Síndrome diafragmático por aneurisma de la aorta torácica *Rev Asoc med argent* 57, 413



## CHAPTER 31

### ANAESTHESIA IN CHEST DISEASE

A I PARRY BROWN

THE PATIENT with chest disease who needs to be anaesthetized presents two problems to the anaesthetist. The anaesthetic may aggravate the chest disease, or the diseased and inefficient respiratory apparatus may make the administration of the anaesthetic more difficult and dangerous. *The first important step is to establish the extent of the respiratory disease.* The site of the lesion, the likelihood of sputum, and the response to treatment can be ascertained on reference to the medical history. The ease with which the respiratory exchange is maintained and the reserve for exercise are points of value in planning the method of anaesthesia.

### SURGICAL EMERGENCIES

First there is the surgical emergency with acute respiratory disease. There is no

to effect a transfer. The vital capacity may be reduced by congestion or effusion, or even by previous treatment, as by the use of an artificial pneumothorax. Additional barriers may be interposed between the ventilating gases and the circulating blood as, for example, widespread mucus in a tracheo-bronchitis. Air may be drawn into areas of lung without a corresponding blood supply as in the case of a giant air cyst or in chronic emphysema. These patients become sub-oxygenated on mixtures which would be quite adequate for the normal patient. The anaesthetic is less quickly absorbed and excreted so that both the induction and the recovery period are prolonged.

*The patient who knows that he has a chest disease fears that the anaesthetic will affect him adversely.* This is particularly true in patients with pulmonary tuberculosis. Such patients when faced with an operation for acute appendicitis are genuinely, perhaps with reason, more frightened of the anaesthetic than of the operation. This state of nervous tension makes the induction more difficult and must be allayed by reassurance, which can only come from a careful pre-operative assessment.

In this group of surgical emergencies time is not available for any thorough

limited in serious chest disease. If the patient is dyspnoeic or orthopnoeic, the position on the table will be intolerable. Many of the techniques of local analgesia used for abdominal operations involve the paralysis of the intercostal muscles,

which would increase the respiratory embarrassment. Cases with uncontrollable coughing are difficult to manage at the time of operation if spinal analgesia is chosen. They are subjected to a greater than normal risk of post-spinal headache, for the persistent coughing encourages the leak of cerebrospinal fluid through the dural puncture. If some of these points contra-indicate the use of local analgesia then cyclopropane is probably the best of the general anaesthetic agents being non-irritant to the respiratory tract and the administration is with high oxygen concentrations. The respiration is quiet under its action, a rapid return of the protective reflexes after the operation follows as cyclopropane is quickly eliminated.

A woman was admitted to hospital suffering from lobar pneumonia, who had been in labour for 24 hours and had reached the stage of exhaustion. She was cyanosed and orthopnoeic with a respiratory rate of 40 per minute. It was obvious that she was unable to deliver herself and as the cervix was nearly fully dilated it was decided to deliver her with high forceps. After premedication with atropine, anaesthesia was induced with a mixture of cyclopropane 20 per cent and oxygen 80 per cent. On this mixture her respiratory distress was relieved, and her colour improved. As she lost consciousness it was possible to lower her into the horizontal position without a return of the cyanosis. High forceps were applied in the lithotomy position and the woman was safely delivered. The return of consciousness was rapid and as the splinting of the diaphragm by the uterus was relieved she was able to breathe easily and made a rapid recovery under medical treatment.

Whilst such a case suited the technique almost as well as the technique suited the case it must be remembered that these cases can be difficult and it is unwise to use an unfamiliar technique. It is safer to use a routine method, provided that it is physiologically sound, than to use an agent which however ideal suffers the disadvantage of unfamiliarity. Nitrous oxide, oxygen and ether provides a sequence which although humdrum is safe during the administration and which carries little extra risk of post-operative complications if steps are taken to prevent the spread of secretions whilst the reflexes are inactive and ciliary movement is reduced.

An even more dangerous error than using an appropriate anaesthetic agent but with insufficient experience is to strain a weak agent beyond its proper capabilities. Nitrous oxide has great advantages as an anaesthetic agent for it is non-toxic, and rapidly eliminated, but it is weak and can only produce more than the light planes of anaesthesia by depriving the patient of oxygen. The old finding of "fit for gas" has no place in anaesthesia for cases with chest disease.

### SPREAD OF SECRETION

Unsound technique greatly increases the danger of the spread of secretions. Forced hyperpnoea by the use of carbon dioxide may simplify and accelerate the induction of anaesthesia in the healthy patient, but it is obviously inapplicable in these cases. The cilia of the respiratory mucosa are easily damaged. Rough instrumentation may destroy them and leave the patient partially deprived of this mechanism for the removal of secretions. The cuffed endotracheal tube particularly when lubricated by a greasy ointment containing a powerful surface anaesthetic is injurious and should only be used when essential. The spraying of the larynx

and bronchial tree with local analgesia solutions carries with it the danger that secretions may spread whilst reflexes are depressed, and if it is necessary to use them steps must be taken to control secretions. In all except acute emergencies this control can start pre-operatively. Postural drainage and suitable medical treatment can reduce the amount of secretion so that the risk of flooding new areas of lung is greatly reduced. Premedication with atropine or scopolamine by keeping the respiratory tract clear from mucus reduces the volume of any secretions and so the likelihood of soiling, but at the price of making viscid plugs which are harder to cough up.

### Control of spread

The secretions may be controlled during the operation by suction by posture, or by the isolation of the diseased area of lung. The removal of secretions by suction is the most immediately attractive solution, for if the anaesthetic gases are being given through an endotracheal tube the only change from normal routine is the occasional passage of a suction catheter down the endotracheal tube. The disadvantages are that it interferes with the smooth administration for, the anaesthetic gases are lost during the periods of aspiration. It requires deeper anaesthesia to prevent coughing when the end of the suction catheter touches the carina than is needed for many operations.

It may fail because of the silent spread of small quantities of sputum or because large volumes need such constant attention that no time is left to give the anaesthetic. This blind suction may make worse the condition in the lung. If the catheter is passed into small bronchi all the air may be drawn from a segment which will fail to re-expand. Haemorrhage may be caused particularly if the disease is an adenoma or carcinoma of the bronchus. Despite these disadvantages intrabronchial suction remains a most valuable method of controlling secretions and of preventing the contamination of fresh areas of lung. It should be applied at intervals determined by the amount of secretion and at stages known to be associated with the release of sputum as for example when the patient is moved on to the operating table. The relative lengths of the endotracheal tube and of the suction catheter should be known so that the position of the sucker may be estimated.

A suitable position on the table may eliminate the danger of inhalation of sputum. The secretions may be retained in the diseased area where they cause no reflex disturbance or interference with respiration, or a position can be chosen in which any sputum that is released will run up towards the mouth rather than be absorbed in new areas of lung.

The former is useful in short procedures

with the child half sitting up and partly on his left side. In this position the left lower lobe is dependent and secretions were retained in the diseased area. Nitrous oxide, oxygen and ether were used and a smooth induction was obtained without coughing. When the child was fully anaesthetized a naso tracheal tube was introduced and through

this a fine gum-elastic suction catheter was passed. Nearly an ounce of sputum was aspirated, although it had caused no disturbance whilst retained in the diseased area of lung. The suction catheter was removed and the operation proceeded normally.

Protection is possible by the postural method if the operation is performed in the prone position or in the lateral position if a Trendelenburg tilt is used, but in the supine position, unless an extreme head down tilt is used, the secretions will not gravitate towards the mouth.

The control of secretions by endobronchial tamponade is not usually applicable to cases undergoing general surgery but rather to cases for lung surgery where the diseased area will be removed, for the mucosa at the site of the obstruction will be injured and further damage may be done in the isolated area of the lung.

This danger from the inhalation of secretions is not limited to the duration of the operation but continues throughout the subsequent period of depressed reflexes. The patient may not have such constant supervision during the recovery period so that a technique which allows a quick return to full reflex activity will be advantageous. The worst anaesthetics in this respect are ether, when the anaesthesia is deep, and the short-acting intravenous barbiturates, when repeated administration is needed to maintain anaesthesia for long periods. The advisability of increasing the rate of excretion of ether by the inhalation of carbon dioxide and oxygen at the end of the operation should be considered but there is a danger that this forced over breathing may spread secretions into small bronchi which they will occlude. This method can give a false sense of security after a long deep ether anaesthesia, for during the period of over-ventilation the concentration of ether in the blood is lowered and reflex activity returns. There is afterwards a reaction when the breathing is very shallow, during this period ether is mobilized from body tissues and the blood concentration rises again.

### DANGER OF COLLAPSE OF LUNGS

Diminished movement of the chest wall may lead to collapse of the underlying lung. This restricted movement may be due to pain as after any upper abdominal operation, or due to paralysis of some of the respiratory musculature. Reference has already been made to this undesirable side-action of spinal analgesia. The muscle relaxants which spare the diaphragm but paralyse the intercostal musculature produce paradoxical movement of the apices of the lung, which is inefficient and likely to cause sub-oxygenation in patients with respiratory disease. The artificial respiration which will correct this is likely to spread secretions unless adequate steps are taken to control them.

### AGGRAVATION OF EXISTING DISEASE

Besides these local effects an anaesthetic can aggravate pre-existing diseases by its general toxic action. In pulmonary tuberculosis a critical balance may be maintained between the invasive powers of the organism and the resistance of the host. If this resistance is lowered, as it may well be by an operation, an exacerbation of the disease is to be expected. The lowered resistance is in part due to surgical trauma, in part to blood loss and in part to the anaesthetic. The anes-

thetist can help to reduce the trauma by providing complete muscular relaxation, and some general effects of the trauma can be reduced by blocking painful stimuli with local analgesia. The careful replacement of blood loss is especially worth while in these cases. An estimate of the blood loss as by weighing the used swabs helps in assessing the volume of the transfusion required for the risk of causing pulmonary oedema by over transfusion is greater with pre-existing respiratory disease. Those drugs which interfere with the metabolic processes of the body are most likely to lower the resistance of the patient. On this count the prolonged deep administration of ether or chloroform should be avoided, but it is only fair to add that some authorities use ether with excellent results for all thoracic operations including those for tuberculosis. Prolonged sub oxygenation, or bouts of anoxia from "incidents" during the anaesthetic are probably worse.

### CHOICE OF TECHNIQUE

The chest disease may make some ordinary anaesthetic techniques unsuitable. The

respiration cease to work as consciousness is lost and respiratory arrest may follow. In such a case a preliminary use of cocaine in the respiratory tract will facilitate immediate intubation, and as the endotracheal tube passes the obstruction it gives immediate relief. It is sometimes difficult to pass the obstruction without entering the right bronchus. If this occurs it becomes much easier for air to enter the right than the left lung. This causes differences in pleural pressure on the two sides and mediastinal shift results.

The fear of spasm is ever present in patients who give a history of asthma, and an induction with thiopentone may precipitate a severe laryngospasm. Fortunately ether is a reliable agent for such cases. Instead of more coughing and spasm as might be expected, the ether relaxes the bronchial musculature and the breathing

Indeed there is a marked relief of  
indicated by  
the adminis-  
tration of cocaine to the respiratory tract before the induction of anaesthesia, or by the use of an intravenous transfusion with a procaine solution.

### PROBLEMS PRESENTED BY PNEUMOTHORAX CASES

The patient with a pneumothorax may present difficulties. If the proposed operation requires the patient to lie in the lateral position with the pneumothorax undermost, the rising diaphragm on the underside and the sagging mediastinum may cause the pleural pressures in the artificial pneumothorax to rise so high that

In non-  
Too often  
rux has a  
larger than usual refill pre operatively in the hope of minimizing the effect of the anaesthetic on the underlying lung.

# ADDED PROBLEMS IN CHEST OPERATIONS

If a chest operation is proposed some of the problems particularly those of the control of the secretions and the maintenance of adequate ventilation become more urgent and special techniques must be used. Most chest operations involve an open pleura and in all paradoxical respiratory movement is likely to occur in areas of lung unsupported by a rigid chest wall.

This paradoxical movement makes respiration inefficient for some of the effort of inspiration draws air from the paradoxically moving portion of the lung whilst in expiration a proportion of the air which should be expelled from the trachea will distend the unsupported lung. This pendulum air takes no part in the respiratory exchange and may be the vehicle for dispersing secretions through previously healthy lung.

The mechanism of this paradoxical movement is that unsupported lung moves in response to pressure changes in the bronchi. This pressure is high during expiration and less than atmospheric during inspiration. These pressure differences are small with quiet unforced breathing. A patient well premedicated and free from anxiety who receives a careful local analgesia of the chest wall will breathe so quietly that the area of pleura exposed in a thoracoplasty for tuberculosis will move very little. If the patient becomes tense perhaps from inadequacy of the local analgesia the breathing becomes strained and paradoxical movement becomes apparent. The effort of working the valves of an anaesthetic machine increases the pressure differences so it is difficult to give a light general anaesthetic to cover any deficiencies in the local analgesia without causing marked paradoxical movement.

Nevertheless local analgesia must be accepted as a most useful technique for operations on the chest wall most notably for thoracoplasty and for the drainage of empyemas.

This rhythm of pressure changes can be reversed if with the patient in an apnoeic state respiratory exchange is maintained by an intermittent positive pressure anaesthesia. This controlled respiration ensures that all areas of lung fill during inspiration and empty in expiration thus eliminating the pendulum air with its effect of a greatly increased respiratory dead space.

The use of controlled respiration is not essential for chest surgery but it greatly increases the scope of surgery by keeping the patient fit and well oxygenated and by providing quiet conditions in which to operate. It does however demand a more careful control of secretions for the positive pressure of inspiration could easily drive secretions into small bronchioles which would be obstructed. In overcoming this difficulty besides the methods already outlined of posture and intra bronchial suction resort is often made to bronchial tamponade.

The anaesthetic may be given into one main bronchus the other lung being isolated from the anaesthetic circuit by an inflated balloon which encircles the tube. This is carried out more easily if the tube can be placed on the left side for the left main bronchus can accommodate the balloon and allow free drainage from the right lung outside the anaesthetic tube. When the endobronchial tube is passed into the right side the balloon must be partly in the trachea for the distance from the carina to the orifice of the right upper bronchus is too short to take the balloon.

As a result secretions are retained in the left lung, which threaten catastrophe should at any time the tube become displaced. The movement of these endobronchial tubes must always be feared. The position of the inflated cuff must be accurate to within a centimetre. The tube is fixed at the proximal end. During dissection traction is made on the hilum of the lung, and the tube becomes displaced into the trachea.

An alternative is to obstruct the bronchus of the diseased lobe or lung with a balloon which encircles a fine drainage tube, whilst the anaesthetic is given through an endotracheal tube into the remainder of the lung. These drainage blockers give good control of secretions and allow the diseased area of lung to collapse during the operation. There are two commonly used types of blocker. The

net gives a better adhesion than rubber to the mucosa of the bronchus. This balloon is bulky and can only be passed through an adult 11 millimetre bronchoscope. The Magill blocker is more delicately made and can be passed through a smaller bronchoscope which gives it a wider applicability, but more skill is required to prevent its becoming displaced during the operation. These drainage blockers can be used for pneumonectomies, and lower lobectomies on either side, but are not reliable for upper lobectomies as it is only occasionally possible to introduce them into the upper lobe orifice.

under lung. The removal of secretions is completed by intermittent intra bronchial suction, and by a post-operative bronchoscopy.

If the secretions are controlled, and paradoxical movement is prevented by the use of controlled ventilation, the anaesthesia required for thoracic surgery is simple. An induction with thiopentone and maintenance with nitrous oxide and oxygen will suffice if curare is given to allow the over ventilation necessary to

the patient appears to be of value in preventing post operative respiratory complications.

### SUMMARY

To summarize, it may be said that local analgesia should be considered if it is suitable for the operation proposed and if respiratory distress is not a marked feature of the disease. Routine general anaesthesia is safer than experimenting with new methods and in particular ether despite its irritant properties is safer than the injudicious use of the short acting barbiturates. In acute respiratory infections with dyspnoea cyclopropane is valuable to those experienced in using it. Light general anaesthesia used with muscle relaxants is almost universally adopted for chest surgery and is quite suitable for general surgery in the presence of chest disease if secretions are well controlled and respiration is assisted.

## CHAPTER 32 PHYSIOTHERAPY FOR CHEST DISEASES

JOCELYN M W REED

THE NEED for expiratory breathing exercises for asthmatic and bronchitic patients became widely recognized in about the year 1930 and their value was proved in the asthma research clinics. Since then the necessary training in relaxation and in the development of an easy good carriage or posture has been intensified with improvement in results. Thoroughness with careful attention to detail are essential for good results in the physical treatment of chest conditions since the movements of the chest are comparatively small and are carried out in a series of joints set at varying angles. The patient therefore has no understanding of the capabilities of his chest under voluntary control. The conditions in which the treatment is of a similar type may be grouped together. The first group includes bronchitis, emphysema and asthma, the second comprises pneumonia and pleurisy, the third lung abscess, collapse and bronchiectasis and the fourth pulmonary tuberculosis.

### BRONCHITIS, EMPHYSEMA AND ASTHMA

Patients suffering from these conditions generally respond best to treatment in groups although the main breathing exercises and particular faults of posture as well as training in relaxation need individual attention and correction which can be given before and after the general part of the class. Adult patients derive psychological benefit from finding others in a worse condition than themselves and from hearing of the improvement made by others. With children there is scope for imitation and competition in class work.

The aims of treatment are the same in these three conditions: (1) to gain relaxation, (2) to promote breathing control and (3) to develop good posture and balance.

#### Attainment of relaxation

The accessory muscles of inspiration are over-toned and tightened to a varying degree. The pectorals, sternomastoids, the upper fibres of trapezius, the scaleni and all the adductor muscles of the arm are involved. To many patients relaxation of these muscles is difficult or even impossible at first and in bad cases there is a general inability to relax fully any of the large muscle groups of the whole body. Those needing a great deal of careful training in relaxation belong to the anxiety type, mainly asthmatics. They are never mentally and therefore never physically relaxed. Some of them cannot keep up with their work and the more they worry the more tense they become. All adult patients and some children need training in general relaxation as well as in local relaxation of their over-used muscles of inspiration. It is important that a simple explanation of the necessity and importance of relaxation should be



given in a confident and friendly manner before the treatment starts, but when the patient has become accustomed to his surroundings

*General relaxation*—The patient lies on his back with a pillow under his head, another under his knees, and one under the whole of each arm. The arms are slightly bent at the elbows, with the palms of the hands downwards, so that the sides of the thumbs almost touch the body.

The patient is shown, the instructor demonstrating with her own arm, the difference between tension and relaxation of the muscles. It helps him to understand what is meant by relaxation if he is allowed to hold her arm. First the arm is stiffened so that the muscles feel hard, then little by little, the muscles are allowed to relax, so that the patient feels how the arm gradually gets heavier and heavier, until he finally feels its complete relaxed weight. Attention is directed in turn to large muscle-groups, starting with those of the hand, forearm, shoulder, neck and so on. The patient "talks" to his muscles in his mind telling them gently and frequently, always on a sighing out breath, "let go". This is not a breathing exercise but just natural breathing, which, gradually, will be found to become slower and deeper and more diaphragmatic. The patient needs to practise daily for about half an hour at a regular time, and it will be found that the general well being as well as the breathing improves. This method is based on a combination of those described by Jacobson (1938) and Fink (1945). It is simple and successful with chest patients when taught with patience and perseverance.

*Local relaxation*—Local relaxation of the tightened inspiratory muscles is gained by rhythmic swinging exercises for the arm and shoulder muscles, and by rhythmic stretching and relaxing shoulder-girdle and spinal movements. This will be done during the class part of the treatment.

### Breathing control

It is not uncommon to find these patients apparently breathing entirely with their upper chests, and for screening to prove that the diaphragmatic movements are poor, sometimes even paradoxical.

These breathing exercises are designed first to train relaxation and to control the overworking upper chest, so that it is reduced in size and the stretched lung tissue and bronchial tubes are rested. This is done by relaxation training and by slow expiratory breathing exercises.

Secondly, they develop in the patient a longer expiratory phase of breathing, to help him in his expiratory difficulty, and to counteract his habit of over inspiration. The residual air will thus be reduced. Thirdly, the exercises develop fully lower costal and diaphragmatic movements so that the patient can make full use of the bases of his lungs. In emphysema, when the patients have raised barrel shaped chests, this procedure is modified. They should keep entirely to expiratory breathing without any attempt to gain further expansion.

Long expiratory diaphragmatic breathing shown in Fig. 184 and expiratory lateral basal expansion exercises shown in Figs. 185–186 are the main ones used to develop longer expiration with the fullest use of the lung bases and of the lower half of the chest generally. Expiration should last for twice as long as inspiration. Good lower costal movement is actually more easily acquired if practised for one

FIG 184—Long expiratory diaphragmatic breathing exercise (a) Full expiration with good contraction of upper abdominal muscles and relaxation of upper chest, (b) inspiration with relaxation of abdominal muscles. Note control of upper chest which is still in relaxed position (B, courtesy Institute for Diseases of the Chest)



side separately for a time, as shown in Fig 185, before adding bilateral expansion as in Fig 186

Short expiratory diaphragmatic breathing is used to overcome breathlessness or to prevent or stop an asthmatic attack developing. The exercise is described in Fig 187. Positions are arranged so that gravity helps relaxation of the abdominal muscles on inspiration. Positions shown in Fig 187 (a) and (b) are used for children and some younger adults. It can be modified for adults into a sitting, leaning forward position. The position in which the body is in a simple resting posture, slightly rotated forward, is used also for teaching diaphragmatic breathing in difficult cases (Fig 187c).

Control of breathing is practised during strenuous exercise and during ordinary activities such as going upstairs. Short expiratory diaphragmatic breathing is

given in a confident and friendly manner before the treatment starts, but when the patient has become accustomed to his surroundings

*General relaxation*—The patient lies on his back with a pillow under his head, another under his knees, and one under the whole of each arm. The arms are slightly bent at the elbows, with the palms of the hands downwards, so that the sides of the thumbs almost touch the body.

The patient is shown, the instructor demonstrating with her own arm, the difference between tension and relaxation of the muscles. It helps him to understand what is meant by relaxation if he is allowed to hold her arm. First the arm is stiffened so that the muscles feel hard, then little by little, the muscles are allowed to relax, so that the patient feels how the arm gradually gets heavier and heavier, until he finally feels its complete relaxed weight. Attention is directed in turn to large muscle-groups, starting with those of the hand, forearm, shoulder, neck and so on. The patient "talks" to his muscles in his mind, telling them gently and frequently, always on a sighing out breath, "let go". This is not a breathing exercise but just natural breathing, which, gradually, will be found to become slower and deeper and more diaphragmatic. The patient needs to practise daily for about half an hour at a regular time, and it will be found that the general well-being as well as the breathing improves. This method is based on a combination of those described by Jacobson (1938) and Fink (1945). It is simple and successful with chest patients when taught with patience and perseverance.

*Local relaxation*—Local relaxation of the tightened inspiratory muscles is gained by rhythmic swinging exercises for the arm and shoulder muscles, and by rhythmic stretching and relaxing shoulder-girdle and spinal movements. This will be done during the class part of the treatment.

### Breathing control

It is not uncommon to find these patients apparently breathing entirely with their upper chests, and for screening to prove that the diaphragmatic movements are poor, sometimes even paradoxical.

These breathing exercises are designed first, to train relaxation and to control the overworking upper chest, so that it is reduced in size and the stretched lung tissue and bronchial tubes are rested. This is done by relaxation training and by slow expiratory breathing exercises.

Secondly, they develop in the patient a longer expiratory phase of breathing, to help him in his expiratory difficulty, and to counteract his habit of over-inspiration. The residual air will thus be reduced. Thirdly, the exercises develop fully lower costal and diaphragmatic movements so that the patient can make full use of the bases of his lungs. In emphysema, when the patients have raised barrel-shaped chests, this procedure is modified. They should keep entirely to expiratory breathing without any attempt to gain further expansion.

Long expiratory diaphragmatic breathing shown in Fig. 184 and expiratory lateral basal expansion exercises shown in Figs. 185–186 are the main ones used to develop longer expiration with the fullest use of the lung bases and of the lower half of the chest generally. Expiration should last for twice as long as inspiration. Good lower costal movement is actually more easily acquired if practised for one

FIG 184—Long expiratory diaphragmatic breathing exercise (a) Full expiration with good contraction of upper abdominal muscles and relaxation of upper chest (b) inspiration with relaxation of abdominal muscles. Note control of upper chest which is still in relaxed position (By courtesy Institute for Diseases of the Chest)



side separately for a time, as shown in Fig 185 before adding bilateral expansion as in Fig 186

Short expiratory diaphragmatic breathing is used to overcome breathlessness or to prevent or stop an asthmatic attack developing. The exercise is described in Fig 187. Positions are arranged so that gravity helps relaxation of the abdominal muscles on inspiration. Positions shown in Fig 187 (a) and (b) are used for children and some younger adults. It can be modified for adults into a sitting leaning forward position. The position in which the body is in a simple resting posture, slightly rotated forward, is used also for teaching diaphragmatic breathing in difficult cases (Fig 187c).

Control of breathing is practised during strenuous exercise and during ordinary activities such as going upstairs. Short expiratory diaphragmatic breathing is




FIG 185—Unilateral basal exercises for bronchitis and collapse of right lower lobe

left palm




FIG 186—Bilateral basal expansion exercise (a) End of slow expiratory phase with pressure of both palms applied in lateral costal area, (b) inspiration with full expansion outwards of both sides against slight palmar pressure

used and when going upstairs is timed to the movements. This is first taught on the level.

Breathing is arranged rhythmically and slowly, in time with some of the postural muscle strengthening exercises to help breathing control. The patient is trained gradually to tolerate more and more exercise without becoming breathless.

The psychological approach to the asthmatic patient is slightly different from that to the emphysematous or bronchitic patient. These asthmatic patients have to be inspired with confidence in the treatment, and to learn that they can do something themselves to prevent an attack from coming on, instead of having to



FIG 187 —Pos tions for overcoming breathlessness (a) Contraction of upper abdominal muscles on expiration (quickly) (b) relaxation of abdominal muscles on inspiration (quickly) (c) position for short diaphragmatic breathing exercise



rely entirely on outside help. Much depends therefore on the personality of the physiotherapist. The patient is taught to rely on the short expiratory diaphragmatic exercise to prevent the development of an attack or even to stop one that may have developed.

#### Development of good posture and balance

This is the third aim. It is achieved by postural muscle strengthening exercises and by developing good posture through particular attention to it during the mobility exercises in class. Remedial exercises are given for individual posture deformities.

It is essential to develop a well shaped thorax and good carriage otherwise good lower-chest breathing will not be maintained.

The patient must stand so that the upper abdominal muscles can be relaxed. An imaginary plumb-line dropped from the shoulder should pass through the hip (iliac crest just above the anterior iliac spine) and on down to the forepart of the foot where the weight is carried. This makes it possible for the abdominal muscles to be relaxed and thus allow the patient to breathe diaphragmatically.

He must learn to carry himself easily and loosely.

Constant enquiry into the amount and regularity of practice is essential. Asthmatic and emphysematous patients need much encouragement to help them control breathlessness and thus prevent the development of attacks.

For all chest patients the vital-capacity estimates, chest measurements and careful assessment of chest movements and postural deformities are made before starting treatment. The vital capacity is measured at regular intervals. These tests stimulate the patient to further effort as well as being a gauge of improvement.

*Children with catarrhal bronchitis*—For these, training in nose instead of mouth breathing is necessary. Limbering up, strengthening postural and balance exercises are needed more than relaxation training. Most of the limbering-up exercises are given in the form of rhythmic swinging or "stretch relax" exercises, these bring about relaxation of the tightened accessory muscles of inspiration.

Strong remedial exercises, some in kneeling positions and some in crawling positions, are used for bigger children. Good posture is encouraged by the use of mirrors. Later, more difficult balance exercises replace the mirror correction.

### Other physical treatment

Many of these patients are given a course of general ultra-violet light, administered on alternate days for 20 doses for its general stimulating effect. The dose is 1000 rads.

the horny dead layers of cells on the surface. Another course may be given after a 2 or 3 months' gap.

Short-wave diathermy is beneficial in the treatment of nasal sinusitis partly for its bactericidal effect and partly to produce hyperaemia. It can be applied to the chest wall as a form of treatment for asthma, emphysema and bronchitis, in order to increase the circulation.

## PNEUMONIA AND PLEURISY

### Pleurisy

Since pleurisy which does not result from pneumonia is most often of tuberculous or carcinomatous origin, breathing exercises should not be given.

### Pneumonia

It is most important that breathing exercises should be given early after pneumonia to help re-expand partially collapsed bases, to prevent the development of fibrosis and to expedite resolution. Modified exercises can safely be given during broncho-pneumonia, and this is done in many London hospitals and thoracic units with good results.

The best form of inspiratory localized breathing exercises are called "pressure expansion exercises". Pressure is used to gain localization of movement and to train the inspiratory muscles by giving them resistance. The exercises are used to develop full use of the lung bases, particularly on the affected side. These exercises are described in detail later, because they are the same as those used for surgical chest conditions.

## LUNG ABSCESS, COLLAPSED LOBES AND BRONCHIECTASIS

Physical treatment in these conditions is directed to draining the muco purulent secretions arising from the abscess or the bronchiectatic bronchi, and to regaining expansion in those parts of the chest where respiration has been restricted because of the pathological condition

The elimination of muco purulent secretion is facilitated by postural drainage and expiratory breathing exercises, and by tapôtment or percussion treatment consisting in clapping and pressure-vibration

The posture required for successful drainage is decided by consulting the bronchograms. It is of importance to find out which bronchi are draining an abscess. One or two variations of the apparently anatomically correct posture must often be tried out before good drainage occurs (Fig 188)

Clapping and pressure-vibrations, during long expirations, are the most effective forms of mechanical stimulus to elimination of secretions. They are carried out over the affected areas, for bronchiectasis, during the first few minutes of postural drainage. For lung abscess clapping should not be allowed, because it might cause haemoptysis, fine vibrations, with a little pressure during prolonged expiration, are used instead. In cases in which severe haemoptysis complicates

ed should be practised in expiratory manner

After this time, if there is still restricted movement, they can be practised in the full inspiratory manner, as described for post-operative surgical patients

## PULMONARY TUBERCULOSIS

Patients about to get up after prolonged rest in bed gain considerable benefit from physiotherapy. It is particularly useful for "poor breathers", for patients suffering from nervous tension, and in cases of long standing pleurisy

*Inadequate respiration*

The poor or bad breathers are those who have suffered from breathlessness and who have therefore developed a shallow upper-chest type of breathing. This type of respiration is detrimental in patients with bilateral lesions. Patients with a low vital capacity who are awaiting thoracoplasty also improve with controlled breathing exercises

*Controlled breathing*

This controlled breathing consists in careful, gentle, localized development of the area of the chest where the patient has sound lung, combined with training in complete relaxation and control of the lesion areas. A patient trained thoroughly and carefully once or twice a day takes between 4 and 8 weeks to get such good control that, when asked to take a deep breath, he only expands the lower chest

The patient with a low vital capacity is trained in diaphragmatic breathing and in careful localized lateral basal expansion of each side, at first separately. The vital capacity can usually be raised by 400-700 millilitres in about 3 weeks

*Nervous tension*

Restless, nervous, worried patients can be helped by training in general physical






FIG 183—Postural drainage positions (a) For drainage of the posterior segment of the right upper lobe. (Note the quarter turn or lift backwards for rotation)

(e)



(f)

F1

(1943) and to practical experience with a large number of cases )

relaxation of the same type as is taught to asthmatic patients. The general sense of physical well being induced brings about a release of nervous tension.

### Pleurisy

Patients with a history of long standing tuberculous pleurisy are sometimes left with a painful stiff chest having restricted movement. This condition can be eased by careful localized expansion exercises of the lateral basal area of the affected side, combined with good training in diaphragmatic breathing.

### Strengthening exercises

Graduated strengthening exercises for all the main anti gravity muscle groups

are important after a long period of rest in bed. They should be started weeks before the patient is to get up, so that the strength of the foot, leg, back and neck exercises can be gradually increased. This prevents poor posture and flat feet. In suitable wards these exercises can be performed in a graduated class.

## PHYSIOTHERAPY IN SURGICAL CONDITIONS

Pre-operative training in breathing control is of importance in almost all surgical chest conditions, acute empyema being the only exception, but it is in the early post-operative period that localized breathing exercises are essential for the patient's uneventful and speedy recovery. In this period post-operative collapse can be prevented by establishing good functioning of the remaining lung or lobes. Between 1 and 2 weeks' pre-operative training is necessary to establish good control and develop localized expansion movements, unless the patient has a low vital capacity, when 3 or 4 weeks may be necessary.

### Pre-operative physiotherapy

Pre-operative exercises in chest surgery can be divided into three main groups (1) localized breathing, (2) posture and (3) shoulder movements.

#### *Localized breathing exercises*

Localized breathing exercises are important to establish breathing control in any cases of thoracotomy for a removal of cysts and non-malignant tumours, after empyema drainage or decortication, and in all cases of lung resection.

In order to bring about good localized movement of any one part, the rest of the chest must be relaxed. Training in physical relaxation is therefore necessary to gain breathing control.

In all non-tuberculous and non-malignant chest conditions, chest rehabilitation includes the development of full equal bilateral chest movements, and restoration of the ability of each patient to undertake his normal activities and exercise without breathlessness.

#### *Postural training*

Secondly, a good balanced stance or posture should be taught, as previously described for bronchitic and asthmatic patients. This can also be achieved in bed, sitting in a good position, with pillows, arranged to help to maintain it. Small corrective active movements should be practised to prevent each possible post-operative fault. Pre-operative and post-operative exercises which teach "limbering-up" and postural muscle-strengthening, should be interspersed with the localized breathing exercises.

#### *Arm and shoulder movements*

Thirdly, arm movements must be maintained by relaxed active shoulder movements taught pre-operatively. They include scapular movements with a relaxed arm, gleno-humeral movements with the scapula controlled, and combined shoulder girdle movements. The movements are isolated and controlled in order not to stretch the incision and stitches, and to maintain good functioning with minimal physical exertion in the early post-operative period. The latter reason is particularly important for tuberculous patients.

*Postural drainage*

In all cases of bronchiectases and lung abscess, intensive pre-operative postural drainage forms the fourth and most important part of their treatment. The patient's lung should be as dry as possible before operation, so that post operative collapse, caused by an overflow of secretions into neighbouring bronchi, may be prevented.

**Localized breathing exercises***Diaphragmatic breathing*

Prior to operation, diaphragmatic breathing is of great importance in all types of major chest surgery, because it develops the use of the lung bases and helps to keep them clear of mucus in the early post-operative period. Localization of movement to the epigastric and antero lateral lower costal area is taught in conjunction with relaxation and control of the upper chest.

During the major part of their pre operative treatment, patients with bronchiectasis and lung abscess are taught the same slow expiratory type of diaphragmatic exercise as used for asthmatic patients, to help in the elimination of the muco-purulent secretions. In other surgical conditions, the accent of the exercise is equally on expiration with contraction of the upper abdominal muscles and on normal diaphragmatic expansion. The patient should keep his finger-tips resting lightly on his lower front ribs in order to facilitate consciousness of the movement. Actual pressure should never be applied because it produces trick movements. For a few days before operation all patients are taught a shortened form, that is the sharp expiratory diaphragmatic breathing used in cases of asthma. This is an aid to the elimination of post anaesthetic mucous secretions immediately after the operation, especially in cases of lung resection.

*Localized costal-expansion exercises*

The aim of these exercises is to stimulate good expansion of any areas of the chest where movement is poorly developed or impaired through disease or injury, and at the same time to inhibit other chest movements. This inhibition of movement is achieved by relaxation and control, coupled with good expansion of any one part of the chest, it ensures that most of the air inspired will be drawn into the expanding part of the lung.

Pressure is applied over the moving costal area in order to focus the attention of the patient on the site and direction of movement required, and, through the resistance given, to develop muscle tone. Pressure is applied at first by hand and later by using a belt.

For these purposes the chest is divided into four main areas. (1) *The lateral basal area*, in which expansion of the chest is outwards against pressure applied by the palm or knuckles on the side of the lower ribs. (For position of hands see Fig 185. Figure 189 shows the pressure applied by the belt.)

(2) *The upper lateral area*, in which expansion of the chest is outwards against pressure applied by either the palm of the hand or the fingers of the opposite hand, placed high on the side (Fig 190 a and b).

(3) *The pectoro-apical area*, in which expansion of the chest is upwards and forwards against pressure applied by the palmar surfaces of the fingers, placed




FIG. 189—T.W. (age 17 years) has had a right middle and lower lobectomy for bronchiectasis. He is having localized basal expansion exercise. (a) breathing in while expanding lower ribs against the belt. (b) breathing out showing the belt pulled firmly prior to next expansion.

over the upper chest. The tips of the fingers rest just below the clavicle, the opposite hand being used (Fig. 190 c and d).

(4) *The posterior basal area* in which expansion of the chest is mainly backwards against pressure applied by a belt placed across the back, covering the upper posterior basal area.


These localized costal exercises are practised pre-operatively in areas of the chest where movement is poorly developed or has become restricted and after operation when it is desired that full expansion shall be maintained or regained. As soon as movement has improved on the weak side to about three quarters of that on the sound side, bilateral movement is practised. This is necessary to develop the normal co-ordinated movements of both sides of the chest.

The method of applying pressure is first taught by the physiotherapist by applying it herself. It is then practised by the patient with his own manual pressure. Later a belt is used for lower and upper costal expansion exercises because the shoulders can be more relaxed. The belt crosses in front, the piece coming from the left side being held by the right hand and that from the right side by the left hand (Fig. 189).

Localized costal exercises can be practised in an expiratory manner or in an



(a)



(b)



(c)



(d)

Fig. 100. . . . . removal of ne. trophoma (right upper

inspiratory manner. In the former case expiration takes at least twice as long as inspiration and it is accompanied by extra pressure towards the end of the expiratory phase. The exercises are given in this manner prior to operations for bronchiectasis and lung abscess and in cases of asthma and emphysema. The inspiratory manner is used for empyema after thoracotomy, and after lobectomy and pneumonectomy. It is also used to stimulate the expansion of collapsed lobes and in convalescence from pneumonia.

When good expansion is required the patient is taught to breathe in with as strong expansion as possible against applied pressure, maintain the expansion while holding the breath and finally relax while breathing out. This is followed by a pause before repeating the movements.

Each of the three phases are of equal duration. The middle phase however, should not be allowed if a fistula is present.

### Physiotherapy after thoracotomy

Physiotherapy is needed after thoracotomy performed to treat empyema or for removal of tumours, cysts, neurofibroma or foreign bodies (when lung resection is not performed) or for ligation of a patent ductus arteriosus.

*Breathing exercises*—Breathing exercises should be practised carefully for 1 2

expansion exercises of the full inspiratory type are taught for each area of the affected side of the chest. Normal diaphragmatic breathing and bilateral basal expansion are practised to develop full use of the lung bases.

*Postural exercises*—Simple posture corrective movements are taught to help hip, spine and shoulder alignment. These movements are important while the patient is in bed but as soon as possible graded class exercises for muscle strengthening and posture training replace them.

### Physiotherapy before and after pulmonary resection

#### *Surgical conditions other than tuberculosis or carcinoma*

In cases in which lobectomy or pneumonectomy is performed for diseases other than tuberculosis or carcinoma pre-operative treatment includes a course of 3-5 weeks postural drainage with tapotement or percussion massage. This consists in clapping (except when there is a history of haemoptysis) and of pressure vibrations applied during prolonged expiration. These manipulations should be carried out alternately for the first 3-5 minutes of each tipping session. Classes should be held daily for limbering up, localized expiratory breathing exercises and good posture training. During the week before the operation the patient is taught short sharp expiratory diaphragmatic breathing and localized costal expansion exercises in the full inspiratory manner for the remaining sound lobe or lobes which will be needed in the immediate post-operative phase. In preparation for lower lobe lobectomy, expansion exercises are practised in the




FIG. 191—Same patient as shown in FIG. 189. A—Chest expansion exercise.

pectoro apical and upper lateral areas. In the immediate post operative phase good expansion of the upper lobe should be maintained. In 3-4 days when the skiagrams show that there is full upper lobe expansion lateral basal expansion is started to expand the upper lobe downwards. By the end of 2-3 weeks expansion should have increased so that there is no space between the upper lobe and the diaphragm.

For patients undergoing middle lobe lobectomy since the lower lobe readily tends to collapse mainly posterior and lateral basal expansion exercises should be practised in the pre operative and immediate post operative phases.

For patients submitted to pneumonectomy although all areas of the affected side should be well exercised beforehand lateral basal and upper lateral costal expansion of the sound side and diaphragmatic breathing are the important exercises. When these movements are well developed about a week after operation expansion movements on the side of the pneumonectomy are practised to prevent the development of scoliosis. Bilateral expansion exercises for the lower and middle-chest develop normal co-ordinated movements.

#### *Operations for tuberculosis or carcinoma*

When lobectomy or pneumonectomy is performed for tuberculosis or carcinoma



inspiratory manner. In the former case expiration takes at least twice as long as inspiration and it is accompanied by extra pressure towards the end of the expiratory phase. The exercises are given in this manner prior to operations for bronchiectasis and lung abscess and in cases of asthma and emphysema. The inspiratory manner is used for empyema after thoracotomy and after lobectomy and pneumonectomy. It is also used to stimulate the expansion of collapsed lobes and in convalescence from pneumonia.

When good expansion is required the patient is taught to breathe in with as strong expansion as possible against applied pressure, maintain the expansion while holding the breath and finally relax while breathing out. This is followed by a pause before repeating the movements.

Each of the three phases are of equal duration. The middle phase however should not be allowed if a fistula is present.

### *Physiotherapy after thoracotomy*

Physiotherapy is needed after thoracotomy performed to treat empyema or for removal of tumours, cysts, neurofibroma or foreign bodies (when lung resection is not performed) or for ligation of a patent ductus arteriosus.

*Breathing exercises*—Breathing exercises should be practised carefully for 1-2 weeks pre-operatively (except for empyema) and started within 24 hours of the operation so that the lung may be re-expanded before the pleura has time to become thickened or fibrous scar tissue and adhesions can be formed. Localized costal expansion exercises of the full inspiratory type are taught for each area of the affected side of the chest. Normal diaphragmatic breathing and bilateral basal expansion are practised to develop full use of the lung bases.

*Postural exercises*—Simple posture corrective movements are taught to help hip, spine and shoulder alignment. These movements are important while the patient is in bed but as soon as possible graded class exercises for muscle strengthening and posture training replace them.

*Arm and shoulder movements*—Relaxed active shoulder movements are changed to resisted movements and free exercises as soon as possible (Fig. 191). Elevation of the arm should be of full range by the fifth or sixth day.

### *Physiotherapy before and after pulmonary resection*

#### *Surgical conditions other than tuberculosis or carcinoma*

In cases in which lobectomy or pneumonectomy is performed for diseases other than tuberculosis or carcinoma, pre-operative treatment includes a course of 3-5 weeks postural drainage with tapotement or percussion massage. This

is done in the inspiratory manner with the patient in the prone position. The hands are placed over the lower lobes and the upper lobes respectively. The patient is then turned on to his side and the same procedure is repeated. This is followed by a period of rest.

and good posture training. During the week before the operation the patient is taught short sharp expiratory diaphragmatic breathing and localized costal expansion exercises in the full inspiratory manner for the remaining sound lobe or lobes which will be needed in the immediate post-operative phase. In preparation for lower lobe lobectomy, expansion exercises are practised in the



(a)



(b)

Fi

arm on the affected side

pectorapical and upper lateral areas. In the immediate post operative phase good expansion of the upper lobe should be maintained. In 3-4 days when the skiagrams show that there is full upper lobe expansion lateral basal expansion is started to expand the upper lobe downwards. By the end of 2-3 weeks expansion should have increased so that there is no space between the upper lobe and the diaphragm.

For patients undergoing middle lobe lobectomy since the lower lobe readily tends to collapse mainly posterior and lateral basal expansion exercises should be practised in the pre operative and immediate post operative phases.

For patients submitted to pneumonectomy although all areas of the affected side should be well exercised beforehand lateral basal and upper lateral costal expansion of the sound side and diaphragmatic breathing are the important exercises. When these movements are well developed about a week after operation

of the pneumonectomy are practised to

Bilateral expansion exercises for the co-ordinated movements

#### *Operations for tuberculosis or carcinoma*

When lobectomy or pneumonectomy is performed for tuberculosis or carcinoma

**inspiratory manner** In the former case expiration takes at least twice as long as inspiration, and it is accompanied by extra pressure towards the end of the expiratory phase. The exercises are given in this manner prior to operations for bronchiectasis and lung abscess, and in cases of asthma and emphysema. The inspiratory manner is used for empyema, after thoracotomy, and after lobectomy and pneumonectomy. It is also used to stimulate the expansion of collapsed lobes and in convalescence from pneumonia.

When good expansion is required the patient is taught to breath in with as strong expansion as possible against applied pressure, maintain the expansion while holding the breath, and finally relax while breathing out. This is followed by a pause before repeating the movements.

Each of the three phases are of equal duration. The middle phase however, should not be allowed if a fistula is present.

### **Physiotherapy after thoracotomy**

Physiotherapy is needed after thoracotomy performed to treat empyema or for removal of tumours, cysts, neurofibroma or foreign bodies (when lung resection is not performed) or for ligation of a patent ductus arteriosus.

**Breathing exercises**—Breathing exercises should be practised carefully for 1-2 weeks pre-operatively (except for empyema) and started within 24 hours of the operation, so that the lung may be re-expanded before the pleura has time to become thickened or fibrous scar tissue and adhesions can be formed. Localized costal expansion exercises of the full inspiratory type are taught for each area of the affected side of the chest. Normal diaphragmatic breathing and bilateral basal expansion are practised to develop full use of the lung bases.

**Postural exercises**—Simple posture corrective movements are taught to help hip, spine and shoulder alignment. These movements are important while the patient is in bed, but as soon as possible graded class exercises for muscle strengthening and posture training replace them.

**Arm and shoulder movements**—Relaxed active shoulder movements are changed to resisted movements and free exercises as soon as possible (Fig 191). Elevation of the arm should be of full range by the fifth or sixth day.

### **Physiotherapy before and after pulmonary resection**

#### *Surgical conditions other than tuberculosis or carcinoma*

In cases in which lobectomy or pneumonectomy is performed for diseases other than tuberculosis or carcinoma, pre-operative treatment includes a course of 3-5 weeks postural drainage with tapotement or percussion massage. This consists in clapping (except when there is a history of haemoptysis) and of pressure-vibrations applied during prolonged expiration. These manipulations should be carried out alternately for the first 3-5 minutes of each tipping session. Classes should be held daily for limbering up, localized expiratory breathing exercises and good posture training. During the week before the operation, the patient is taught short sharp expiratory diaphragmatic breathing and localized costal expansion exercises in the full inspiratory manner for the remaining sound lobe, or lobes, which will be needed in the immediate post-operative phase. In preparation for lower lobe lobectomy, expansion exercises are practised in the




FIG 192 — L.M. (aged 30 years) after second stage right 8th rib thoracoplasty (a) Correction of posture in front of mirror sitting up away from pillows between 2-3 weeks after operation (b) lateral deviation movement of neck against resistance to correct natural tendency to left deviation

of the long incision, the postural and arm movements are isolated to a single movement at a time

All the factors concerned with the maintenance of good posture are demonstrated with the use of a mirror placed at the foot of the bed (Fig 192a—here the patient is seen 3 weeks after second stage thoracoplasty). Corrective movements are practised for any present faults and to prevent those likely to occur after operation (Fig 192b—here neck correction is being practised 3 weeks after right second stage thoracoplasty).

Diaphragmatic breathing is practised in the patient's own time and with complete relaxation of the upper chest. This is the most important breathing exercise, ensuring good use of the base of the lung on the thoracoplasty side and helping to prevent paradoxical breathing in the upper chest after operation.

Localized postero-lateral basal expansion is practised on the side for thoracoplasty. A little pressure is applied to help the patient to feel the movement, but it is applied gently so that it is localized. This exercise is essential for patients

pre operative physical treatment consists in 1-2 weeks' individual training in relaxation and control of the chest, coupled with diaphragmatic breathing and with small localized unilateral expansion movements of the healthy parts of the lungs. In tuberculous cases the amount of good diaphragmatic movement that is possible will depend on whether or not there has been a phrenic avulsion or a pneumoperitoneum is present. Relaxed active shoulder movements and postural corrective exercises are practised before operation but they are of secondary importance to the controlled breathing exercises.

In preparation for upper lobectomy, small pectoro apical expansion movements are practised on the sound side for a few days if there is a large cavity in the upper lobe on the affected side. If there is infiltration without cavity formation in the upper lobe, this pectoro apical expansion can be practised gently on the affected side. After operation, this exercise and upper lateral costal expansion are used to help to expand the lower lobe upwards. As soon as expansion is complete (in about 2 weeks) breathing exercises are stopped. Postural corrective movements and graduated muscle strengthening exercises may be needed for a further 2 weeks and should be supervised until the patient is ambulant. Foot and leg exercises should be practised for 2 weeks before the patient gets up.

When pneumonectomy is performed for carcinoma the patients tend to be elderly, to have stiff chests, and to use the upper parts excessively. In the short pre operative period good diaphragmatic movement and unilateral lower costal expansion of the healthy side are developed. If these breathing exercises are practised frequently in the early post operative phase, there is little danger of collapse of the lower lobe, or of pneumonia caused by infection of retained secretions. Graduated foot and leg exercises should be started on the day after

These patients usually begin exercises are continued and

patients

When pneumonectomy is performed on account of tuberculosis, similar breathing exercises are used but the foot and leg exercises are postponed until a later date. Postural corrective movements need more attention with younger and more mobile patients, some of whom have poor postural muscle tone.

### Physiotherapy for plastic operations

In these operations there are extensive muscle incisions and rib resections, and therefore postural muscle strengthening and training are essential.

#### Thoracoplasty

*Pre operative exercises*—Many of the patients who undergo thoracoplasty have poor muscle tone and posture resulting from long periods of rest in bed and of curtailed physical exertion. Some have a shallow upper chest type of respiration with low vital capacity. Pre operative training is needed for between 1 and 3 weeks. Care should be taken to graduate these movements and exercises in order not to tire the tuberculous patient. For this reason, and to prevent any stretching



(a)



(b)

FIG. 192 — L.M. (aged 30 years) after second stage right 8th rib thoracoplasty (a) Correction of posture in front of mirror sitting up away from pillows between 2-3 weeks after operation (b) lateral deviation movement of neck against resistance to correct natural tendency to left deviation

of the long incision, the postural and arm movements are isolated to a single movement at a time

All the factors concerned with the maintenance of good posture are demonstrated with the use of a mirror placed at the foot of the bed (Fig. 192a—here the patient is seen 3 weeks after second stage thoracoscopy). Corrective movements are practised for any present faults and to prevent those likely to occur after operation (Fig. 192b—here neck correction is being practised 3 weeks after right second stage thoracoscopy).

Diaphragmatic breathing is practised in the patient's own time and with complete relaxation of the upper chest. This is the most important breathing exercise ensuring good use of the base of the lung on the thoracoplasty side and helping to prevent paradoxical breathing in the upper chest after operation.

Localized postero-lateral basal expansion is practised on the side for thoracoplasty. A little pressure is applied to help the patient to feel the movement, but it is applied gently so that it is localized. This exercise is essential for patients

undergoing general anaesthesia. Training in relaxation of shoulder and arm muscles is needed, so that movements can be isolated, so far as is possible, to individual muscle-groups. Movements of the scapula, with relaxed arm-depression and retraction, and gleno-humeral movements with controlled scapula are taught. Those most used are elbow raising to shoulder level in different planes to abduct the arm, extension, and outward rotation.

All shoulder-joint movements are performed with a bent elbow, in order to give the patient a small weight to lift and therefore the least strain on the muscles controlling the scapula. The arm is elevated above the head, with assistance at first (Fig 191a), and later, when the movement is practised by the patient, both arms are raised, the palms meeting (Fig 191b). Ultimately he can separate the arms above the head, and practise raising the affected arm by itself.

*Post operative exercises*—After the second-stage operation, when the ribs beneath the scapula have been removed, elevation of the arm should not be practised for 5-7 days. Until the scapular movements are strong, elevation is painful, and it can shift the scapula when it has been placed forwards with its inferior angle beneath the seventh and eighth ribs.

In the event of post-operative collapse of the lower lobe, postural drainage is arranged for half an hour 3 times a day for a few days. The patient is placed on his sound side with a pillow support for his head and neck, which should not be allowed to drop towards the sound side. The foot of the bed is raised 16 inches from the floor. Diaphragmatic and postero-lateral basal expansion exercises are practised with expiratory emphasis.

Postural muscle-strengthening exercises should be continued until the patient is allowed up, and then should be supervised once or twice a week either in ward classes or individually. Foot and leg exercises are started in the last weeks prior to getting up. When the patient is up, classes in graduated posture training, walking and balance exercises prevent postural deformities arising from bad habits of stance.

### *Extrapleural pneumolysis*

In cases of extrapleural pneumolysis, the postural back muscle incisions are nearly as extensive as in thoracoplasty. There is, therefore, a similar, though less severe, temporary imbalance of the upper thoracic extensor and shoulder retractor muscles. Pre operative training in localized postural corrective movements and diaphragmatic breathing, with relaxation of the upper chest, is practised as for thoracoplasty patients. Usually about 4 periods are sufficient. Because in such cases the tuberculosis is frequently more active, precautions should be taken not to tire the patient. Localized arm movements can be practised once or twice with the opposite arm. Post operative treatment is not usually necessary for more than 2 weeks.

When the patient is up, the arm is supported by a sling. The arm is used for the graft. The deltoid can be practised to prevent atrophy, but neither stretching nor contraction of the muscle used for the graft is allowed for the first 14 days. Movements are then gradually increased. The arm is either kept in a sling or lightly bound to the side, to

## BIBLIOGRAPHY

prevent the patient from using it inadvertently    Postural corrections and breathing exercises are used

### *Posture*

In the physical rehabilitation of 'surgical chest' patients, co operation between nurses and physiotherapist is necessary to maintain good posture in bed and to ensure early coughing to eliminate post anaesthetic secretions and thus prevent collapse of the lower lobes

## BIBLIOGRAPHY

- Fink H D (1945) *Release from Nervous Tension*    London    Allen and Unwin  
Foster Carter A F (1943)    An Outline of Bronchial Anatomy    *Proc R Soc Med* 36, 451  
Jacobson E (1938) *Progressive Relaxation* 2nd ed    Chicago    University of Chicago Press  
Nelson H P (1934)    Postural Drainage of the Lungs    *Brit med J* 2, 251





# INDEX TO BOTH VOLUMES

## A

Abdomen in pneumonia 80 98  
Abortive focus tuberculous 300

### Abscess

cerebral  
bronchopulmonary 149 171

chemotherapy in 223  
chronicity following rupture 214  
classification 197  
clinical features 213 220  
cold agglutinins in diagnosis 221  
convalescence following rupture 214  
cough in 213  
course 213  
cysts and radiological appearances 222  
definition 197  
dental sepsis and 242  
diagnosis 221 223  
differential diagnosis 221  
drainage in mycoses 167  
early stage of 213  
embolism causing 198 199 209  
empyema complicating 222 236 18, 37  
*Entamoeba histolytica* in 236  
eosinophilia in 223  
external drainage in 224  
fistula broncho-pleural diagnosis from 221  
fluke infestation causing 223  
foreign bodies and 238 239 242  
Friedländer's bacillus in 97 211 236  
fulminating following rupture 214  
haematoma and 239  
haemoptysis in 214 215

### Abscess—continued

pulmonary—continued  
healing of 206  
hyper-acute following rupture 214  
idiopathic 198  
infarction and 238 240-241  
influenza simulating, 221  
inhalational pneumonia and 123  
injuries and 239  
investigations in 221  
leucopenia in diagnosis, 221  
lobectomy specimen 201  
mediastinitis caused by 354  
non specific infection 213  
onset 213  
pain in 215  
pathogenesis 198  
pathology 202 211  
percussion therapy 224  
physiotherapy in 223 401  
pleura appearances of 209  
pneumonia and  
diagnosis from 221  
non specific suppurative and 202 239  
pneumococcal 87 230  
staphylococcal 229 230-234  
streptococcal 230  
pneumonitis simulating 221  
pneumothorax caused by 54  
post atelectatic obstruction and 238  
postural drainage in 223  
pyaemia and 238  
radiographic appearances 203 205 207  
209 210 215 218 219 220 222 226-  
227 229  
resection in 225  
residual cyst 218  
resolution 206  
results of treatment 228  
rupture and 202, 214  
signs 215  
specific types of 228 243  
specimen appearances 212  
spread 206 214  
sputum in, 215 221  
*Staphylococcus aureus* in 211  
suppuration management of 227  
surgical treatment 224 227  
symptoms 215  
tomographic appearances, 201 205  
treatment 223-228  
urticaria in 223  
vascular emboli and 198 238  
virus infection simulating 221  
sub-diaphragmatic 380 384  
sub-phrenic empyema and 37, 57  
ACTH  
asthma in 94-95

Figures in bold type refer to Volume Two

# INDEX

ACTH—continued  
beryllium pneumonia in 147

*Actinomyces* 163

*israeli* 168 174

Actinomycosis 163 167

abscess pulmonary and 211 234

carious teeth and 163

causative organisms 163

clinical features 164

culture 167

empyema caused by 18 20 38

necessitatis 39

incidence 163

laboratory diagnosis 164

mediastinitis caused by 355

new bone formation in 164

penicillin in 167

prognosis 167

radiographic appearances 165 166

treatment 167

Adenitis hilar in children 284-289

Adenocarcinoma pulmonary 310

Adenomatosis 330

Adenoma

bronchial 331 335

pulmonary abscess and 237

sebaceous in tuberous sclerosis 201

Adrenal glands and pneumococcal pneumonia 74

Adrenaline

asthma in 92

respiration effect on 23

Agglutination tests in pulmonary tuberculosis 373

Agglutinins

abscess pulmonary in 221

pneumonia pneumococcal in 69

Agranulocytosis thiosemicarbazones causing 450

Agricultural workers actinomycosis and 163

Albuminuria

bronchiectasis in 163

eosinophilia transient pulmonary in 71

PAS causing 441

pneumococcal pneumonia in 80 89

polyarteritis nodosa in 78

streptomycin causing 436

Alcoholism

pneumococcal pneumonia and 75

tuberculosis and 259

Aleudrin in asthma 93

Alimentary tract pneumococcal pneumonia signs in 80

Allergy

asthma and 83-84 91

pneumococcal pneumonia and 71

eosinophilia transient pulmonary and 65

whooping-cough effect on 159

Altitude whooping-cough disease 285

Aluminum and lung disease 318

Amenorrhoea in bronchogenic tuberculosis 318

Aminophylline in asthma 93

Ammonia pneumonia from 141 142

Ammonia pneumonia 225-227

Amoebiasis thoracic 225-227

Amyloid disease

bronchiectasis and 163 173

empyema complicated by 47

Anaemia

histoplasmosis in 185

hypochromic pulmonary tuberculosis and 373

thiosemicarbazones causing 450

Anaesthesia 388 394

abscess pulmonary danger of embolism in 209 213

aggravation of disease by 391

atelectatic dangers of 391

pneumonia and 63 123

pneumoperitoneum artificial in 398

pneumothorax and 392 392

secretion spread and 389

surgical emergencies in 388

technique choice of 392

thoracoplasty in 404

Anatomy

broncho pulmonary 1 23

lobar 3 17

Angiofibroma diaphragmatic 384

Angiography

hamartoma in 339

mediastinal growths in 359

Angiomatosis 217

Aniline pneumonia from 141

Animal protective antibodies pneumococcal pneumonia and 69

Ankylostoma 224

*braziensis* in transient pulmonary eosinophilia 65

Ankylostomiasis schistosomiasis associated with 220

Anoxaemia arterial in pulmonary oedema 129

Anthrax 277

Anthrax broncho pulmonary 101

Antibiotics (see also specific preparations)

antibiotic pulmonary in 213

abscess pulmonary in 213

bronchitis acute catarrhal tracheo- in 48

carcinoma pulmonary in 330

empyema tuberculous in 364

monilia proliferation of flora and 169

*Mycobacterium tuberculosis* sensitivity to 368

rickettsial pneumonia in 122

Anticoagulants 371 244

Antigens

coccidioidomycosis and 184

histoplasmin 185 188

tuberculin 252 254

Antihistamines

asthma in 93

bronchitis chronic in 59

farmer's lung in 180

polyarteritis nodosa in 80

Antimony in schistosomiasis 219 221

Antipyretics in pulmonary tuberculosis 383

Antispasmodics

atelectasis in 144

bronchitis chronic in 58

whooping cough in 159

Figures in bold type refer to Volume Two

- Animal inoculation in pulmonary tuberculosis 368
- Aorta  
   dissecting aneurysm of causing haemorrhax 49  
   hypoplasia of in schistosomiasis 219
- Aphasia in whooping cough 158
- Appendicitis stimulating pneumococcal pneumonia 90
- Argyrosiderosis 283
- Arsenicals  
   eosinophilia in 65, 72  
   pulmonary syphilis in 237
- Arteriosclerosis in pneumococcal pneumonia 80
- Arterius giant cell 79 80
- Arthritis  
   bronchiectasis complicated by 173  
   pneumococcal pneumonia and 74 89
- Asbestos lung disease and 263 267
- Ascaris* 224  
   transient pulmonary eosinophilia in 65
- Ascites haematogenous tuberculosis accompanying, 301
- Aspergillosis 175 177
- Aspergillus fumigatus* 168 175 179
- Asphyxiants causing pneumonia 141
- Aspiration  
   diagnostic  
     carcinoma pulmonary in 324  
     empyema in 24 42  
     sarcoidosis in 187  
     sub-diaphragmatic hernia in 383  
   therapeutic  
     abscess pulmonary sequelae of 211  
     artificial pneumothorax haemorrhagic in 402  
     bronchoscopic  
       pneumococcal pneumonia in 364  
       status asthmaticus in 95  
     empyema in  
       acute 26-28  
       tuberculous 364  
     radiography as adjuvant 417  
     transpleural in hydatid cysts 233
- Assmann focus 301
- Asthma 82 99  
   adrenaline in, 92  
   age incidence 85  
   alcudrine in 93  
   allergy in 83 84  
   allergen avoidance 91  
   allied conditions 87  
   aminophylline in 93  
   anaesthesia in 392  
   angioneurotic oedema and 88  
   antihistamines in 93  
   atelectasis and 141  
   bronchiectasis complicating 89  
   bronchiolar spasm relief of 92  
   bronchitis and 88  
   acute catarrhal tracheo- 47  
   chronic 55 57
- Asthma—continued  
   bronchitis and—continued  
     farmer's lung simulating 177  
     kymographic reading in 32  
   cardiac 90 125  
   cardophyllin in 93  
   clinical features 86-89  
   complications of 88  
   conjunctivitis seasonal and 88  
   cor pulmonale complicating 89  
   definition 82  
   description of attack 86  
   desensitization specific 96  
   diagnosis 90 91  
   differential diagnosis 90  
   eczema and 88  
   emphysema in 107  
     chronic hypertrophic 116  
     complicating 88  
   environment and 85 92  
   eosinophilia and 65 91  
   ephedrine 93  
   exercises breathing 93  
   exertion and 86  
   extrinsic causes 84  
   food idiosyncrasies in 88  
   general protective measures 91  
   hay fever and 87  
   heredity and 85  
   hypnotics in 93  
   incidence 84  
   inhalants in 93  
   intelligence and 85  
   intrinsic causes 84  
   isupren in 93  
   massive collapse complicating 89  
   mechanistic causes 84  
   migraine and 88  
   miners 277  
   morphine and 93  
   nationality and 84  
   occupation and 92  
   periodicity of attacks 87  
   physiotherapy in 395  
   platinum 252 254  
   pneumothorax spontaneous complicating 89  
   precipitating causes of attack 86  
   prognosis 89  
   prurigo and 88  
   psychological factors in 84  
   race incidence 84  
   rhinitis allergic and 87  
   rhinorrhoea seasonal and 87  
   sex incidence 85  
   skin tests in diagnosis 90  
   social status and 85  
   specific treatment 95  
   status asthmaticus, treatment 94 96  
   symptomatic treatment 92  
   treatment 91 98  
   tuberculosis pulmonary complicating, 89  
   urticaria in 88

Figures in bold type refer to Volume Two

## INDEX

- Asthma**—continued  
vital capacity in 88  
**Atelectasis** 137 155  
abscess pulmonary in 202 218 238  
antispasmodics in 144  
asthma and 141  
bronchial occlusion and 199  
bronchitis and 44  
bronchograms in 152  
bronchoscopy in 154  
carcinoma and  
bronchial 140  
pulmonary 322 323  
causation 140  
classification 140  
complications 144 148  
definition 137  
embolism diagnosis from 241  
emphysematous bulla ruptured complicating 148  
fate of lung 154 155  
heart displacement in 139  
histology 137  
honey combing in 150  
hydatid cysts causing 237  
lobar 140  
lobular 140  
massive 140  
mucus plug in 142 143  
penicillin in 144  
physiology of 138  
physiotherapy in 401  
pneumothorax and  
artificial 394  
spontaneous 148  
post-operative 239 148 154  
radiographic appearances 144  
segmental 140  
signs 142  
symptoms 141  
thoracoplasty complicating 410 414  
tracheo bronchitis tuberculous causing 140  
treatment 144  
tuberculosis and 390 144  
whooping cough and 156 159 141  
**Aureomycin**  
abscess, pulmonary in 223  
amoebic infections in 227  
pneumonia in 153  
pneumococcal in 93  
psittacosis 113  
staphylococcal 99  
whooping cough in 159  
**B**  
**Bacillus**  
anthracis pneumonia of 101  
Friedlander's (see Friedlander's *Bacillus*)  
role in haemorrhax 49  
**BAL**, in vanadium pneumonia 149  
**Barbitosis**, 280  
**Barbiturates** causing pneumonia 141  
**Basophilic** causing pneumococcal pneumonia 75  
**B.C.G.** 270  
hypersensitivity 292  
infant inoculation 382  
intradermal injection 292  
multipuncture technique 291  
preliminary testing 292  
reaction to 292  
scarification in 291  
tuberculosis in children in 291  
Belladonna in whooping cough 159  
Benzene causing pneumonia 141  
**Blastomycosis** 188-190  
causative organism 188  
European 171  
mode of infection, 189  
North American 189  
radiographic appearances 189  
South American 190  
treatment 190  
**Blood**  
carbon dioxide transport of 26  
carcinoma metastases and 312  
clotting time streptomycin in 441  
disorders pulmonary manifestations 192  
menstruation 37  
**Blood sedimentation rate**  
pulmonary embolism in 240  
sclerosis in 272  
streptomycin action on 435  
transfusion artificial pneumothorax haemorrhage of in 402  
tuberculosis and  
pulmonary 373  
spread by 294 313 (see also Tuberculosis  
Haematogenous)  
vanadium in demonstration of 149  
**Boeck's sarcoid** 185  
**Bordet-Gengou** culture in whooping cough 158  
**Bornholm disease** 342 343  
**Bradycardia** in pulmonary abscess 221  
**B**  
**Bacillus**  
anthracis pneumonia of 101  
Friedlander's (see Friedlander's *Bacillus*)  
role in haemorrhax 49  
**BAL**, in vanadium pneumonia 149  
**Barbitosis**, 280  
**Barbiturates** causing pneumonia 141  
**Basophilic** causing pneumococcal pneumonia 75

**Bronchial tree—continued**

- nomenclature 3
- respiration and 23
- Bronchiectasis 156-184**
  - abscess cerebral complicating 168, 171
  - aetiology 158
  - albuminuria in 163
  - amyloid disease in 163, 173
  - arthritis infective complicating 173
  - asthma complicated by 89
  - auscultation findings 163
  - bacteriology sputum 162
  - bronchitis
    - acute 163
    - chronic 163
  - catarrhal 163
  - children 160
    - pulmonary complicated by 311
  - children course and 168
  - Chloromycetin in 181
  - clubbing of fingers in 163
  - complications 171 173
  - course 168 171
  - cylindrical 167
  - diagnosis 162 168
  - embolism and 241
  - empyema acute and 18 37
  - experimental work in 158
  - foreign bodies causing 160
  - fusiform 167
  - hepatomegaly in 163
  - histology 162
  - history 156
  - hydatid cysts causing 231
  - lobectomy in 183
  - measles causation and 159
  - mediastinitis caused by 344
  - mediastinal displacement caused by 340
  - penicillin in 175
  - percussion therapy 175
  - pneumectomy in 183
  - pneumonia and
    - lipoid, 150
    - lobar causing 160
    - measles and 111
    - pneumococcal complicating 88
  - postural drainage in 173, 182
  - radiographic appearances 46 47 163, 164, 161, 169 171, 172, 174 181
  - saccular 167
  - signs 162
  - sputum appearances in 162
  - streptomycin in 181
  - sulphonamides in 175
  - surgery in 183
  - symptoms 162
  - tracheo-bronchitis tuberculous and 162
  - tracheotomy and 160
  - treatment 173 183

**Bronchiectasis—continued**

- tuberculosis and
  - children in complicating 284
- fibroid 346
- glandular causation and 159
- haemoptysis and 319
- whooping cough and
  - causation and 159
  - complicating 158 159
- Bronchiolitis**
  - diffuse honey comb lungs caused by 200
  - fibrosa obliterans 53
- Bronchitis**
  - acute tracheo- 43 54
  - catarrhal 43-49
    - aetiology 43
    - antibiotics in 48
    - asthma and 47
    - atelectasis from 44
    - bacteriology 43
    - bronchiectasis complicating 45
    - chronic bronchitis complicating 45
    - clinical picture 44
    - complications 45
    - convalescence 49
    - differential diagnosis 47
    - diphtheria and 52
    - general treatment 48
    - heart failure complicating 45 48
    - mitral stenosis and 43
    - pathology 44
    - pleurisy complicating 45
    - pneumonia and 44 45
    - post-operative 47
    - radiographic appearances 46 47
    - renal disease and 43
    - rheumatic heart disease complicating 45
    - signs 44
    - specific treatment 48
    - sulphonamides in 49
    - symptomatic drug treatment 48
    - symptoms 44
    - treatment 48-49
    - tuberculosis complicating 45
  - asthma complicated by 88
  - byssinosis in 258
  - catarrhal
    - dust inhalation and 52 53
    - enteric fevers in 52
    - fibrinous 50-51
    - gas poisoning in 53 54
    - infants in 49
    - influenza in 51
    - measles in 51
    - physiotherapy in 400
    - plastic 50-51
    - smallpox and 52
    - suppurative 50
    - whooping cough in 51, 156
  - chronic 55 59
    - acute tracheal catarrhal complicating 45
    - aetiology 56
    - antihistamines in 59

- Bronchitis—continued**  
*chronic—continued*  
 antispasmodics in 58  
 asthma and 55 57  
 bronchiectasis diagnosis from 58  
 bronchographic appearances 57  
 clinical picture 55  
 clubbing of fingers in 58  
 differential diagnosis 57  
 emphysema chronic hypertrophic and 107, 116  
 exercise breathing in 59  
 farmer's lung simulating 177  
 first phase, 55  
 pathology 56  
 physical signs 56  
 prognosis 58  
 radiography in 57  
 smoking in 56 58  
 temperature and 56 58  
 treatment 58  
 tuberculosis diagnosis from 58  
 fenestriatum 254  
 gases and 142  
 manganese causing 148  
 physiotherapy and 395  
 pneumonia aspiration and 64  
 post-operative 47  
 purulent influenzal pneumonia and 109  
 schistosomiasis complicated by 219  
 tuberculosis and  
   bronchogenic 343  
   chemotherapy 444  
   fibroid 346  
   pulmonary complicating 378  
   vanadium pneumonia in 148  
 xanthomatosis in 203  
**Bronchodilators in emphysema, 117**  
**Bronchogenic tuberculosis 314 354**  
**Bronchographic appearances**  
 atelectasis 152  
 bronchiectasis 165-166  
 bronchitis chronic 57  
 bronchus  
   displacement of 20  
   obstruction 9  
     children in 287  
 granuloma diffuse cystic 214  
 lung abscess 208-209  
 normal 6  
 supernumerary apical branch 21  
**Broncho moniliasis 170**  
**Broncho pneumonia**  
 abscess pulmonary and 216 221  
 children in 280-284  
 embolism diagnosis from 241  
 farmer's lung simulating 177  
 fluke infestation causing 223  
 measles and, 110  
 reticulosis in 194  
 tuberculous 106-107  
 xanthomatosis in 203
- Bronchoscopy**  
 abscess pulmonary in 221 224 237  
 atelectasis in 144  
 bronchiectasis in 67  
 carcinoma pulmonary in 320  
 hydatid cysts in 233  
 indications for, 391  
 mediastinal growths in 358  
 pneumococcal pneumonia in 71  
 sarcoma pulmonary in 330  
 tuberculosis and  
   children in 291  
   tracheo bronchitis in 346  
 whooping cough in 159  
**Bronchospasm**  
 " 731
- B**  
 adenoma of 331-335  
 anatomy 1 23  
 biopsy in pulmonary tuberculosis 373  
 carcinoma of  
   abscess pulmonary and 237  
   atelectasis caused by 140  
   bronchiectasis caused by 160  
 celloidin cast of 11  
 diagrammatic representation 12  
 displaced 19 20  
 fibrosarcoma of 330  
 mucosa of section 374  
 supernumerary 22  
 ulcer tuberculous lesions and 314  
**Brucella tularensis pneumonia of 102**  
**Brucellosis**  
 eosinophilia transient pulmonary in 65  
 polyarthritis nodosa diagnosis from 80  
**Byssinosis 255-258**
- C**  
 Cadmium causing pneumonia 149  
 Caisson disease oxygen administration and 40  
 Calciferol in sarcoidosis 191  
 Camelote formation 231  
**Candida albicans 167 168 174**  
**Carbon dioxide**  
 administration problems 40-41  
 blood content, in pneumococcal pneumonia 71  
 " 71  
 " 41

Causes of

- bronchial
  - bronchiectasis caused by 160
  - empyema complicating 37
- pulmonary 258 261, 306-330
  - abscess pulmonary complicating 311
  - adenocarcinoma 310
  - aetiology 306
  - antibiotics 330
  - asbestosis and 261
  - aspiration biopsy 324
  - avascular-cell 330
  - bronchiectasis complicating 311
  - bronchogenic metastasis 312
  - bronchography 320
  - bronchoscopy 320
  - chromate working and 261
  - columnar-cell 308
  - cordotomy in 330
  - cough in 316
  - diagnosis 312
  - diaphragmatic infiltration 312
  - differential diagnosis 324
  - direct extension of 312
  - effusion pleural in 330
  - empyema acute causing 18
  - haematogenous metastasis 312
  - haemoptysis in 316
  - haemothorax in 51
  - histology 308
  - incidence 306
  - inoperable diagnosis of 326
  - lymphatic spread 312
  - macroscopic appearance 310
  - mediastinal displacement caused by 350
  - metastasis 312
  - neurectomy intercostal in 329
  - oat-cell 308
  - operability of 325
  - palliation 329
  - pathology 308
  - penicillin in 330
  - physical examination 318
  - physiotherapy in 409
  - pneumonitis suppurative complicating 311
  - pneumothorax caused by 54
  - radiographic appearances 318
  - radiotherapy in 329
  - resection results 328
  - spread 311
  - sputum examination 320
  - squamous-cell 308
  - symptomatology 316
  - tobacco smoking and 261, 307
  - tomography 324
  - treatment 325 330
- Carcinomatous lymphangiosis 312
- Cardophyllin in asthma 93
- Casoni reaction
  - hydatid cysts in 229
  - pulmonary abscess in 223
- Cavitation tuberculous
  - abscess pulmonary and 222
  - drainage of 415-419
- Cellulitis pulmonary pneumococcal causing 70
- Cerebrospinal fluid examination in pulmonary tuberculosis 371
- Cerebrum haemorrhage of in whooping cough 158
- Chemotherapy
  - bronchitis tuberculous in 44
  - laryngitis tuberculous in 445
  - palliative use 443
  - resection in 446
  - sinus tuberculous in 447
  - thoracoplasty in 447
  - tuberculosis in 431-456
  - unstable cases in 444
- Chest wall
  - abscess actinomycosis causing 164
  - tumours 344 349
- Chicken pox pneumonia of 113
- Children
  - atelectasis in 138
  - bronchiectasis in 167
  - bronchitis chronic in 55
  - erythema nodosum pleural tuberculosis and 360
  - haemosiderosis in 248
  - leukaemias in 194
  - pneumonia in
    - staphylococcal 99 232
    - pneumococcal 84 85
  - pyopneumothorax in 232
  - tuberculosis in 270 293
    - haematogenous 295
    - pleural 356
    - pulmonary streptomycin in 432
  - tuberculous sclerosis in 201
- China clay and lung disease 261
- Chloramphenicol (see also Chloromycetin)
  - pneumonia in 133
  - influenza 110
  - pneumococcal 93
  - psittacotic 113
- Chlorides blood content in pneumococcal pneumonia 81
- Chlorine
  - bronchitis acute tracheo-bronchitis from 53
  - pneumonia and 142
- Chloromycetin (see also Chloramphenicol)
  - abscess pulmonary in 223
  - bronchiectasis in 181
  - sub-diaphragmatic hernia in 384
  - whooping cough in 158
- Chloropicrin
  - bronchitis acute tracheo-bronchitis from 53
  - pneumonia from 141
- Chondromas
  - chest wall of 345

Figures in bold type refer to Volume Two



Chondromas—continued  
diaphragmatic 384  
mediastinal 373

emphysema chronic hypertrophic in 108  
irradiation pneumonia in 151  
oxygen saturation reduced and 31  
schistosomiasis in 220

Coal miners aspergillosis of 175  
Cocaine respiration effect on 23  
*Coccidio des immitis* 168 179 180  
Coccidioidin 184

Coccidioidomycosis 180-184  
blood test in 184  
causative organism 180  
diagnosis 184  
mode of infection 180  
primary 180

pulmonary (see Atelectasis)  
therapy 387 396 (see also *Pneumothorax*)

Complement fixation test  
influenzal pneumonia in 110  
pneumococcal pneumonia and 69  
rickettsial pneumonia in 122  
tuberculosis pulmonary in 373

Conjunctiva haemorrhagic in whooping cough 156

Conjunctivitis  
asthma and 88  
pneumococcal pneumonia complicating 89  
thiosemicarbazones causing 450

Conteben 449

Convulsions in whooping cough 158

Cordotomy in pulmonary carcinoma 330

Cor pulmonale  
asthma complicated by 89  
embolism pulmonary causing 239

Cortisone  
asthma in 95  
beryllium pneumonia in 147

Coryza in bronchitis 44 51

Cough  
carcinoma pulmonary in 316  
lung abscess prevention and 200  
pneumonia in 77 131  
tracheo bronchitis in 320 348  
tuberculosis in 320 383

Cylindruria in pneumococcal pneumonia 89  
Cystine streptomycin and inhibition by 433  
Cysts

anaesthesia in 388  
bronchial pneumothorax diagnosis from 57  
hydatid 229 234  
abscess pulmonary and 222 223  
aetiology 229  
aspiration of 233  
bronchoscopy in 233  
radiographic appearances 230 232 233  
treatment 231-234  
mediastinal 358 374  
enterogenous 370 371  
pericardial 367 369  
physiotherapy in 404  
pulmonary 294 305  
acquired 298  
congenital 294 298

## D

D D S in tuberculosis 447 448

Decortication

chronic empyema in 44  
haemothorax in 53  
physiotherapy and 404

Delirium tremens in pneumonia 77 131

Decompression

sickness oxygen administration and 40  
treatment in whooping cough 159

Dental sepsis and pulmonary abscess 242

Dermatitis caused by streptomycin 436

Dermatomyositis histological similarity to  
periarteritis nodosa 75

Dermoids mediastinal 363

Desoxycorticosterone acetate in subacute pulmonary oedema 135

Devil's grip 342

Diabetes

carcinomatous invasion of 312

eventration 375 376

hernia of 376-380

paralysis of 375

pulmonary tuberculosis in 396 397

paresis of 375

pleural effusion and discharges of 13

pneumonia pneumococcal in 71

trichiniasis 376

tumours of 384

Diarrhoea in pulmonary tuberculosis 384

Diazene in tuberculosis, 448

- Diatomaceous earth in lung disease 268  
 Dicumarol  
   irradiation pneumonia in 152  
   pulmonary embolism in 244  
 Diet  
   diabetic in tuberculous states 382  
   pneumonia in 130  
 Digitalis  
   embolism in 245  
   pulmonary oedema heart failure and 131  
 Dihydrostreptomycin 438  
 Diplococcus pneumoniae 67  
 D stomatosis 223 224  
 Drainage  
   cavity 415-419  
   empyema acute in 28 33  
   external pulmonary abscess in 224  
   lymphatic of the lungs 24  
   ostural 13 402-403  
   abscess pulmonary in 228  
   bronchiectasis in 173 182  
   physiotherapy and 405  
   status asthmaticus in 95  
   suppuration, pulmonary in 228  
   tuberculosis pulmonary in 387  
   whooping cough in 159  
 Dust (see also Occupational Diseases)  
   bronchitis and  
     acute tracheo- 52 53  
     chronic 141  
   coccidioidomycosis, in 180  
   empysema and 141  
   farmer's lung in 177  
   pneumonia and 141  
   tuberculosis and 141  
 Dyspepsia in pulmonary tuberculosis 384  
 Dysphagia  
   inhalational pneumonia in 123  
 Mediastinitis in, 355  
 Dyspnoea  
   empysema chronic hypertrophic in 106  
   farmer's lung in 177  
   paroxysmal nocturnal cardiac 125  
   pneumococcal pneumonia in 77  
   tuberculosis in 321
- E**  
 Eczema asthma and 88  
 Effusions  
   abscess pulmonary in following treatment  
     209  
   blood stained in embolism 241  
   pericardial  
     haematogenous tuberculosis accompany  
       ing 301  
     pneumococcal pneumonia complicating 88  
   peritoneal artificial pneumoperitoneum com  
     plicating 399  
   pleural 2 17  
   amount  
     bilateral radiographic appearances 357  
     carcinoma pulmonary in 330
- Effusions—cont nued  
   cardiac disease and 14  
   chyliform 15  
   chylous 15  
   extrapulmonary abscess in 358  
   hepatic disease and 13  
   mediastinal displacement caused by 350  
   new growths and 14  
   pneumococcal pneumonia complicating  
     86  
   pneumothorax artificial and 395  
   polyarteritis nodosa in 78  
   post pneumon 6 12 13  
   post primary 6  
   primary chemotherapy in 445  
   pyrexia prognosis and 361  
   renal disease and 14  
   spinal abscess and 358  
   sub-diaphragmatic abscess complicated by  
     13 380  
   tuberculous 2 12, 355  
   children in 289  
   clinical features 360 361  
   complicating 7 362  
   encysted 362  
   haematogenous 300  
   p-aminosalicylic acid in 361  
   streptomycin in 361  
   treatment 361 362  
   tuberculin and streptomycin in 363  
   vocal fremitus in 323
- Electrocardiography  
   embolism in 243  
   empysema in 107  
   schistosomiasis in 220  
 Electro welding pneumonia and 143  
 Embolctomy 245  
 Embolism pulmonary 238 245  
   aetiology 238  
   anaemia treatment in 245  
   anticoagulants in 244  
   atelectasis diagnosis from 241  
   broncho-pneumonia diagnosis from 241  
   cor pulmonale caused by 239  
   diagnosis 243  
   dicumarol in 244  
   digitalis in 245  
   electrocardiography in 243  
   embolctomy in 245  
   heart failure in 240  
   hypernephroma causing 238  
   incidence 238, 239  
   mitral stenosis causing 240  
   pathology 239 240  
   prognosis 245  
   prothrombin levels in 244  
   radiographic appearances 242, 243  
   sarcomatous radiographic appearances 240  
   sedimentation rate in, 240  
   signs 240 243  
   strophanthin in 245  
   symptoms 240 243  
   treatment 243

**Embolism pulmonary—continued**

- Tronexan in 244
- tuberculosis primary causing 238
- vasodilators in 245
- venesection in 245
- vitamin K in 244

**Emetine in thoracic amoebiasis 227**

**Emphysema**

- acute vesicular 119
- anaesthesia in 388
- arteriosclerotic 118
- asthma complicated by 88
- atelectasis production in 138
- atrophic 118
- beryllium pneumonia in 147
- byssinosis in, 258
- chronic hypertrophic 101 118
  - aetiology 114
    - air sacs inadequate ventilation of 104
    - alveolar stress in 115
    - asthma and 107, 116
    - atrophic pulmonary changes 115
    - auscultation in 109
    - breathing capacity maximal 112
    - bronchitis, chronic and, 107, 116, 117
    - bronchodilators in 117
    - carbon dioxide tolerance to 113
    - chest appearance 108
    - clinical picture 108-109
    - clubbing of fingers in 108
    - dead space pathological 105
    - degenerative pulmonary changes 115
    - diagnosis 110-114
    - dyspnoea in 106, 117
    - elasticity loss of 102
    - electrocardiographic changes in 107
    - ephedrine in 117
    - expiratory theory of 114
    - functional pathology 104 107
    - gas mixing 113
    - gross changes in 101
    - haemoptysis complicating 108
    - haemo respiratory exchange 104
    - heart failure in 108
    - heredity and 116
    - history 107
    - hypertension arterial complicating 108
    - increased volume in 105
    - inspiratory theory of 114
    - interstitial emphysema complicating 108
    - katharometer measurement in 113
    - lung volume 113
    - lymphatic circulation 107
    - morbidity anatomy 101 104
    - muscular inadequacy theory of 104
    - nutrition pulmonary impairment of 114
    - palpation in 109
    - papilloedema complicating 108
    - percussion in 109
    - physical signs 108-110
    - pneumoperitoneum in 117

**Emphysema—continued**

- chronic hypertrophic—continued
    - pneumothorax
      - artificial in 117
      - spontaneous 55, 108
    - polycythaemia secondary 108
    - pulmonary circulation 106
    - radiographic appearances 111
    - treatment 117 118
    - vascular factors in 104 105, 107
    - ventilation, unequal distribution 105
    - vital capacity in 112
  - compensatory 119
  - complimentary 119
  - embolism in 241
  - focal 119-120
  - honey comb lungs caused by 207
  - hypertrophic 116-117
    - interstitial acute 120-121
    - localized 119-120
    - lobar obstructive 120
    - mediastinal 350
      - spontaneous pneumothorax complicated by 56
  - non-obstructive 118
  - obstructive 101
    - oxygen administration repercussion of 40
  - physiotherapy in 395
  - schistosomiasis pulmonary complicated by 219
  - senile 118-119
  - silicosis and 272
  - tracheo-bronchitis in 349
  - tuberculosis fibroid and 346
  - vicarious 119
  - whooping cough in 156
- Empyema 18-48**
- acute 18 38
    - abscess and
      - cerebral 38
      - pulmonary 209 236 18 37
      - subphrenic 37
    - actinomycosis and 164 18 20 38
    - aetiology 18
    - anaerobic organisms in 20
    - aspiration in
      - diagnostic 24
      - therapeutic 26-28
    - bacteriology 20
    - bronchiectasis and 18, 37
    - carcinoma and
      - bronchogenic 37
      - pulmonary 18
    - chemotherapy in 26-28
    - chest wall infections complicating 38
    - clinical picture 21
    - Clostridium welchii* in 20
    - complications 38
    - encysted pulmonary abscess and 222
    - exercises in 33
    - fistula and
      - broncho pleural 38
      - formation 21

**Empyema—continued**

**acute—continued**

foreign bodies causing 18  
haemorrhage intercostal complicating 38  
Friedländer's bacillus pneumonia complicating 97

intercostal tube drainage 32  
interlobar pulmonary abscess and 222  
mediastinum position in 23  
metapneumonic 21  
morbid anatomy 20  
non pulmonary derivation 18  
oesophageal lesions causing 20  
osteomyelitis causing 20  
pathology 18

pneumonia and 18  
bacterial non pneumococcal 95  
pneumococcal complicating 85 86  
PAS in 364 446

pneumothorax in  
artificial 325  
spontaneous 363 18  
post-operative care 33-36  
pulmonary derivation 18  
pyaemia complicating of 37  
radiographic appearances 18 22, 24 26  
rib resection in 28-32  
septicaemia complication of 37  
sex incidence 20

signs 23-24  
streptomycin in 364 446  
streptococci in 20  
streptodornase in 36  
streptokinase in 36  
symptoms 23 24  
synpneumonic 21  
transpleural surgery and 20  
treatment 364 26 38  
wounds causing 18

chronic 38-47  
aetiology 38 39  
amyloid disease complicating 47  
aspirat on diagnostic 42  
brain abscess complicating 47  
cavity estimation 43  
clinical picture 41-43  
complications 47  
decortication 44  
fistula broncho pleural causing 41  
latent 41

nephritis complicating 47  
osteoarthritis complicating 47  
pathology 38  
persistent 41  
plastic chest wall operations 45  
radiographic appearances 40, 42, 46  
re-drainage 43  
Robert's flap operations 45  
Schede's operation 45  
signs 41  
treatment 43-47

**Empyema—continued**

diaphragmatic 21  
interlobar 21  
necrotic 39  
physiotherapy in, 404 412  
pleural calcification in 63  
staphylococcal radiographic appearance 29  
sub-diaphragmatic abscess complicated by 380

thoracic non tuberculous 18  
tuberculous investigation for 43  
Encephalitis caused by torulosis 173  
Encephalopathy hypertensive in polyarteritis nodosa 78

Endometriomas  
diaphragmatic 384  
haemothorax caused by 49  
chest wall of 344

*Entamoeba histolytica* 225  
eosinophil a transient pulmonary in 65  
pulmonary abscess in 236

Enteric fevers  
bronchitis acute tracheal and 52  
pneumonia in 103

Enteritis tuberculous thiosemicarbazones in 450

**Eosinophilia**

transient pulmonary 65 73  
aetiology 65-66  
albuminuria in 71  
asthma in 65  
asthmatics and 66  
brucellosis in 65  
clinical features 66 71  
definition 65  
drug reactions and 65 66  
food sensitivity and 65  
granulomatosis and 66  
hepatomegaly in 71  
history 65  
morbid anatomy 66  
plant sensitivity and 65  
pleural effusion in 71  
pollens causing 66  
polyarteritis nodosa and 65  
radiographic appearance 67-69  
rheumatic fever in 65  
serum injections and 65 66  
splenomegaly in 71  
thiouracil causing 65  
tropical 65 71

Ephedrine  
asthma in, 93  
emphysema chronic hypertrophic in 117  
respiratory effect of 23

Epilepsy in tuberculous sclerosis 201

Epinephrine in gasping 145

Epistaxis in whooping cough, 156

Epituberculosis, pneumonia and 103 104

# INDEX

Erythema  
*multiforme*  
 coccidioidomycosis and 181  
 exudativum pneumonia of 113  
 nodosum  
 coccidioidomycosis and 181  
 tuberculosis and  
 pleural 360  
 pulmonary in children 290  
 Erythrocyte sedimentation rate  
 coccidioidomycosis in, 184  
 moniliasis in 171  
 mycoses in 167  
 tracheo bronchitis non obstructive 348  
 tuberculosis pulmonary in 374  
 Escharotics in cavity drainage 413  
 Erythema in whooping cough 159  
 Eventration diaphragmatic 375 376  
 Ewing's tumour of ribs 349  
 Exercise (see also Physiotherapy)  
 breathing  
 asthma in 93  
 chronic bronchitis in 59  
 emphysema in 117  
 status asthmaticus in 95  
 — — — 241

## F

Faeces, examination of in pulmonary tuberculosis 370  
 Fallot's tetralogy reduced oxygen saturation and 31  
 Farmer's lung 177 180 254  
 Farm workers aspergillosis of 175  
 Fasciola hepatica in transient pulmonary eosinophilia 65  
 Fever  
 streptomycin causing 435  
 tuberculosis pulmonary treatment of 383  
 Fibroma  
 mediastinal 373  
 sub-ungual tuberculous sclerosis in 201  
 Fibromyosarcoma diaphragmatic, 384  
 Fibrosarcoma bronchial 330  
 Fibrosis  
 bacterial non pneumococcal pneumonia and 95  
 lipo pneumonia in 150  
 Filar a bancrofti and lung disease 225  
 Fistula broncho pleural 396 38 42  
 Fluke infestations of the lungs 223-224  
 Fluoroscopy  
 diaphragmatic eventration in 376  
 mediastinal displacement in 350  
 pleurisy dry in 2  
 Foreign bodies  
 bronchiectasis caused by 160  
 pulmonary abscess caused by 238 239

Fouadin in fluke infestation 221 224  
 French chalk in lung disease 262  
 Friedländer's bacillus  
 lung abscess in 211 236  
 pneumonia of 96 97  
 Fungus diseases (see also specific infections)  
 aspergillosis 175 177  
 actinomycosis 163 167  
 blastomycosis 188 190  
 classification on 162  
 coetid oodomycosis 180-184  
 dimorphic fungi 180-192  
 farmer's lung 177  
 filamentous like organisms, 163 167  
 histoplasmosis 185 188  
 laboratory investigations 162 163  
 moniliasis 167 171  
 nocardiosis 163 167  
 sarcoidosis diagnosis from, 187  
 sporotrichosis 190 192  
 torulosis 171 175  
 treatment 163  
 ubiquitous fungi 175 180  
 yeast like organisms 167 173

## G

Gallop ng consumption 343  
 Ganglioneuromas, mediastinal 360  
 Gases lung irritant 142 146  
 Gassing  
 bronchitis acute tracheal from 53  
 pneumonia from 141  
 treatment 144  
 — — — effect on following radiograph c  
 311 315  
 Gold salts in pulmonary tuberculosis 383  
 Grain handlers aspergillosis of 175  
 Granuloma  
 beryllium 146  
 diffuse cystic causing honey combing 207  
 eosinophilic of bone 203  
 Granulomatosis  
 beryllium diagnosis from sarcoidosis 190  
 pulmonary causing honey combing 207  
 transient pulmonary eosinophilia in 66  
 Graphite and lung disease 276-277

## H

Haemangio endothelioma diaphragmatic 384  
 Haematogenous tuberculosis 294 313  
 Haematoma and pulmonary abscess 239  
 Haematuria  
 polyarteritis nodosa in 78  
 pneumococcal pneumonia in 89

F figures in bold type refer to Volume Two

- Haemophilus influenzae*  
 bronchitis in 43  
 pneumonia in 43 99 109  
 pertussis in whooping-cough 154
- Haemopneumothorax* 49, 54
- Haemoptysis*  
 abscess pulmonary in 214 215  
 carcinoma in  
 bronchial 237  
 pulmonary 316  
 emphysema chronic hypertrophic in 108  
 pneumonia in  
 pneumococcal 77  
 tuberculous lobar 105  
 vanadium 148  
 tuberculosis in 319 383
- Haemosiderosis* 248 251  
 idiopathic 248 250  
 mitral stenosis and 250 251  
 schistosomiasis diagnosis from 221
- Haemothorax*, 49 53  
 decortication in 53  
 empyematous streptokinase and streptodornase in 36  
 radiographic appearances 50 51  
 surgical treatment 53  
 symptomatology 51 52  
 thoracotomy 53  
 traumatic 49
- Halogens pneumonia* from 142
- Hamartoma* 335 340  
 chondromatous 335 337  
 vascular 337 340
- Hand-Schüller-Christian disease* 203
- Hay fever and asthma* 87
- Headache in pneumococcal pneumonia* 77
- Heart*  
 apex beat displacement in tuberculosis 322  
 auricular fibrillation in pneumococcal pneumonia 79  
 dilatation right sided in whooping-cough 156  
 disease  
 asthma complicated by 89  
 biliary 216 222  
 congenital diagnosis from schistosomiasis 221  
 embolism in 240  
 honey comb lungs and 200  
 pleural effusion and 14  
 displacement in atelectasis 91 139  
 failure  
 bronchitis acute tracheal in 45 48  
 emphysema chronic hypertrophic in 108  
 honey comb lungs and 214  
 irradiation pneumonia and 151  
 oedema acute pulmonary and 125 131  
 pneumothorax artificial and 391
- Heart—cont. nued*  
 myomas of in tuberous sclerosis 201  
 pneumonia in 79 131  
 xanthomatosis in 203  
 polyarteritis nodosa symptoms in 78  
 Hiccup and syndrome in sarcoidosis 185  
 Hepatosplenomegaly Egyptian 216
- Heredity*  
 asthma and 85  
 emphysema and 116
- Hernia diaphragmatic* 376 380
- Herpes in pneumococcal pneumonia* 78
- Hexamine in gassing* 145
- Hist test in influenza pneumonia* 110
- Histoplasma capsulatum* 185
- Histoplasmosis* 185 188  
 biopsy in 188  
 causative organism 185  
 clinical features 186  
 haematogenous tuberculosis and 300  
 incidence 186  
 laboratory diagnosis 188  
 mode of infection 185  
 prognosis 188  
 radiographic appearances 186 187  
 treatment 188
- Hodgkin's disease* 172 51 367
- Honey comb lungs* 200 215  
 aetiology 203  
 bronchiolitis diffuse causing 200  
 bronchography in 214  
 clinical features 209 215  
 course 209  
 developmental group 200  
 differential diagnosis 215  
 emphysema causing 207  
 granuloma diffuse cystic causing 207  
 granulomatosis causing 200  
 heart disease and 200 214  
 infantile biliary cirrhosis of 207  
 lymphangitis diagnosis from 215  
 pathogenesis 200  
 pleurodesis artificial in 215  
 pneumoconiosis diagnosis from 215  
 pneumothorax spontaneous complicating 209  
 radiographic appearances 201 214  
 radiotherapy 215  
 sarcoidosis diagnosis from 215  
 tomograms in 214  
 treatment 214  
 tuberous sclerosis 201 205

# INDEX

Hypertension  
arterial in haemothorax 51  
eosinophilia transient pulmonary in 71  
oedema subacute and 132 134  
malignant polyarteritis nodosa diagnosis from 80

## I

Immunization  
natural pneumococcal pneumonia and 75  
whooping-cough in 160  
Ind gestion bronchogenic tuberculosis and 318  
Industry tuberculosis in 258  
Infants (*see also* Children)  
biliary cirrhosis of 207  
pneumonia in  
lipoid 149  
measles and 110  
staphylococcal empyema caused by 18  
pneumothorax spontaneous in 55  
syphilis pulmonary in 235  
whooping cough 154 161

Infarction (*see also* Embolism)  
abscess pulmonary and 198 238  
myocardial in polyarteritis nodosa 78

Influenza  
abscess pulmonary diagnosis from 221  
bronchitis acute tracheal and 51  
pneumonia and 62 107 110  
Insomnia in pneumonia 77  
relief of 130  
Insulin in pulmonary tuberculosis 384  
*Iodamoeba butschlii* 225  
Iodides

aspergillosis in 177  
blastomycosis in, 190  
farmer's lung in 180  
moniliasis in 171  
nocardiosis in 167  
sporotrichosis in 192  
syphilis pulmonary in 237  
torulosis in 173  
*Iso* nicotinic acid hydrazide in tuberculosis 451  
Isupren in asthma 93

## J

Jaundice in pneumococcal pneumonia 77  
complication of 89  
thiosemicarbazones causing 450  
Jungling's syndrome in sarcoidosis 185

## K

Kaolin and lung disease 261  
Kartagena syndrome 159  
radiographic appearance 297  
Katharometry in chronic hypertrophic emphysema 113  
Kidney  
bronchitic disease and 43  
myomas of in tuberculous sclerosis 201

Kidney—*continued*  
pleural effusion and disease of 14  
pneumococcal pneumonia and 74  
polyarteritis nodosa symptoms in 78  
tuberculous deposits in 301  
*Klebsiella pneumoniae* in pulmonary abscess 236

Kymography,  
harmatoma 339  
respiration and 32  
Kyphosis  
emphysema in 108, 111, 119  
tuberculosis in, 322

## L

Laryngitis  
irritant gases and 142  
tuberculosis and  
chemotherapy 443 445  
haematogenous 304  
pneumothorax artificial and 391  
thiosemicarbazones in 450

Larynx  
closure in prevention of lung abscess 21  
swabbing of in pulmonary tuberculosis 3  
Leiomyoma mediastinal 373  
Leiomyosarcoma diaphragmatic 384  
Letterer Siwe's disease 203

Leucocytosis  
eosinophilia transient pulmonary in 71  
neutrophilic in mycoses 167  
pneumococcal pneumonia in 80  
pulmonary tuberculosis in 373

Leucopenia  
abscess pulmonary in 221  
histoplasmosis in 185  
tuberculosis in  
haematogenous 295  
pulmonary 373

Leukaemia 194  
Lewisite causing acute tracheal bronchitis 53

Lipidol  
fluke infestation treatment 224  
pneumonia caused by 151

Lipoma  
diaphragmatic 384  
mediastinal 373  
Lipostol streptomycin and inhibitor of 433

Liver  
cirrhosis of in haemothorax 51  
enlargement of  
bronchiectasis in 163  
eosinophilia transient pulmonary in 71  
histoplasmosis in 185  
pneumococcal pneumonia in 74  
fatty infiltration thiosemicarbazones causing 450  
pleural effusion and disease of 13  
psittacotic pneumonia in 112  
tuberculoma deposits in, 301

Lobation abnormalities of 17 19  
 Lobectomy 426 (see also Lung resection)  
 bronchiectasis in, 183  
 hamartoma in 340  
 physiotherapy following 408

## Lobes

azygos vein of 17  
 branches of  
 anterior  
 basal of lower 15  
 of left upper 12  
 of right upper 8  
 apical  
 of lower 13  
 of right upper 8  
 apico-posterior of left upper 11  
 cardiac of lower 15  
 inferior of left upper 12  
 lateral  
 basal of lower 15  
 of right middle 8  
 lingula of left upper 12  
 medial  
 basal of lower 15  
 of right middle 11  
 posterior  
 basal of lower 5  
 of right upper 8  
 superior of left upper 12  
 upper division of left upper 11  
 left upper 11 13  
 lower 13  
 right,  
 axillary area 8  
 lateral area 8  
 middle 8 11  
 upper 3 8  
 Löffler's syndrome 65 73  
 Lymph glands  
 enlargement of streptomycin causing 435  
 histoplasmosis in 188  
 tracheo-bronchial 346  
 tuberculosis spread by 311  
 Lymphadenoma 194  
 Lymphadenopathy in blastomycosis 190  
 Lymphangioma mediastinal 367  
 Lymphangitis reticularis tuberculosa 311  
 Lymphoblastoma 366  
 Lymphogranulomatosis benigna 135  
 Lymphosarcoma 194  
 mediastinal 366

## M

Malnutrition and pneumococcal pneumonia 75  
 Manganese pneumonia from 148  
 pneumococcal 75  
 Mantoux test 252  
 bronchopneumonia of children, in 281  
 tuberculosis in  
 children, in 277  
 haematogenous 295

## Measles

bronchiectasis caused by 159  
 bronchitis acute tracheal and 51  
 pneumonia and 62  
 whooping cough following 155  
 Mediastinitis  
 acute 353 355  
 chronic 355 356  
 Mediastinotomy 355  
 Mediastinum diseases of 350 356  
 actinomycosis affecting 164  
 adenoma of 373  
 bronchogenic tumours of 362  
 chondroma of 373  
 cysts of 358 374  
 dermoids of 363  
 displacement of 350  
 dyspnoea and 321  
 empyema tuberculous in 363  
 pneumococcal pneumonia in 71  
 emphysema of 350  
 enterogenous cysts of 370 371  
 fibroma of 373  
 ganglioneuromas of 360  
 goitre intrathoracic and 371 373  
 leiomyoma of 373  
 lipomas of 373  
 lymphangoma of 367  
 lymph node enlargement in psittacotic pneumonia 112  
 lymphoblastoma 366  
 lymphoid tissue tumours of 366-367  
 lymphosarcoma 366  
 meningocoele of 373  
 myxoma of 373  
 neurilemmomas 360  
 neurofibromas of 360  
 neuromas of 360  
 osteitis fibrosa of 373  
 pericardial cysts of 367 369  
 teratomas of 363  
 thymoma of 369 370  
 tumours of 358 374  
 xanthoma of 373  
 Melana  
 mediastinal cysts in 371  
 polyarteritis nodosa in 77  
 Melanoptysis 278  
 Melanosis 277  
 Meningism in pneumococcal pneumonia 80  
 Meningitis,  
 chorion lymphocytic pneumonia of 113  
 moniliasis and 169  
 pneumonia and  
 Friedländer's bacillus complicated by 97  
 pneumococcal 74 89 90  
 torulosis resembling 171  
 tuberculosis haematogenous and, 312  
 meningocoele mediastinal 373  
 Meningo-encephalitis from torulosis 173  
 Mentality and tuberculosis 319  
 bronchogenic 333  
 pulmonary treatment of 379



- Mercury in pulmonary syphilis 237  
 Mesotheliomas of chest wall 344  
 Meteorism in pneumonia 77 80 131  
 Methyl bromide  
     pneumonia from 141  
     pulmonary oedema and 142  
 Mica and lung disease 268  
 Migraine and asthma 88  
 Mitral stenosis  
     bronchitis acute tracheal and 43 45  
     embolism in 240  
     haemosiderosis and 250 251  
     haemoptysis causing 319  
     haemothorax in 391 51  
     schistosomiasis diagnosis from 221  
 Mondaldi drainage following thoracoplasty 419  
 Moniliasis 167 171  
     causative organism 167  
     clinical features 169  
     laboratory diagnosis 170  
     mode of infection 169  
     prognosis 171  
     radiographic appearances 170  
     treatment 171  
 Moro's tuberculin test 252  
 Morphine  
     asthma and 93  
     gassing in 146  
     pneumonic pain in 130  
 Mustard gas causing acute tracheal bronchitis 53 54  
 Myalgia epidemic 342  
*Mycobacterium tuberculosis*  
     p-aminosalicylic acid in 438-441  
     culture 367  
     dihydrostreptomycin in 438  
     microscopic examination of 365  
     PAS sensitivity to 372  
     resistance to 266  
     streptomycin in 431-438  
         sensitivity to 372  
 Myocardium  
     fatty in whooping cough 156  
     pneumococcal pneumonia and 74  
 Mycosis  
     fungoides 194  
     pulmonary 162  
 Myelomas of chest wall 347  
 Myosarcoma diaphragmatic 384  
 Myxzone 449  
 Myxoma mediastinal 373

N

- Nasopharyngitis manganese causing 148  
 Necator 224  
     americanus in transient pulmonary eosinophilia 65  
*Neisseria*  
     catarrhalis  
         bronchitis in 43  
         pneumococcal pneumonia in 93

- Neisseria*—continued  
     pharyngitis  
         bronchitis, in, 43  
         pneumococcal pneumonia in 93  
 Neoarsphenamine in eosinophilia 72  
 Neoplasia pulmonary 306-341  
 Nephritis  
     empyema complicating 47  
     pneumococcal pneumonia complicating 89  
     polyarteritis nodosa diagnosis from 80  
 Neurectomy intercostal in pulmonary carcinoma 329  
 Neurilemmomas of chest wall 345  
 Neuntis peripheral in pneumococcal pneumonia 90  
 Neurofibromas 340  
 Neuromas  
     chest wall 345  
     mediastinal 360  
 Nickel carbonyl causing pulmonary oedema 141 143  
 Niemann Pick disease 198  
 Nikethamide in gassing, 146  
 Nitrogen causing pneumonia 141  
 Nitrous fumes  
     bronchitis acute tracheal from 53  
     pneumonia from 141 143  
     pulmonary oedema from 143  
 Nocardiosis 163-167  
     causative organisms 163  
     clinical features 164  
     incidence 163  
     penicillin in 167  
     prognosis 167  
     radiographic appearances 166  
     treatment 167

O

- Occlusion bronchial pulmonary abscess secondary to 236  
 Occupation and lung disease 252 293  
     aluminium in 285  
     anthracosis 277  
     argyrosiderosis 283  
     asbestos in 263-267

- bronchitis  
     byssinosis in 258  
     fenestecium 254  
     byssinosis 255 258  
     carcinoma pulmonary 258-261  
     china clay and 261  
     chromate working and 261  
     cotton causing 255  
     diatomaceous earths 268  
     emphysema  
         byssinosis in 258  
         chronic hypertrophic 116

Figures in bold type refer to Volume Two

# INDEX

Occupation and lung disease—*continued*  
 farmer's lung 254  
 French chalk 262  
 graphite 276-277  
 kaolin 261  
 melanosis 277  
 mica 268  
 organic vegetable materials 254  
 platinum asthma 252 254  
 pneumoconiosis 277 287  
 coal miner's 276-280  
 pneumonia pneumococcal 75  
 schistosomiasis 271  
 siderosis 283  
 silicates in 261 269  
 silicosis 269 276  
 silimanite 268  
 stannosis 283  
 talc 262  
 tobacco smoking and 261  
 tripoli 268  
 tuberculosis pulmonary 386  
 vanadiosis 283

Occupational therapy in pulmonary tuberculosis 378  
 Oedema pulmonary 123 135  
 acute 125  
 central nervous system lesions and 131 132  
 heart failure and 125-131  
 angioneurotic asthma and 88  
 chronic 135  
 classification 124 125  
 hydrostatic pressure relationships 123  
 hypostatic congestion in 135  
 irradiation pneumonia in 151  
 lymphatic drainage in 124  
 methylbromide causing 142  
 nitrous fumes causing 143  
 osmotic factors 123  
 phosgene causing 144  
 physiological factors 123 124  
 subacute 132 135  
 fluid balance disturbance in 134 135  
 hypertension and 132 134

Oesophagus  
 lesions of causing empyema 20  
 pouch of inhalational pneumonia and 124  
 Oesophagoscopy in mediastinal growths 358  
 Old tuberculin international standard 277  
 Oliguria complicating pneumococcal pneumonia 89  
 Oral hygiene pneumococcal pneumonia and 89  
 Ornithosis pneumonia of 62 112  
 Osmium, pneumonia from 141 149  
 Osteitis fibrosa of mediastinum 373  
 Osteoarthropathy pulmonary empyema complicated by 47  
 Osteomyelitis empyema caused by 20  
 Otitis media complicating pneumococcal pneumonia 89  
 Oxyacetylene welding and pneumonia 143

Oxygen  
 administration problems of 40-41  
 saturation  
 pneumococcal pneumonia in 81  
 reduced effects of 31  
 therapy  
 gassing in 145  
 pneumonia in 131  
 Oxyuris 224

## P

Pancoast syndrome chest wall tumours and 345  
 Pancreas in pneumococcal pneumonia 74  
 Papilloedema in emphysema 108  
 Paracoccidiosis *brasilensis* 189  
 Paragonimiasis 223-224  
 Parkinson's disease simulating manganese pneumonia 148  
 Parotitis complicating pneumococcal pneumonia 89

## PAS

absorption 439  
 albuminuria caused by 441  
 antagonists 440  
 antipyretic action 441  
 blood clotting time and 441  
 distribution 440  
 dosage 439  
 effusion pleural in 361  
 empyema tuberculous in 364 446  
 excretion 440  
 fever caused by 441  
 gastro-intestinal upsets from 440  
 history of 438  
 kidneys effect on 441  
 mode of action 440  
*Mycobacterium tuberculosis* sensitivity to 368 372  
 pharmacology 439  
 pleurisy tuberculous in 12  
 rashes caused by 441  
 resistance 441  
 streptomycin and, 441-447  
 toxicity 440  
 tuberculosis in children in 290  
*Pasteurella*  
*pestis* pneumonia of 101  
*tularensis* pneumonia of 102  
 Paul Bunnell test in pneumonia 113  
 Pel-Ebstein syndrome 367  
 Penicillin  
 abscess pulmonary in 223  
 actinomycosis, in 167  
 atelectasis in 144  
 bronchiectasis in 175  
 carcinoma pulmonary in, 330  
 nocardiosis, in 167  
 pneumonia in  
*Bacillus anthracis* 101  
 bacterial non-pneumococcal, 96

Figures in bold type refer to Volume Two  
 [ 17 ]

## Penicillin—continued

- pneumonia in—continued
  - influenza 110
  - measles and 111
  - pneumococcal 92 133
  - staphylococcal 99
- polyarteritis nodosa in 80
- post pneumonic effusion in 13
- pyopneumothorax in 232
- sub diaphragmatic hernia in 383
- suppuration pulmonary in 228
- syphilis pulmonary in 237

## Percussion

- examination in tuberculosis 323
- therapy
  - abscess pulmonary in 224

PERITONITIS 14 66 91

## Peritonitis

- artificial pneumoperitoneum complicating 399

PHLEBO-TOMY 233, 241

## Phosgene

- bronchitis acute tracheo from 53
- pneumonia and 141 143
- pulmonary oedema caused by 144

## Phrenic nerve

- lobectomy and in pulmonary tuberculosis 426

## paralysis of 375

- pulmonary tuberculosis in 396-397
- pleural tuberculosis in, 361

## Phthisis pneumonic acute lobar 105 106

## Physiotherapy 395-413

- abscess pulmonary in 223 401
- arm and shoulder movements in surgical conditions 404
- asthma in 395
- atelectasis in 401
- bronchiectasis in 401
- bronchitis in 395
- catarrhal 400
- carcinoma pulmonary and 409
- costal expansion exercises 405
- diaphragmatic breathing 397, 405
- emphysema in 395
- empyema in 412
- lobectomy and 408
- pleurisy in 400 403
- pneumolysis extra pleural and 412
- pneumonectomy and 408

## Physiotherapy—continued

- pneumonia in 400
- post operative exercises 412
- posture in 399 404
- pre operative 404
- surgical conditions in 404
- thoracoplasty and 401, 412
- thoracotomy and 408
- tuberculosis pulmonary in 401, 409
- von Pirquet test in tuberculosis of children 270

## Plasmocytomas

- chest wall 347
- mediastinal 373

## Pleura

- actinomycosis involvement in 164
- adhesions of tuberculosis and 356
- calcification of 60-64
  - aetiology 60
  - cholesterol thorax occurring in 61
  - clinical features 63
  - empyema in 63
  - incidence 60
  - pathology 61-63
  - pneumothorax spontaneous 60
  - pseudochylothorax in 61
  - radiograph c appearances 61-62
  - scleroderma in 61
  - treatment 63
  - tuberculosis haematogenous in 63
- effusion of
  - Mycobacterium tuberculosis* demonstration in 371
  - radiographic appearances of 356
  - tuberculous 355 356
- perforation of 358
  - thoracoplasty complicating 409
- polyarteritis nodosa signs in 78
- shock to in artificial pneumothorax 393
- tuberculosis of 355-364

## Pleurectomy in tuberculous empyema 364

## Pleurisy 1 17

- aetiology 1
- allergic phenomenon of 6
- amoebiasis thoracic in 227
- Bornholm disease diagnosis from 342
- bronchitis acute tracheo catarrhal complicating 45
- clinical picture 1 2
- dry 1
  - pneumococcal pneumonia and 72
- effusion and 2
- fibrino-purulent pneumococcal pneumonia and 74
- flake infestation causing 223
- idiopathic 358-360
- physiotherapy in 400 403
- pneumococcal pneumonia and complicating 86
- sero-fibrinous pneumococcal pneumonia and 74
- tuberculosis 355 5-12
  - artificial pneumothorax in 11
  - clinical course 6

Figures in bold type refer to Volume Two

*Pleurisy—continued*  
 tuberculosis—*continued*  
   complications 7  
   diuretics in 11  
   effusion associated with 355  
   fluid sampling 9  
   guinea pig inoculations and 9  
   involvement stage of 355  
   management of fluid 10  
   oxygen consumption and 39  
   PAS in 12  
   phrenic interruption in 11  
   pneumococcal pneumonia and diagnosis from 91  
   pneumoperitoneum in 11  
   prognosis 7  
   streptomycin in 12  
   thoracentesis in 9  
   treatment 9 12  
   types 5

Pleurodynia epidemic 342  
 Pleurograms in acute empyema 34 35  
 Pneumatococles 300

Pneumococcus  
   influenzal pneumonia in 109  
   pneumonia of 67 93 (*see also* Pneumonia  
   post-operative pulmonary lesions in 85  
   types 68  
   typing of 67

Pneumoconiosis 277 287  
   argyro-siderosis 283  
   baratosis, 280  
   coal miner's 277 280  
   course 279  
   heart failure in 278  
   pathology 277  
   radiographic appearances 281 282  
   thrombosis pulmonary arterial in 278  
   treatment 280  
   tuberculo-toxaemia in 278

focal emphysema in 120  
 honeycomb lungs caused by 215  
 siderosis 283  
 stannosis 283  
 vanadiosis, 283

Pneumectomy 426 (*see also* Lung Resection)  
   bronchial carcinoma and abscess pulmonary in 237  
   bronchiectasis in 183  
   hamartoma in 340  
   mycoses in 167  
   physiotherapy and 408

**Pneumonia**  
   acute specific classification, 62  
   alba 235  
   anaesthesia danger of 123  
   aspiration, 122 129  
   abscess pulmonary and, 242  
   aetiology 62  
   clinical effects, 64  
   diagnosis 128

**Pneumonia—continued**  
   aspiration—*continued*  
     diffuse 122 123  
     treatment 135  
   leukaemia in 194  
   localized, 123-129  
     abscess pulmonary relation to 128  
     aetiology 123  
     clinical picture 125  
     pathogenesis 126  
     prognosis 128  
     signs, 125  
     symptoms 125  
     treatment 129 135  
   pneumococcal pneumonia diagnosis from 91  
   radiographic appearances 124 125 126 127

*Bacillus anthracis* 101  
 bacterial 60-103  
 bronchial (*see* Broncho-pneumonia)  
 bronchitis and 144  
 caseous 105-106  
   tuberculous radiographic appearances 342

chemical 141 153  
 asphyxiants 141  
 beryllium 146-147  
 cadmium 149  
 centrally acting poisons 141  
 chromic acid 149  
 gases, 142, 144  
 irritants 141 142  
 manganese 148  
 osmium 149  
 vanadium 148 149  
 chicken pox, of 113  
 chono-meningitis lymphocytosis of 113  
 classification 60  
 definition, 60  
 deglutition 123  
 enteric fevers in 102  
 epistuberculous, 103 104  
 erythema multiforme exudativum, of 113  
 fibrinous, 67

Friedländer's bacillus 96-97  
   incidence 66  
   radiographic appearances, 237  
   glandular fever of 113  
   haemolytic streptococcal 94 96  
   incidence 66

*Haemophilus influenzae* 99-101  
 haemothorax in 51  
 hypostatic 63 122  
 inclusion body 119  
 influenza and 62  
 influenzal 62, 107 110  
   bacteriology 109  
   clinical types, 107 108  
   diagnosis, 109  
   fulminating 108  
   mortality 108  
   pathology 109

**Pneumonia—continued**

**influenzal—continued**

pneumococcal pneumonia diagnosis from 91

treatment 110

inhalation 63 123

abscess pulmonary and 242

irradiation 151 153

leukaemoid reactions in children 198

Lupiodol and 151

lipoid 149 151

lobar 67

bronchiectasis caused by 160

caseous 341 343

measles and 62 110-111

mediastinitis caused by 354

mononucleosis infection of 113

nomenclature 60

*Pasteurella pestis* 101 102

peribronchial thrush and 169

physiotherapy in 400

pneumococcal 67 93

abdomen in 80 98

abscess pulmonary and 230

complicating 87

diagnosis from 221

formation and 199

abortive 84

aetiology determination of 93

adrenal glands changes in 74

age and incidence 74

albuminuria complicating 89

alcoholism and 75

alimentary system signs in 80

allergy and pathogenesis 71

anatomical changes in 72

animal experiments and 69 73

anorexia in 76

appendicitis diagnosis from 90

arteriosclerotic changes in 80

arthritis and 74 89

aspiration pneumonia diagnosis from 91

atelectasis and 71 91

atypical 84

aureomycin in 92

auricular fibrillation in 79

bacteraemia prognosis and 92

basic slag and 75

blood

examination in 80

pressure in 79

bronchiectasis complicating 88

bronchoscopic aspiration 71

carbon dioxide

blood content 81

inhalations in 71

cardiovascular system signs in 79

carnification in 88

catarrhal infections and 75

central 85

cerebrospinal fluid examination 89

chest signs in 78 79

chloramphenicol in 92

**Pneumonia—continued**

**pneumococcal—continued**

chloride

blood content in 81

excretion 80

clinical picture 84-85

complications 86-90

conjunctivitis complicating 89

consolidation signs of in 78

conveyance of 75

cough in 77

cylindruria in 89

delirium tremens in 77

differential diagnosis 90-92

disorientation in 77

distribution 71

double 85

dyspnoea 77

effusions complicating 86

pericardial 88

empyema complicating 85 86

endocarditis and 74 88

engorgement lobar in 72

exhaustion and 75

expectoration 77

exposure in 75

fastigium 76

febrile 80

gastro intestinal symptoms in 77

geographical distribution 74

grey hepatization in 73

haematuria in 89

haemoptysis in 77

headache in 76 77

heart signs in 79

hepatic swelling in 74

herpes in 78

immunity from 69

natural 75

incidence 66

infancy in 84

infarction pulmonary diagnosis from 90

influenzal pneumonia diagnosis from 91

insomnia 77

jaundice in 77 89

laboratory investigations in 80-83

leucocyte count prognosis and 92

leucocytosis in 80

lobar involvement 72 78

malnutrition and 75

manganese smelting and 75

massive 85

meningism in 80

meningitis and 74 88 90

meteorism in 77 80

migratory 85

morbid anatomy 71 74

mortality figures 92

myocardial changes in 74

nephritis complicating 89

nervous symptoms 77 80

neutrit peripheral in 90

nitrogen retention in 89

*Figures in bold type refer to Volume Two*

## Pneumonia—continued

- pneumococcal—continued
  - occupation and 75
  - oliguria in, 89
  - onset 76
  - oral hygiene and 89
  - otitis media complicating 85 89
  - oxygen saturation in 81
  - pain in 77
  - pancreatic changes in 74
  - parotitis, acute complicating 89
  - pathogenesis of 69
  - penicillin in 92
  - pericarditis and 74 88 91
  - peritonitis and 74 89
  - personal factors in 75
  - physical signs 78
  - pleurisy in 72, 74, 86 91
  - pneumoconiosis complicating 88
  - prognosis 92 93
  - "pseudo-crisis" 76
  - pyrexia in 76
  - race incidence 74
  - radiographic appearances 82 83
  - red hepatization in 72
  - relapse 85
  - renal changes in 74
  - resolution 73
  - delayed 87
  - respiratory catarrhal infections and 75
  - rigor in 76
  - scarlet fever diagnosis from 90
  - seasonal factors 74
  - serum therapy 92
  - sex incidence 74
  - social conditions and 74
  - splenic enlargement in 74
  - Streptococcus haemolyticus* in 85
  - sulphonamides in 92
  - susceptibility to factors of 74 76
  - symptoms 76-78
  - tachypnoea in 78
  - terminal 85
  - termination 77
  - thrombophlebitis in 90
  - trauma role of 75
  - tuberculosis pulmonary complicating 89
  - typhoid fever diagnosis from 90
  - typhus fever diagnosis from 90
  - urine in 80
  - vasomotor peripheral disturbances in, 80
  - virus pneumonia diagnosis from 91
- pneumonotropic viral 113 119
- post-operative 63
- primary atypical 113 114
- pseudo-lobar 236
- psittacosis-ornithosis group 62 65
- psittacosis 111 113
- rickettsial 62, 65 119 122
- secondary 63
- smallpox of 113
- staphylococcal 97 99
- abscess, pulmonary and, 229 230 234

## Pneumonia—continued

- staphylococcal—continued
    - bacteriology 97
    - children in 232
    - clinical picture, 98
    - empyema acute caused by 18
    - incidence 66 97
    - mortality 99
    - pathogenesis 97
    - pathology 98
    - prognosis 99
    - radiographic appearances 235
    - signs 98 99
    - symptoms 98
    - treatment 99
  - Stevens Johnson syndrome of 113
  - streptococcal pulmonary abscess and 199
  - 230
  - suppurative pulmonary abscess in 202
  - non-specific lung abscess and 239
  - terminal 63
  - treatment 129 136
  - aureomycin 133
  - cardiovascular system in 131
  - chloramphenicol 133
  - convalescence and 135
  - cough 131
  - dangers of 134
  - diet in 130
  - digitalis control of 131
  - general 129
  - insomnia relief of 130
  - meteorism 131
  - nursing 129
  - oxygen administration 131
  - pain relief of 130
  - penicillin 133
  - specific 132 135
  - sulphonamides 133
  - tetracycline 133
  - tuberculous
    - acute 103 107
    - lobar 105 106
  - tularaemic 102
  - undulant fever in 103
  - verminous schistosomal 217
  - viral 111 122
  - pneumococcal diagnosis from 91
- Pneumonitis**
- abscess, pulmonary in 202 219 221
  - beryllium pneumonia diagnosis from 146
  - coccidiodomycosis in 184
  - definition 60
  - suppurative
    - abscess pulmonary and 242
    - carcinoma pulmonary complicated by 311
    - nocardiosis in 167
    - sub-diaphragmatic abscess complicated by 380
  - pneumoconiosis complicating pneumococcal pneumonia, 83

# INDEX

## Pneumoperitoneum

- artificial 397 399
- diaphragmatic eventration in 376
- mediastinum growths of diagnosis of 358
- pleural tuberculosis in 361
- pneumothorax spontaneous complicating 54
- spontaneous
  - emphysema chronic hypertrophic in 117
  - pleurisy tuberculous in 11
  - sub-diaphragmatic abscess causing 380
- Pneumothorax
  - artificial
    - adhesions revealed by 358
    - age in 391
    - air embolism complicating 393
    - anaesthesia 392 392
    - apparatus for 392
    - asthma and 391
    - atelectasis and 394
    - complication 393 394
    - contra indications 390
    - effusion pleural and 395
    - emphysema chronic hypertrophic in 117
    - emphysema tuberculous and 395
    - extrapleural 399-403
    - fistula broncho pleural and 396
    - haemothorax complicating 393 49
    - heart disease and 391
    - hydatid cysts in 233
    - indications 390
    - induction 392
    - management 394
    - medastinal growths diagnosis of 358
    - needles for 392
    - pleurisy tuberculous in 11
    - position for 392
    - pregnancy and 391
    - radiographic appearances 350
    - refilling 392
    - site 392
    - tracheo bronchitis and 349
    - traumatic complicating 393
    - tuberculosis in
      - bronchogenic 344
      - haematogenous 312
      - pleural 358
  - spontaneous 54-59
    - abscess and
      - pulmonary causing 54
      - sub phrenic diagnosis from 57
    - aetiology 54 56
    - aluminium exposure in 285, 286
    - artificial complicating 393
    - aspiration biopsy causing 324
    - asthma complicated by 89
    - atelectasis complicated by 148
    - benign 54
    - carcinoma pulmonary causing 54
    - clinical symptoms 56
    - cysts and
      - bronchial diagnosis from 57
      - pulmonary causing 304

## Pneumothorax—continued

- spontaneous—continued
  - cystic lung localized causing 55
  - diagnosis 57
  - emphysema chronic hypertrophic and 55, 108
- empyema and
  - acute 18
  - tuberculous 363
  - honey comb lungs complicated by 209
  - newborn in 55
  - oesophagocopy causing 54
  - pleura and
    - calcification of 60
    - perforation 147 358 359
  - pneumoperitoneum artificial compli
    - cated by 54
  - prognosis 58
  - radiographic appearance 56
  - tension 58
  - tracheotomy causing 54
  - traumatic 54
  - treatment 58
  - thyroidectomy causing 54
- Polomyelitis whooping cough immunization and 160
- Polyarteritis nodosa 74 81
- aetiology 75
- antihistamines in 80
- arteritis temporal diagnosis from 80
- brucellosis diagnosis from 80
- cardiac symptoms 78
- clinical manifestations 77-78
- definition 74
- diagnosis 78-80
- endocarditis subacute bacterial diagnosis
  - from 78
- eosinophil a transient pulmonary and 65
- epigastric discomfort in 77
- haematemesis in 77
- history 74
- hypertension diagnosis from 80
- locomotor signs 78
- melaena in 77
- morbid anatomy 74 75
- nephritis diagnosis from 80
- neurological symptoms 78
- penicillin in 80
- prognosis 80
- pulmonary signs in 78
- renal symptoms 78
- salicylates in 80
- sarcoidosis diagnosis from 78
- sections 75, 76
- sulphonamides in 80
- treatment 80
- tuberculosis diagnosis from 78
- Postural drainage
  - foreign bodies intrabronchial in 218
  - whooping cough in 159
- Pottenger's sign in mediastinitis 355
- Precipitins pneumococcal pneumonia and 69

- Pregnancy  
 artificial pneumothorax and 391  
 tuberculosis pulmonary and, 381
- Premenstrual temperature in pulmonary tuberculosis 318
- Promin in tuberculosis 447 448
- Promizole in tuberculosis 447 448
- Promithrombin levels  
 embolism in 244  
 irradiation pneumonia in 152
- Prurigo and asthma 88
- Pseudochylothorax in pleural calcification 61
- Pseudo-tuberculosis 185
- Psittacosis  
 pneumonia of 62, 111 113  
 tuberculosis pulmonary and 381
- Purified protein derivatives 252 270  
 children in 277
- Pyæmia  
 acute empyema complicating 37  
 staphylococcal pneumonia and 98
- Pyopneumothorax 54  
 children in 232  
 Friedländer's bacillus pneumonia in 97  
 influenza pneumonia in 109  
 radiological appearances 233 19  
 staphylococcal pneumonia in 98  
 sub-diaphragmatic abscess complicated by 380
- Pyothorax in staphylococcal pneumonia 99
- R**
- Racial factors,  
 asthma in 84  
 haematogenous tuberculosis in 295  
 pneumococcal pneumonia in, 74  
 sarcoidosis in 190
- Radioactive isotopes in intrathoracic goitre 371
- Radiography  
 diagnostic (see Radiographic appearances)  
 mass, 254-255
- Radiographic appearances  
 abscess pulmonary 203 205 207 209 210  
 215-218 219 220 226 227 301  
 actinomycosis 165 166  
 adenoma bronchial 333, 335  
 aluminium exposure 287  
 amoebiasis thoracic 227  
 asbestosis exposure 266  
 aspergillosis 176  
 atelectasis 144 145 148, 149 153  
 post-operative 239  
 bagasse exposure 255 256  
 beryllium sarcoi 147  
 blast injuries 246  
 blastomycosis, North American 189  
 Bornholm disease 342  
 bronchoectasis 347 161, 163 164 169 171  
 172, 174 181  
 bronchitis and 46 47
- Radiographic appearances—*continued*  
 bronchitis  
 acute tracheal 44 46 47 50  
 chronic 57  
 broncho-pneumonia  
 children in 281 282 283  
 tuberculous 107  
 byssinosis 259  
 carcinoma  
 bronchial 322  
 pulmonary 260 318  
 collapse due to 323  
 emphysema in 321  
 hilar 314 315  
 left lower lobe 313  
 peripheral 317 323 327  
 prostatic metastasis 332  
 sub-hilar 319  
 carcinomatous lymphangiosis 313  
 catheterization 47  
 chondroma of chest wall 346  
 coccidioidomycosis 182 183  
 consolidation pulmonary 9 10 14  
 cysts  
 hydatid 230 232 233  
 mediastinal pericardial coelomic 368  
 mesothelial diaphragmatic 385  
 pulmonary 299 301 303  
 congenital 296 297  
 dermoids mediastinal 364 365  
 diaphragm humping of in hepatic amoebiasis 226  
 dysplasia fibrous 348  
 effusion,  
 encysted 362  
 pleural 359 3 4  
 tuberculous, 356  
 embolism pulmonary 242  
 emphysema  
 chronic hypertrophic 111 112  
 mediastinal 352  
 empyema  
 acute 18, 22, 24 26  
 chronic 40 46  
 eosinophilia transient pulmonary 67 69  
 epituberculosis 352  
 eventration of diaphragm 378 379  
 farmer's lung 178 254  
 fluke infestation 224  
 foreign body ex, pulmonary abscess secondary to 242  
 goitre intrathoracic 371  
 granuloma  
 diffuse cystic, 208, 211 212  
 eosinophilic, 206  
 haemosiderosis  
 idiopathic 248 249  
 mitral stenosis and 250  
 haemothorax 40 41  
 hamartoma  
 chondromatous 337  
 vascular 338 339  
 Hand-Schüller-Christian disease 206

Figures in bold type refer to Volume Two



Radiographic appearances—continued

- herna
  - diaphragmatic 381, 383
  - mediastinal 351
- histoplasmosis 186-187
- Hodgkin's disease 367
- honey comb lungs 201 202 204 206, 208 214
- infarction pulmonary, 240 241
- Kartagena's syndrome 297
- leukaemia 197 198
- lobectomy 422
- lymphadenoma 193, 195
- lymphosarcoma 196
- mediastinitis 356
- mica exposure 268
- moniliasis 170
- myeloma of chest wall 348
- neurofibroma 361
- nitrous fumes gassing in 54
- nocardiosis 166
- oedema pulmonary 144
  - acute 129
  - heart failure and 129 130
  - sub-acute hypertension and 133
- paralysis diaphragmatic 377
- pleura calcification of 61-62
- pleurisy 18
  - dry 2
- pneumoconiosis coalminers 281 282
- pneumonecrosis 425
- pneumonia 14
  - aspiration 124-127
  - beryllium 147
  - caseous tuberculous 342
  - epituberculous 103
  - Friedlander's bacillus 97 237
  - irradiation 151
  - lipoid 150
  - pneumococcal 82
  - ricketsial 120-121
  - staphylococcal 98 235
  - tuberculous lobar 106
  - viral 116-117
- pneumoperitoneum 421
- pneumothorax
  - artificial 350 416 421
  - extrapleural 400-401
  - spontaneous 359 56-57
- polyarteritis nodosa 78
- pyopneumothorax 233 19
- round worm infection 224
- sarcoidosis 78 188-189
- sarcoma
  - cannon ball 332
  - embolic 240
- schistosomiasis
  - cardiac 219 220
  - pulmonary 218 219
- siderosis 284
- silicosis 272 276, 284
- silimanite exposure 268
- syphilis pulmonary 235 236

Radiographic appearances—continued

- talc exposure 263
- teratomas mediastinal 364
- thoracoplasty 405-407, 411 418
  - bilateral 409
  - lateral 425
- thymoma mediastinal 370
- thyroid petrosternal 372
- torulosis 172
- tuberculosis
  - bronchogenic 335 338 340
  - fibro-caseous 345
  - haematogenous 297 299 306-310
  - lymphogenous 311
  - minimal lesions 337
  - pulmonary 38 411
    - early subclavicular infiltrate 328 330
    - emphysematous chronic diffuse 304
    - nodular 326 328 330
    - primary lesion in children 280-281
  - tubercous sclerosis 204 205
  - whooping cough 157
- Radiotherapy
  - carcinoma pulmonary in 329
  - Ewing's tumour, in 349
  - lymphadenoma in 194
  - lymphosarcoma in 194
  - sarcoma pulmonary in 330
  - von Recklinghausen's disease 360
- Rectum prolapse of whooping cough complicating 158
- Relaxation in physiotherapy 396
- Renal function and chemotherapy 372
- Resection lung chemotherapy in 446-447
- Respiration
  - alveoli and 23
  - gas exchange in 26-29
  - anoxia and 31
  - aortic reflexes regulation and 30
  - applied physiology of 23-42
  - bronchial tree and 23
  - carbon dioxide
    - administration problems of 40-41
    - blood transport 26-27
    - respiratory centre action on 30
  - carotid reflexes regulation and 30
  - catarrhal infections of, pneumococcal pneumonia and 75
  - cerebral cortex and regulation 29
  - clinical application 31-40
  - dead space 37
  - dynamics of 35
  - intrapulmonary mixing 37
  - kymography in 32
  - lobule lung schematic representation 24
  - lymphatic drainage and 24
  - muscles of aspiration pneumonia in 65
  - nervous control and 29
  - oxygen
    - administration problems of 40-41
    - blood transport of 27-29
    - consumption 38
    - pain and 30

# INDEX

Respiration—*continued*  
 regulation of 29 31  
 residual air 34  
 spirometry in 28 32  
 total lung capacity 34  
 tracheo-bronchitis obstructive and 348 349  
 upper respiratory tract and 73  
 vascular shunts and 31  
 ventilation pulmonary mechanism of 31  
 vital capacity 33

Reticulo-endothelial system and lung disease 192 199

Reticuloses 192 199  
 Gaucher's disease 194  
 general considerations 192  
 glandular enlargement 192  
 haemothorax in 51  
 leukaemias 194  
 leukaemoid reactions 198  
 lymphadenoma 194  
 lymphosarcoma 194  
 mediastinum in 367  
 mycosis fungoides 194  
 Niemann Pick disease 198  
 radiographic appearances 193, 195 198  
 Rhabdomyosarcoma diaphragmatic 384  
 Rheumatism bronchiectasis complicated by 173

Rhinitis allergic and asthma 87

Rhinorrhoea and asthma 87

Rib

fixation in tuberculosis 322

resection in empyema 28 32

Rickettsia pneumonia of 62 119 120

Robert's flap operation in empyema 45

Round worm infections 224 225

## S

Salicylates in polyarteritis nodosa 80  
 Sanatorium treatment 378  
 Sanocrysin in pulmonary tuberculosis 383  
 Sarcoidosis 185 191, 330 331  
 aetiology 186 187  
 aspiration hepatic in 187  
 beryllium 147  
 granulomatosis diagnosis from 190  
 biopsy in diagnosis 187  
 caliciferol in 191  
 definition 185  
 diabetes insipidus in 185  
 diagnosis 187  
 diaphragmatic 384

osteogenic of chest wall 345

Sarcoidosis—*continued*  
 polyarteritis nodosa and diagnosis from 78  
 racial factors in 190  
 radiographic appearances 188 189  
 treatment 191  
 urinary beryllium content in 190

Scarlet fever  
 haemothorax in 51  
 pneumococcal pneumonia simulating 90

Schaumann's disease 185

Schede's operation in chronic empyema 45

Schistosomiasis 271

Schistosomiasis, 216-222

cardiac 219-221

pulmonary 216-220

Scleroderma

periarthritis nodosa histological similarity to 75

pleural calcification in 61

pulmonary fibrosis in 61

Sclerosis tuberosa 201 203

Scoliosis

thoracoplasty prevention of following 414  
 tuberculosis in 322

Sedatives in whooping cough 159

Serological tests in pulmonary tuberculosis 373

Siderosis 283

Silicates and lung disease 261 269

Silicosis 269 276

Simon focus 300

bronchogenic tuberculosis in 336

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

— — — — — " " " "

Social factors pneumococcal pneumonia and 74

Sodium Amytal in insomnia of pneumonia 130

Sodium Luminal in pneumonic delirium 131

Spine tuberculous abscess of pleural effusion and 358

Spirometry  
 respiration and 32  
 total lung capacity subdivisions of 28

Spleen  
 enlargement of  
 eosinophilia transient pulmonary in, 71  
 haemistogenous tuberculosis in 295  
 histoplasmosis in 185  
 pneumococcal pneumonia in 74  
 rickettsial pneumonia in 120  
 putrefactive pneumonia in 112  
 tuberculoma deposits in 301

Sputum  
 bronchiectasis in 162  
 bronchitis in 43  
 carcinoma pulmonary in 320  
 monilia in 170  
 pneumonia in  
 bacterial 95  
 Friedländer's bacillus 96  
 pneumococcal 77

# INDEX

- Sputum—continued**  
pneumonia in—*continued*  
staphylococcal 98  
tuberculous lobar 106  
tracheo bronchitis in 348  
tuberculosis in  
bronchogenic c 343  
pulmonary 320 369
- Sporotrichosis** 190-192  
causative organism 190-192  
*Sporotrichum schenckii* 190
- Stannosis** 283
- Staphylococcus aureus**  
haemothorax in 49  
lung abscess in 211  
pneumonia in 93 97 98
- Status asthmaticus** 94 95
- Stevens-Johnson syndrome** pneumonia and, 113
- Subophen in schistosomiasis** 221
- Sulboestrol in pulmonary oedema** 135
- Streptococcus**  
haemolyticus  
bronchitis in 43  
empyema in 20  
pneumonia in 93  
pneumococcal 85  
mucosus 67  
pneumoniae 67  
viridans in bronchitis 43
- Streptodornase and streptokinase** 36
- Streptomyces griseus** 431
- Streptomycin**  
abscess pulmonary in 223  
absorption 433  
albuminuria in 436  
antagonists 433  
bronchiectasis in 181  
cavity drainage and 415  
cysteine action on 433  
dermatitis from 436  
distribution 433  
effusion pleural in 361  
empyema tuberculous in 364 446  
excretion 433  
Friedländer's bacillus pneumonia in 97  
history 431  
Liposomal action on 433  
mode of act on 433  
*Mycobacterium tuberculosis* sensitivity to, 368 372  
oral administration 432  
PAS and 441-447  
*Pasteurella pestis* pneumonia in 102  
pharmacology 432  
pleurisy tuberculous in 12  
pregnancy and 434  
resection in 426  
resistance 436  
sulphones and 438  
sub-diaphragmatic hernia in 383  
toxicity 433  
tuberculin and in pleural effusion 363
- Streptomycin—continued**  
tuberculosis in  
bronchogenic 431-438  
haematogenous 312  
iliary resistance in 437  
pulmonary  
children in 290 432  
dosage 432  
urinary changes and 436  
vestibular disturbances from 434  
vitamin deficiencies from 436  
whooping cough in 139
- Strongyloides** 224
- Strophanthin in pulmonary embolism** 245
- Sulphadiazine in nocardiosis** 167
- Sulphamezathine**  
*Pasteurella pestis* pneumonia in 102  
staphylococcal pneumonia in 99
- Sulphetone in pulmonary tuberculosis**, 447 448
- Sulphonamides**  
abscess pulmonary in 228  
actinomycosis in 167  
blastomycosis in 190  
bronchiectasis in 175  
bronchitis acute tracheal catarrhal in 48  
eosinophilia transient pulmonary in 65  
pneumonia in 133  
bacterial non pneumococcal in 95  
pneumococcal 91  
staphylococcal 99  
polyarteritis nodosa in 80  
sub-diaphragmatic hernia in 383  
torulosis in 173
- Sulphones**  
antagonists of 448  
history 447  
pharmacology 448  
results of treatment 449  
streptomycin resistance and 438  
toxicity 442
- Sulphur dioxide causing pneumonia** 141 142
- Syphilis pulmonary** 235-237
- T
- Tachycardia**  
hydatid cysts in 231  
polyarteritis nodosa in 78
- Tachypnoea in pneumococcal pneumonia** 78
- Taenia echinococcus** 229
- Talc and lung disease** 262
- Teeth carious actinomycosis and** 163
- Teratomas mediastinal**, 363
- Terramycin**  
amoebic infections in 227  
pneumonia in 133  
subdiaphragmatic hernia in 384  
whooping cough in 159
- Theophylline in pulmonary oedema** 131
- Thiacetazone** 449
- Thioparazone** 449
- Thiosemicarbazones** 450
- Thoracentesis in tuberculous pleurisy**, 9



Tuberculosis—*continued*bronchogenic—*continued*

- stabilization of disease 337
- subclavicular nodular early infiltrate 326
- superinfection exogenous 332 334
- symptoms of 327
- treatment indications 338

## carriers 262

## cattle contacts 266

## chemotherapy in 431-456

## p-aminosalicylic acid 438-441

## dihydrostreptomycin 438

## nicotinaminic analogues 451

## streptomycin 431-438

## sulphones 447-449

## thiosemicarbazones 449-451

## children in 270-293

## adenitis hilar complicating 284

## adult type infiltration 284

## B C G in 291

## broncho pneumonia and 280-284

## bronchoscopy in 291

## disseminated lesions 290

## incidence 270

## Mantoux test 277

## miliary 284

## percutaneous test in 277

## pleural effusion in 289

## prevention 291 293

## primary lesions 277

## clinical manifestations 279

## course 278

## glandular involvement 278

## initial reaction 278

## management 290-291

## PAS in 290

## pathology 278

## radiographic evidence 279

## situation 277

## streptomycin 290

## treatment 290-291

## purified protein derivatives in 277

## tuberculin tests 270

## comparisons in 277

## von Pirquet test in 270

## contacts 262

## crude death rates 247

## D D S in 447 448

## dasone in 448

## droplet infection 260

## employment and 259

## epidemiology 245 273

## extrapleural chemotherapy in 447

## factory contacts 266

## fibrocaceous 344-346

## fibroid 346

## general surveys 255

## genito-urinary 303

## haematogenous 294 313

## abortive forms of 300

## acute miliary 295 296

## Assmann focus of 301

## calcification in 299 163

Tuberculosis—*continued*haematogenous—*continued*

## chronic miliary 296-300

## collapse therapy in 312

## extra pulmonary foci lung changes in 302 303

## genito urinary manifestations 312

## ground glass appearance radiographic 306-307

## management 312

## serous membranes implication 312

## Simor focus in 300

## streptomycin in 312

## suprenals involvement in 312

## treatment 311 313

## vaccination in 312

## hospital contacts 264

## husband wife relationship 263

## immunology 345 373

## industry in 258

## institutional contacts 265

## lymphogenous 311

## meningeal thiosemicarbazones in 450

## miliary

## chemotherapy in 442

## children in 284

## honey comb lungs caused by 215

## leukaemoid reactions in 198

## thiosemicarbazones in 450

## mortality 249 247 251 260

## natural resistance 268

## nicotinaminic analogues 451

## non-caseating 185

## notification 251

## nurses and 264

## open air work and 259

## phagocytes in resistance 268

## pleural 255 264

## aetiology 355

## age groups in 356

## bronchial block complicating 362

## bronchogenic lesions relation to 356

## children and 356

## clinical features 360-361

## discovery of 360

## effusion in 355 364

## erythema nodosum and 360

## glands tuberculous complicating 362

## phrenic interruption 361

## pneumoperitoneum in 361

## pneumothorax artificial in 358

## post primary lesions relation to 356

## prognosis 361

## radiographic appearances 356 357

## stage of involvement 355

Figures in bold type refer to Volume Two

- Tuberculosis—continued**  
**pleural—continued**  
 tracheo-bronchitis complicating 362  
 treatment 361 362  
 portals of entry 260  
 primary,  
   chemotherapy in 442  
   pulmonary embolism caused by 238  
 Promin in, 447 448  
 Promizole in 447, 448  
 prophylaxis 270 273  
 proportional mortality 248  
 pulmonary  
   abortion therapeutic and 382  
   actinomycosis simulating 164  
   agglutination tests 373  
   amenorrhoea in 318  
   animal inoculation in 368  
   asthma complicated by 89  
   atelectasis in 321  
   auscultation in 323  
   B.C.G. vaccination in children 382  
   biopsy in 373  
   blood picture 373  
   breast feeding in 382  
   breath sounds in 323  
   bronchiectasis and 319  
   cavitation and 319  
     abscess and 222  
   cerebrospinal fluid examination in 371  
   chemotherapy 442-445  
     tests before 371 373  
   chest pain in 321  
   clinical pathology 365 375  
   complement fixation tests, 373  
   cough in 320  
   diabetes and 382  
   dyspnoea in 321  
   emphysematous chronic diffuse 303 311  
   erythrocyte sedimentation rates, 373 374  
   excretions microscopic examination, 365  
   exudates microscopic examinations 365  
   faeces in 370  
   fibrosis pulmonary in 321  
   fluke infestation simulating 223  
   gastric lavage in 370  
   haemoptysis in 319  
   haemothorax in 51  
   heart beats in 322  
   histology 373  
   histoplasmosis and 186  
   hypochromic anaemia in 373  
   inspection in 322  
   kyphosis and 322  
   laryngeal swabbing in 370  
   leucocyte count in 373  
   local effects symptoms of 316 319  
   mediastinal displacement and 321  
   mentality and 319  
   mitral stenosis causing haemoptysis, 319  
   monocyte count in 374  
   *Mycobacterium tuberculosis* culture of 367  
   onset 315 324
- Tuberculosis—continued**  
**pulmonary—continued**  
 original discovery 315  
 palpation 322  
 PAS sensitivity and 368  
 percussion in 323  
 physical signs 321  
 physiotherapy in 401, 409  
 pleural fluid in 323 371  
 pneumococcal pneumonia complicating, 87  
 pregnancy and 381  
 puerperal relapse in 381  
 pulse rate in 318  
 radiographic appearances 318  
 rales in 324  
 reticulosis in 194  
 rhonchi in 323  
 rib fixation in 322  
 scoliosis in 322  
 serology 373  
 shadows discovery of 316  
 silicosis and 272  
 sputum in 320 369  
 stomach washing in 370  
 streptomycin sensitivity and 368  
 temperature in 317  
 torulosis and 172  
 toxæmia symptoms of 316 317  
 treatment 376-430  
   active relaxation 389  
   after-care 384  
   antipyretics in 383  
   anxiety and 386  
   atelectasis in 390  
   cavitated lesions 387 389  
   cavity drainage 415-419  
   chemotherapy 431-456  
   climate in 377  
   cod liver oil in 384  
   collapse therapy 397 396  
   cough in 383  
   diaphragmatic paralysis 396 397  
   diarrhoea in 384  
   domiciliary 379  
   dyspepsia 384  
   exercise in 378  
   fever 383  
   fresh air in 377  
   gold salts in 383  
   haemoptysis 383  
   and gestation 384  
   insulin in 384  
   lesure and 387  
   lobectomy 426  
   mentality and 379  
   occupational therapy in 378  
   occupations and 386  
   passive relaxation in 389  
   pneumonecstomy 426  
   pneumoperitoneum, artificial 397 399  
   pneumothorax artificial 380-396  
     extrapleural 399-401

**Tuberculosis—continued**  
**pulmonary—continued**  
**treatment—continued**  
 posture in 387  
 resection in 419-427  
 responsibility and 386  
 rest in 376-377  
 sanatorium 378  
 Sanocrysin in 383  
 specific 382 383  
 symptomatic 383 384  
 thoracoplasty 403-413  
 travelling and 385  
 tuberculin 382-383  
 tumours causing haemoptysis 319  
 ulcers and 319  
 urine examination in 370  
 ventilation and 385  
 village settlements 378  
 vitamin deficiency in 384  
 vocal fremitus absence of 323  
 weight loss in 384  
 wheezing in 321  
 working conditions and 385  
 Ziehl Nielsen staining 365  
 renal polyarteritis diagnosis from 78  
 resistance to 266-270  
 sarcoidosis diagnosis from 187  
 skeletal 303  
 standardized death rate 248  
 students and 265  
 Sulphetron 447 448  
 surgical 302  
 total deaths 247  
 tracheo bronchial chemotherapy in 444  
 tuberculin surveys 252  
 village settlements 263  
 war time and 259  
 world prevalence of 245  
*Tuberculo-toracemia in pneumococcosis* 278  
 Tularaemia pneumonia of 102  
 Tumours (*see also* specific neoplasms)  
 bronchial  
     adenoma 331 335  
     fibro-sarcoma 330  
 chest wall 344 349  
 diaphragmatic 384-386  
 effusion pleural and 360 14  
 mediastinal 358 374  
 neurogenic 340  
 non malignant physiotherapy and 404  
 pulmonary 306-341

U

Ulcer of tongue in whooping cough 158  
 Undulant fever in pneumonia 103  
 Upper respiratory tract respiration and 23  
 Urine examination  
     pneumococcal pneumonia in 80  
     tuberculosis pulmonary in 370  
     vanadium demonstration of 149

V

Vaccination (*see also* specific vaccines)  
     haematogenous tuberculosis in 312  
     whooping cough in 160  
 Vaccines skin testing  
     blastomycosis in 190  
     moniliasis in 171  
 Venesection  
     embolism in 245  
     mediastinitis in 356  
 Village settlements 263 378  
 Vital capacity 33  
     asthma in 88  
     emphysema, in 112  
     oedema pulmonary in 128 129  
     silicosis in 272  
 Vitamin

W

Wasserman test in pulmonary syphilis 237  
 Weinberg complement fixation test in hydatid cysts 229  
 Westergren erythrocyte estimation method 375  
 Whooping-cough 154-161  
     adenitis hilar simulating 285  
     aetiology 154  
     altitude treatment 159  
     antispasmodics in 159  
     atelectasis in 141  
     atropine in 159  
     aureomycin in 159  
     bacteriology 154  
     belladonna in 159  
     blood count in 158  
     bromoform in 159  
     bronchiectasis caused by 159  
     bronchitis acute tracheal and 51  
     catarrhal stage 156  
     Chloromycetin in 158  
     clinical course 156-158  
     complications 158 159  
     death rates 154  
     decompression treatment 159  
     ephedrine in 159  
     epidemiology 154  
     Eumydrin in 159  
     general measures 158  
     immunization 160  
     incidence 155

Figures in bold type refer to Volume Two

# INDEX

Whooping-cough—*continued*  
 incubation period, 155  
 isolation in 160  
 laboratory findings 158  
 leukaemoid reactions in 158  
 notification rate 154  
 paroxysmal stage 156  
 pathology 155  
 phenobarbitone in 159  
 pneumothorax complicating 157  
 prevention 160  
 radiographic appearances 157 158  
 sedatives in 159  
 streptomycin in 159

Whooping-cough—*continued*  
 terramycin in 159  
 treatment 158 159  
 Wintrobe erythrocyte estimation method, 374  
 Wool-sorter's disease 101

## X

Xanthoma medullary 373  
 Xanthomatosis 203-207  
 X rays  
     therapy (see Radiotherapy)  
     diagnosis (see Radiographic appearances)

Figures in bold type refer to Volume Two





